




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : C07D 207/34, 231/14, 233/90, 249/06, 257/04, A61K 31/40, 31/41, C07D 401/12, 403/12, 409/12, 413/12</p>	<p>A1</p>	<p>(11) International Publication Number: <b>WO 98/28269</b> (43) International Publication Date: 2 July 1998 (02.07.98)</p>
<p>(21) International Application Number: PCT/US97/22895 (22) International Filing Date: 15 December 1997 (15.12.97) (30) Priority Data: 08/769,859 23 December 1996 (23.12.96) US 08/879,944 20 June 1997 (20.06.97) US (71) Applicant: THE DUPONT MERCK PHARMACEUTICAL COMPANY [US/US]; 1007 Market Street, Wilmington, DE 19898 (US). (72) Inventors: PINTO, Donald, Joseph, Phillip; 39 Whitson Road, Newark, DE 19702 (US). PRUITT, James, Russell; 237 Skycrest Drive, Landenberg, PA 19350 (US). CACCIOLA, Joseph; 105 Pattie Drive, Newark, DE 19702 (US). FEVIG, John, Matthew; 291 Church Road, Lincoln University, PA 19352 (US). HAN, Qi; 2609 Marhill Drive, Wilmington, DE 19810 (US). ORWAT, Michael, James; 118 South Colts Neck Way, Hockessin, DE 19707 (US). QUAN, Mimi, Lifen; 113 Venus Drive, Newark, DE 19711 (US). ROSSI, Karen, Anita; 120A Emery Court, Newark, DE 19711 (US).</p>		<p>(74) Agent: VANCE, David, H.; The Dupont Merck Pharmaceutical Company, Legal/Patent Records Center, 1007 Market Street, Wilmington, DE 19898 (US). (81) Designated States: AM, AU, AZ, BR, BY, CA, CN, CZ, EE, HU, IL, JP, KG, KR, KZ, LT, LV, MD, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, UA, VN, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: NITROGEN CONTAINING HETEROAROMATICS AS FACTOR Xa INHIBITORS</p>		
<div style="text-align: center;">  <p>(I)</p> </div>		
<p>(57) Abstract</p> <p>The present application describes nitrogen containing heteroaromatics and derivatives thereof of formula (I) or pharmaceutically acceptable salt or prodrug forms thereof, wherein J is N or NH and D may be C(=NH)NH<sub>2</sub>, which are useful as inhibitors of factor Xa.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		



TITLE


Nitrogen Containing Heteroaromatics as Factor Xa Inhibitors

FIELD OF THE INVENTION

This invention relates generally to nitrogen containing heteroaromatics which are inhibitors of trypsin-like serine protease enzymes, especially factor Xa, pharmaceutical compositions containing the same, and methods of using the same as anticoagulant agents for treatment and prevention of thromboembolic disorders.


BACKGROUND OF THE INVENTION

WO 95/18111 addresses fibrinogen receptor antagonists, containing basic and acidic termini, of the formula:




wherein R<sup>1</sup> represents the basic termini, U is an alkylene or heteroatom linker, V may be a heterocycle, and the right hand portion of the molecule represents the acidic termini. The presently claimed compounds do not contain the acidic termini of WO 95/18111.

In U.S. Patent No. 5,463,071, Himmelsbach et al depict cell aggregation inhibitors which are 5-membered heterocycles of the formula:



wherein the heterocycle may be aromatic and groups A-B-C- and F-E-D- are attached to the ring system. A-B-C- can be a wide variety of substituents including a basic group attached to an aromatic ring. The F-E-D- group, however, would appear to be an acidic functionality which differs from the present invention. Furthermore, use of these compounds as inhibitors of factor Xa is not discussed.

Baker et al, in U.S. Patent No. 5,317,103, discuss 5-HT<sub>1</sub> agonists which are indole substituted five-membered heteroaromatic compounds of the formula:




5

wherein R<sup>1</sup> may be pyrrolidine or piperidine and A may be a basic group including amino and amidino. Baker et al, however, do not indicate that A can be a substituted ring system like that contained in the presently claimed heteroaromatics.

10

Baker et al, in WO 94/02477, discuss 5-HT<sub>1</sub> agonists which are imidazoles, triazoles, or tetrazoles of the formula:



15

wherein R<sup>1</sup> represents a nitrogen containing ring system or a nitrogen substituted cyclobutane, and A may be a basic group including amino and amidino. Baker et al, however, do not indicate that A can be a substituted ring system like that contained in the presently claimed heteroaromatics.

20

Tidwell et al, in *J. Med. Chem.* 1978, 21(7), 613-623, describe a series of diarylamidine derivatives including 3,5-bis(4-amidinophenyl)pyrrole. This series of compounds was tested against thrombin, trypsin, and pancreatic kallikrein. The presently claimed invention does not include these types of compounds.

25

Activated factor Xa, whose major practical role is the generation of thrombin by the limited proteolysis of prothrombin, holds a central position that links the intrinsic and extrinsic activation mechanisms in the final common

30

pathway of blood coagulation. The generation of thrombin, the final serine protease in the pathway to generate a fibrin clot, from its precursor is amplified by formation of prothrombinase complex (factor Xa, factor V,  $\text{Ca}^{2+}$  and phospholipid). Since it is calculated that one molecule of factor Xa can generate 138 molecules of thrombin (Elodi, S., Varadi, K.: *Optimization of conditions for the catalytic effect of the factor IXa-factor VIII Complex: Probable role of the complex in the amplification of blood coagulation.*

*Thromb. Res.* 1979, 15, 617-629), inhibition of factor Xa may be more efficient than inactivation of thrombin in interrupting the blood coagulation system.

Therefore, efficacious and specific inhibitors of factor Xa are needed as potentially valuable therapeutic agents for the treatment of thromboembolic disorders. It is thus desirable to discover new factor Xa inhibitors.


#### SUMMARY OF THE INVENTION

Accordingly, one object of the present invention is to provide novel nitrogen containing aromatic heterocycles which are useful as factor Xa inhibitors or pharmaceutically acceptable salts or prodrugs thereof.

It is another object of the present invention to provide pharmaceutical compositions comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

It is another object of the present invention to provide a method for treating thromboembolic disorders comprising administering to a host in need of such treatment a therapeutically effective amount of at least one of the compounds of the present invention or a pharmaceutically acceptable salt or prodrug form thereof.

These and other objects, which will become apparent during the following detailed description, have been achieved by the inventors' discovery that compounds of formula (I):




I

or pharmaceutically acceptable salt or prodrug forms thereof,  
 wherein A, B, D, E, G, J, M, R<sup>1a</sup>, R<sup>1b</sup>, s and m/z are defined  
 5 below, are effective factor Xa inhibitors.

#### DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

[1] Thus, in a first embodiment, the present invention  
 provides novel compounds of formula I:

10



I

or a stereoisomer or pharmaceutically acceptable salt thereof,  
 wherein;

15

ring M contains, in addition to J, 0-3 N atoms, provided that  
 if M contains 2 N atoms then R<sup>1b</sup> is not present and if M  
 contains 3 N atoms then R<sup>1a</sup> and R<sup>1b</sup> are not present;

20 J is N or NH;

D is selected from CN, C(=NR<sup>8</sup>)NR<sup>7</sup>R<sup>9</sup>, NHC(=NR<sup>8</sup>)NR<sup>7</sup>R<sup>9</sup>,  
 NR<sup>8</sup>CH(=NR<sup>7</sup>), C(O)NR<sup>7</sup>R<sup>8</sup>, and (CR<sup>8</sup>R<sup>9</sup>)<sub>t</sub>NR<sup>7</sup>R<sup>8</sup>, provided that D  
 is substituted meta or para to G on E;

25

E is selected from phenyl, pyridyl, pyrimidyl, pyrazinyl,  
 pyridazinyl, and piperidinyl substituted with 1 R;

alternatively, D-E-G together represent pyridyl substituted  
 30 with 1 R;

R is selected from H, halogen, (CH<sub>2</sub>)<sub>t</sub>OR<sup>3</sup>, C<sub>1-4</sub> alkyl, OCF<sub>3</sub>, and  
 CF<sub>3</sub>;

G is absent or is selected from  $\text{NHCH}_2$ ,  $\text{OCH}_2$ , and  $\text{SCH}_2$ , provided that when s is 0, then G is attached to a carbon atom on ring M;

5

Z is selected from a  $\text{C}_{1-4}$  alkylene,  $(\text{CH}_2)_r\text{O}(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{C}(\text{O})(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{O}(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{OC}(\text{O})(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{OC}(\text{O})\text{O}(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{OC}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})\text{O}(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{S}(\text{O})_p(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{SO}_2\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{SO}_2(\text{CH}_2)_r$ , and  $(\text{CH}_2)_r\text{NR}^3\text{SO}_2\text{NR}^3(\text{CH}_2)_r$ , provided that Z does not form a N-N, N-O, N-S,  $\text{NCH}_2\text{N}$ ,  $\text{NCH}_2\text{O}$ , or  $\text{NCH}_2\text{S}$  bond with ring M or group A;

15

$\text{R}^{1a}$  and  $\text{R}^{1b}$  are independently absent or selected from  $-(\text{CH}_2)_r\text{R}^{1'}$ ,  $\text{NCH}_2\text{R}^{1'}$ ,  $\text{OCH}_2\text{R}^{1'}$ ,  $\text{SCH}_2\text{R}^{1'}$ ,  $\text{N}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ ,  $\text{O}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ , and  $\text{S}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ , or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2  $\text{R}^4$  and which contains from 0-2 heteroatoms selected from the group consisting of N, O, and S;

20

$\text{R}^{1'}$  is selected from H,  $\text{C}_{1-3}$  alkyl, halo,  $(\text{CF}_2)_r\text{CF}_3$ ,  $\text{OR}^2$ ,  $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(\text{O})\text{R}^{2c}$ ,  $\text{OC}(\text{O})\text{R}^2$ ,  $(\text{CF}_2)_r\text{CO}_2\text{R}^{2c}$ ,  $\text{S}(\text{O})_p\text{R}^{2b}$ ,  $\text{NR}^2(\text{CH}_2)_r\text{OR}^2$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})\text{NHR}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2a}$ ,  $\text{OC}(\text{O})\text{NR}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ,  $\text{NR}^2\text{SO}_2\text{R}^{2b}$ ,  $\text{C}_{3-6}$  carbocyclic residue substituted with 0-2  $\text{R}^4$ , and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2  $\text{R}^4$ ;

30

$\text{R}^{1''}$  is selected from H,  $\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  $\text{S}(\text{O})\text{R}^{2b}$ ,  $\text{S}(\text{O})_2\text{R}^{2b}$ , and  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ;

35

$\text{R}^2$ , at each occurrence, is selected from H,  $\text{CF}_3$ ,  $\text{C}_{1-6}$  alkyl, benzyl,  $\text{C}_{3-6}$  carbocyclic residue substituted with 0-2  $\text{R}^{4b}$ ,

and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

5 R<sup>2a</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, C<sub>1-6</sub> alkyl, benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2 R<sup>4b</sup>, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

10 R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkyl, benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2 R<sup>4b</sup>, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting  
15 of N, O, and S substituted with 0-2 R<sup>4b</sup>;

R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkyl, benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2 R<sup>4b</sup>, and 5-6 membered heterocyclic system  
20 containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a 5 or 6 membered saturated, partially saturated or unsaturated ring  
25 substituted with 0-2 R<sup>4b</sup> which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

30 R<sup>3</sup>, at each occurrence, is selected from H, C<sub>1-4</sub> alkyl, and phenyl;

R<sup>3a</sup>, at each occurrence, is selected from H, C<sub>1-4</sub> alkyl, and phenyl;

35 A is selected from:  
C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4</sup>, and

5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4</sup>;

5 B is selected from:

X-Y, NR<sup>2</sup>R<sup>2a</sup>, C(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>C(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>,

C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4a</sup>, and

5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4a</sup>;

X is selected from C<sub>1-4</sub> alkylene, -CR<sup>2</sup>(CR<sup>2</sup>R<sup>2b</sup>)(CH<sub>2</sub>)<sub>t</sub>-, -C(O)-, -C(=NR)-, -CR<sup>2</sup>(NR<sup>1</sup>R<sup>2</sup>)-, -CR<sup>2</sup>(OR<sup>2</sup>)-, -CR<sup>2</sup>(SR<sup>2</sup>)-, -C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)-, -S(O)<sub>p</sub>-, -S(O)<sub>p</sub>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>S(O)<sub>p</sub>-, -S(O)<sub>2</sub>NR<sup>2</sup>-, -NR<sup>2</sup>S(O)<sub>2</sub>-, -NR<sup>2</sup>S(O)<sub>2</sub>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>S(O)<sub>2</sub>NR<sup>2</sup>-, -NR<sup>2</sup>S(O)<sub>2</sub>NR<sup>2</sup>-, -C(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)-, -C(O)NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -NR<sup>2</sup>C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)NR<sup>2</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>C(O)-, -NR<sup>2</sup>C(O)O-, -OC(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)NR<sup>2</sup>-, -NR<sup>2</sup>-, -NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>-, O, -CR<sup>2</sup>R<sup>2a</sup>O-, and -OCR<sup>2</sup>R<sup>2a</sup>-;

Y is selected from:

(CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a N-N, O-N, or S-N bond,

C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4a</sup>, and

5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4a</sup>;

R<sup>4</sup>, at each occurrence, is selected from =O, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, -CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>C(O)NR<sup>2</sup>R<sup>2a</sup>, CH(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, NHC(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>-C<sub>1-4</sub> alkyl, NR<sup>2</sup>SO<sub>2</sub>R<sup>5</sup>, S(O)<sub>p</sub>R<sup>5</sup>, (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>, NCH<sub>2</sub>R<sup>1</sup>′, OCH<sub>2</sub>R<sup>1</sup>′, SCH<sub>2</sub>R<sup>1</sup>′, N(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>′, O(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>′, and S(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>′,

alternatively, one  $R^4$  is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S;

- 5  $R^{4a}$ , at each occurrence, is selected from  $=O$ ,  $(CH_2)_rOR^2$ , halo,  $C_{1-4}$  alkyl,  $-CN$ ,  $NO_2$ ,  $(CH_2)_rNR^2R^{2a}$ ,  $(CH_2)_rC(O)R^{2b}$ ,  $NR^2C(O)R^{2b}$ ,  $C(O)NR^2R^{2a}$ ,  $NR^2C(O)NR^2R^{2a}$ ,  $CH(=NR^2)NR^2R^{2a}$ ,  $NHC(=NR^2)NR^2R^{2a}$ ,  $SO_2NR^2R^{2a}$ ,  $NR^2SO_2NR^2R^{2a}$ ,  $NR^2SO_2-C_{1-4}$  alkyl,  $NR^2SO_2R^5$ ,  $S(O)_pR^5$ , and  $(CF_2)_rCF_3$ ;

10

alternatively, one  $R^{4a}$  is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-1  $R^5$ ;

- 15  $R^{4b}$ , at each occurrence, is selected from  $=O$ ,  $(CH_2)_rOR^3$ , halo,  $C_{1-4}$  alkyl,  $-CN$ ,  $NO_2$ ,  $(CH_2)_rNR^3R^{3a}$ ,  $(CH_2)_rC(O)R^3$ ,  $NR^3C(O)R^{3a}$ ,  $C(O)NR^3R^{3a}$ ,  $NR^3C(O)NR^3R^{3a}$ ,  $CH(=NR^3)NR^3R^{3a}$ ,  $NH^3C(=NR^3)NR^3R^{3a}$ ,  $SO_2NR^3R^{3a}$ ,  $NR^3SO_2NR^3R^{3a}$ ,  $NR^3SO_2-C_{1-4}$  alkyl,  $NR^3SO_2CF_3$ ,  $NR^3SO_2$ -phenyl,  $S(O)_pCF_3$ ,  $S(O)_p-C_{1-4}$  alkyl,  $S(O)_p$ -phenyl, and  $(CF_2)_rCF_3$ ;

20

$R^5$ , at each occurrence, is selected from  $CF_3$ ,  $C_{1-6}$  alkyl, phenyl substituted with 0-2  $R^6$ , and benzyl substituted with 0-2  $R^6$ ;

25

$R^6$ , at each occurrence, is selected from H, OH,  $(CH_2)_rOR^2$ , halo,  $C_{1-4}$  alkyl, CN,  $NO_2$ ,  $(CH_2)_rNR^2R^{2a}$ ,  $(CH_2)_rC(O)R^{2b}$ ,  $NR^2C(O)R^{2b}$ ,  $NR^2C(O)NR^2R^{2a}$ ,  $CH(=NH)NH_2$ ,  $NHC(=NH)NH_2$ ,  $SO_2NR^2R^{2a}$ ,  $NR^2SO_2NR^2R^{2a}$ , and  $NR^2SO_2C_{1-4}$  alkyl;

30

$R^7$ , at each occurrence, is selected from H, OH,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkylcarbonyl,  $C_{1-6}$  alkoxy,  $C_{1-4}$  alkoxycarbonyl,  $(CH_2)_n$ -phenyl,  $C_{6-10}$  aryloxy,  $C_{6-10}$  aryloxycarbonyl,  $C_{6-10}$  arylmethylcarbonyl,  $C_{1-4}$  alkylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl,  $C_{6-10}$  arylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl,  $C_{1-6}$  alkylaminocarbonyl, phenylaminocarbonyl, and phenyl  $C_{1-4}$  alkoxycarbonyl;

35



R<sup>8</sup>, at each occurrence, is selected from H, C<sub>1-6</sub> alkyl and (CH<sub>2</sub>)<sub>n</sub>-phenyl;

5 alternatively, R<sup>7</sup> and R<sup>8</sup> combine to form a 5 or 6 membered saturated, ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

10 R<sup>9</sup>, at each occurrence, is selected from H, C<sub>1-6</sub> alkyl and (CH<sub>2</sub>)<sub>n</sub>-phenyl;

n, at each occurrence, is selected from 0, 1, 2, and 3;

15 m, at each occurrence, is selected from 0, 1, and 2;

p, at each occurrence, is selected from 0, 1, and 2;


r, at each occurrence, is selected from 0, 1, 2, and 3;

20 s, at each occurrence, is selected from 0, 1, and 2; and,

t, at each occurrence, is selected from 0 and 1;

25 provided that D-E-G-(CH<sub>2</sub>)<sub>s</sub>- and -Z-A-B are not both benzamidines.

30 [2] In a preferred embodiment, the present invention provides novel compounds of formulae Ia-Ih:



wherein, groups D-E- and -Z-A-B are attached to adjacent atoms on the ring;

- 5 Z is selected from a  $\text{CH}_2\text{O}$ ,  $\text{OCH}_2$ ,  $\text{CH}_2\text{NH}$ ,  $\text{NHCH}_2$ ,  $\text{C}(\text{O})$ ,  $\text{CH}_2\text{C}(\text{O})$ ,  $\text{C}(\text{O})\text{CH}_2$ ,  $\text{NHC}(\text{O})$ ,  $\text{C}(\text{O})\text{NH}$ ,  $\text{CH}_2\text{S}(\text{O})_2$ ,  $\text{S}(\text{O})_2(\text{CH}_2)$ ,  $\text{SO}_2\text{NH}$ , and  $\text{NHSO}_2$ , provided that Z does not form a N-N, N-O,  $\text{NCH}_2\text{N}$ , or  $\text{NCH}_2\text{O}$  bond with ring M or group A;
- 10 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2  $\text{R}^4$ :  
 phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, and isoindazolyl;
- 20
- 25 B is selected from: Y, X-Y,  $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ , and  $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ ;


X is selected from C<sub>1-4</sub> alkylene, -C(O)-, -C(=NR)-, -CR<sup>2</sup>(NR<sup>2</sup>R<sup>2a</sup>)-, -C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)-, -C(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)-, -C(O)NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -NR<sup>2</sup>C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)NR<sup>2</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>C(O)-, -NR<sup>2</sup>C(O)NR<sup>2</sup>-, -NR<sup>2</sup>-, -NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>-, O, -CR<sup>2</sup>R<sup>2a</sup>O-, and -OCR<sup>2</sup>R<sup>2a</sup>-;

Y is NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R<sup>4a</sup>;

cyclopropyl, cyclopentyl, cyclohexyl, phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl, benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl, benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, and isoindazolyl;


alternatively, Y is selected from the following bicyclic heteroaryl ring systems:



30

K is selected from O, S, NH, and N.

[3] In a more preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf:



5

wherein;

Z is selected from a C(O), CH<sub>2</sub>C(O), C(O)CH<sub>2</sub>, NHC(O), C(O)NH, C(O)N(CH<sub>3</sub>), CH<sub>2</sub>S(O)<sub>2</sub>, S(O)<sub>2</sub>(CH<sub>2</sub>), SO<sub>2</sub>NH, and NHSO<sub>2</sub>,  
 10 provided that Z does not form a N-N or NCH<sub>2</sub>N bond with ring M or group A.

15 [4] In an even more preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

E is phenyl substituted with R or 2-pyridyl substituted with R;  
 20 R;

D is selected from NH<sub>2</sub>, C(O)NH<sub>2</sub>, C(=NH)NH<sub>2</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>NHCH<sub>3</sub>, CH(CH<sub>3</sub>)NH<sub>2</sub>, and C(CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>, provided that D is substituted meta or para to ring M on E; and,

25

R is selected from H, OCH<sub>3</sub>, Cl, and F.

[5] In a further preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, and 6-(2-amino-2-propyl)pyrid-2-yl.

[6] In another even more preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

Z is C(O)CH<sub>2</sub> and CONH, provided that Z does not form a N-N bond with group A;

A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R<sup>4</sup>; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R<sup>4a</sup>;

R<sup>4</sup>, at each occurrence, is selected from OH, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, and (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>;

R<sup>4a</sup> is selected from C<sub>1-4</sub> alkyl, CF<sub>3</sub>, S(O)<sub>p</sub>R<sup>5</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and 1-CF<sub>3</sub>-tetrazol-2-yl;

R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl, phenyl, and benzyl;

X is CH<sub>2</sub> or C(O); and,

5

Y is selected from pyrrolidino and morpholino.

[7] In another further preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

15

B is selected from the group: 2-CF<sub>3</sub>-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF<sub>3</sub>-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

25

[8] In another even more preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

30

E is phenyl substituted with R or 2-pyridyl substituted with R;

35

D is selected from NH<sub>2</sub>, C(O)NH<sub>2</sub>, C(=NH)NH<sub>2</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>NHCH<sub>3</sub>, CH(CH<sub>3</sub>)NH<sub>2</sub>, and C(CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>, provided that D is substituted meta or para to ring M on E; and,

R is selected from H, OCH<sub>3</sub>, Cl, and F;

Z is C(O)CH<sub>2</sub> and CONH, provided that Z does not form a N-N bond with group A;

5 A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R<sup>4</sup>; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino,  
1,2,3-triazolyl, and imidazolyl, and is substituted with  
10 0-1 R<sup>4a</sup>;

R<sup>4</sup>, at each occurrence, is selected from OH, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, and (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>;

15 R<sup>4a</sup> is selected from C<sub>1-4</sub> alkyl, CF<sub>3</sub>, S(O)<sub>p</sub>R<sup>5</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and 1-CF<sub>3</sub>-tetrazol-2-yl;

R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl, phenyl, and benzyl;

20 X is CH<sub>2</sub> or C(O); and,

Y is selected from pyrrolidino and morpholino.

25 [9] In another further preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

30 D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-

aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, 6-(2-amino-2-propyl)pyrid-2-yl;

- 5 A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,
- 10 B is selected from the group: 2-CF<sub>3</sub>-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF<sub>3</sub>-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl,
- 15 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

[10] In a still further preferred embodiment, the present invention provides a novel compound of formula IIa.

[11] In another still further preferred embodiment, the present invention provides a novel compound of formula IIb.

25

[12] In another still further preferred embodiment, the present invention provides a novel compound of formula IIc.

30

[13] In another still further preferred embodiment, the present invention provides a novel compound of formula IID.

35 [14] In another still further preferred embodiment, the present invention provides a novel compound of formula IIe.



[15] In another still further preferred embodiment, the present invention provides a novel compound of formula II<sub>f</sub>.

5 [16] In another even more preferred embodiment, the present invention provides novel compounds of formulae II<sub>a</sub>-II<sub>f</sub>, wherein;

D is selected from C(=NR<sup>8</sup>)NR<sup>7</sup>R<sup>9</sup>, C(O)NR<sup>7</sup>R<sup>8</sup>, NR<sup>7</sup>R<sup>8</sup>, and CH<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>,  
10 provided that D is substituted meta or para to ring M on E;

E is phenyl substituted with R or pyridyl substituted with R;

15 R is selected from H, Cl, F, OR<sup>3</sup>, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, OCF<sub>3</sub>, and CF<sub>3</sub>;

Z is selected from C(O), CH<sub>2</sub>C(O), C(O)CH<sub>2</sub>, NHC(O), and C(O)NH,  
provided that Z does not form a N-N bond with ring M or  
group A;

20 R<sup>1a</sup> and R<sup>1b</sup> are independently absent or selected from  
-(CH<sub>2</sub>)<sub>r</sub>-R<sup>1'</sup>, NCH<sub>2</sub>R<sup>1''</sup>, OCH<sub>2</sub>R<sup>1''</sup>, SCH<sub>2</sub>R<sup>1''</sup>, N(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1'</sup>,  
O(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1'</sup>, and S(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1'</sup>, or combined to form  
a 5-8 membered saturated, partially saturated or  
25 unsaturated ring substituted with 0-2 R<sup>4</sup> and which  
contains from 0-2 heteroatoms selected from the group  
consisting of N, O, and S;

R<sup>1'</sup>, at each occurrence, is selected from H, C<sub>1-3</sub> alkyl, halo,  
30 (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>, OR<sup>2</sup>, NR<sup>2</sup>R<sup>2a</sup>, C(O)R<sup>2c</sup>, (CF<sub>2</sub>)<sub>r</sub>CO<sub>2</sub>R<sup>2c</sup>, S(O)<sub>p</sub>R<sup>2b</sup>,  
NR<sup>2</sup>(CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)<sub>2</sub>R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>,  
SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and NR<sup>2</sup>SO<sub>2</sub>R<sup>2b</sup>;

A is selected from one of the following carbocyclic and  
35 heterocyclic systems which are substituted with 0-2 R<sup>4</sup>;  
phenyl, piperidinyl, piperazinyl, pyridyl,  
pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,

pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,  
isothiazolyl, pyrazolyl, and imidazolyl;

5 B is selected from: Y, X-Y,  $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ , and  
 $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ ;

X is selected from  $\text{CH}_2$ ,  $-\text{CR}^2(\text{CR}^2\text{R}^{2b})(\text{CH}_2)_t-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{C}(=\text{NR})-$ ,  
 $-\text{CH}(\text{NR}^2\text{R}^{2a})-$ ,  $-\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{NR}^2\text{C}(\text{O})-$ ,  $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{NR}^2-$ ,  
and O;

10 Y is  $\text{NR}^2\text{R}^{2a}$ , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following  
carbocyclic and heterocyclic systems which are  
15 substituted with 0-2  $\text{R}^{4a}$ ;

phenyl, piperidinyl, piperazinyl, pyridyl,  
pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,  
pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyll,  
thiazolyl, isothiazolyl, pyrazolyl, imidazolyl,  
20 oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,  
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,  
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,  
1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,  
1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

25  $\text{R}^4$ , at each occurrence, is selected from =O, OH, Cl, F,  $\text{C}_{1-4}$   
alkyl,  $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  
 $\text{CH}(=\text{NH})\text{NH}_2$ ,  $\text{NHC}(=\text{NH})\text{NH}_2$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ,  $\text{NR}^2\text{SO}_2\text{-C}_{1-4}$  alkyl,  
 $\text{NR}^2\text{SO}_2\text{R}^5$ ,  $\text{S}(\text{O})_p\text{R}^5$ , and  $(\text{CF}_2)_r\text{CF}_3$ ;

30  $\text{R}^{4a}$ , at each occurrence, is selected from =O, OH, Cl, F,  $\text{C}_{1-4}$   
alkyl,  $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  
 $\text{CH}(=\text{NH})\text{NH}_2$ ,  $\text{NHC}(=\text{NH})\text{NH}_2$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ,  $\text{NR}^2\text{SO}_2\text{-C}_{1-4}$  alkyl,  
 $\text{NR}^2\text{SO}_2\text{R}^5$ ,  $\text{S}(\text{O})_p\text{R}^5$ ,  $(\text{CF}_2)_r\text{CF}_3$ , and 1- $\text{CF}_3$ -tetrazol-2-yl;

35  $\text{R}^5$ , at each occurrence, is selected from  $\text{CF}_3$ ,  $\text{C}_{1-6}$  alkyl,  
phenyl substituted with 0-2  $\text{R}^6$ , and benzyl substituted  
with 0-2  $\text{R}^6$ ;

R<sup>6</sup>, at each occurrence, is selected from H, =O, OH, OR<sup>2</sup>, Cl, F, CH<sub>3</sub>, CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, CH(=NH)NH<sub>2</sub>, NHC(=NH)NH<sub>2</sub>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

5 R<sup>7</sup>, at each occurrence, is selected from H, OH, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkoxy, C<sub>1-4</sub> alkoxy carbonyl, benzyl, C<sub>6-10</sub> aryloxy, C<sub>6-10</sub> aryloxy carbonyl, C<sub>6-10</sub> arylmethylcarbonyl, C<sub>1-4</sub> alkylcarbonyloxy C<sub>1-4</sub> alkoxy carbonyl, C<sub>6-10</sub> arylcarbonyloxy C<sub>1-4</sub> alkoxy carbonyl, C<sub>1-6</sub> alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C<sub>1-4</sub> alkoxy carbonyl;

10 R<sup>8</sup>, at each occurrence, is selected from H, C<sub>1-6</sub> alkyl and benzyl; and

15 alternatively, R<sup>7</sup> and R<sup>8</sup> combine to form a morpholino group; and,

20 R<sup>9</sup>, at each occurrence, is selected from H, C<sub>1-6</sub> alkyl and benzyl.

25 [17] In a another further preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

E is phenyl substituted with R or 2-pyridyl substituted with R;

30 R is selected from H, Cl, F, OCH<sub>3</sub>, CH<sub>3</sub>, OCF<sub>3</sub>, and CF<sub>3</sub>;

Z is selected from a C(O)CH<sub>2</sub> and C(O)NH, provided that Z does not form a N-N bond with group A;

35 R<sup>1a</sup> is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>, OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2c</sup>, CH<sub>2</sub>C(O)R<sup>2c</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

R<sup>1b</sup> is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>, OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2c</sup>, CH<sub>2</sub>C(O)R<sup>2c</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

5

A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R<sup>4</sup>; phenyl, pyridyl, pyrimidyl, furanyl, thiophenyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, and imidazolyl;

10

B is selected from: Y and X-Y;

X is selected from CH<sub>2</sub>, -CR<sup>2</sup>(CR<sup>2</sup>R<sup>2b</sup>)-, -C(O)-, -C(=NR)-, -CH(NR<sup>2</sup>R<sup>2a</sup>)-, -C(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)-, -NR<sup>2</sup>C(O)NR<sup>2</sup>-, -NR<sup>2</sup>-, and O;

15

Y is NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R<sup>4a</sup>;

20

phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

25

30

R<sup>2</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

35

R<sup>2a</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, OCH<sub>3</sub>, CH<sub>3</sub>,  
benzyl, and phenyl;

5 R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, OCH<sub>3</sub>, CH<sub>3</sub>,  
benzyl, and phenyl;

alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a 5 or 6 membered  
saturated, partially unsaturated, or unsaturated ring  
which contains from 0-1 additional heteroatoms selected  
10 from the group consisting of N, O, and S;

R<sup>3</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, and  
phenyl;

15 R<sup>3a</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, and  
phenyl;

R<sup>4</sup>, at each occurrence, is selected from OH, Cl, F, CH<sub>3</sub>,  
CH<sub>2</sub>CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>,  
20 and CF<sub>3</sub>;

R<sup>4a</sup>, at each occurrence, is selected from OH, Cl, F, CH<sub>3</sub>,  
CH<sub>2</sub>CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>,  
S(O)<sub>p</sub>R<sup>5</sup>, CF<sub>3</sub>, and 1-CF<sub>3</sub>-tetrazol-2-yl;

25 R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl,  
phenyl substituted with 0-2 R<sup>6</sup>, and benzyl substituted  
with 1 R<sup>6</sup>;

30 R<sup>6</sup>, at each occurrence, is selected from H, OH, OCH<sub>3</sub>, Cl, F,  
CH<sub>3</sub>, CN, NO<sub>2</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

R<sup>7</sup>, at each occurrence, is selected from H, OH, C<sub>1-3</sub> alkyl,  
C<sub>1-3</sub> alkylcarbonyl, C<sub>1-3</sub> alkoxy, C<sub>1-4</sub> alkoxycarbonyl,  
35 benzyl, phenoxy, phenoxycarbonyl, benzylcarbonyl, C<sub>1-4</sub>  
alkylcarbonyloxy C<sub>1-4</sub> alkoxycarbonyl, phenylcarbonyloxy  
C<sub>1-4</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylaminocarbonyl,  
phenylaminocarbonyl, and phenyl C<sub>1-4</sub> alkoxycarbonyl;

R<sup>8</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, and benzyl;  
and,

5 alternatively, R<sup>7</sup> and R<sup>8</sup> combine to form a morpholino group;

R<sup>9</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, and benzyl.

10 [18] In a another still further preferred embodiment, the present invention provides novel compounds of formulae IIa-IIf, wherein;

R<sup>1a</sup> is absent or is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>,  
15 OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>,  
CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2c</sup>, CH<sub>2</sub>C(O)R<sup>2c</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

R<sup>1b</sup> is absent or is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>,  
OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>,  
20 CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2b</sup>, CH<sub>2</sub>C(O)R<sup>2b</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

A is selected from one of the following carbocyclic and  
heterocyclic systems which are substituted with 0-2 R<sup>4</sup>;  
phenyl, pyridyl, and pyrimidyl;

25 B is selected from: Y and X-Y;

X is selected from -C(O)- and O;

30 Y is NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a O-N bond;

alternatively, Y is selected from one of the following  
carbocyclic and heterocyclic systems which are  
substituted with 0-2 R<sup>4a</sup>;

35 phenyl, piperazinyl, pyridyl, pyrimidyl,  
morpholinyl, pyrrolidinyl, imidazolyl, and 1,2,3-  
triazolyl;

- R<sup>2</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;
- 5 R<sup>2a</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;
- R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, OCH<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;
- 10 R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, OCH<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;
- alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a ring system selected from pyrrolidinyl, piperazinyl and morpholino;
- 15 R<sup>4</sup>, at each occurrence, is selected from Cl, F, CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, and CF<sub>3</sub>;
- R<sup>4a</sup>, at each occurrence, is selected from Cl, F, CH<sub>3</sub>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>5</sup>, and CF<sub>3</sub>; and,
- 20 R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub> and CH<sub>3</sub>.
- 25 [19] Specifically preferred compounds of the present invention are selected from the group:
- 1-(3-amidinophenyl)-2-[[ (2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]pyrrole;
- 30 1-(3-amidinophenyl)-2-[[ (2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]pyrrole;
- 35 1-(3-amidinophenyl)-2-[[ (2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]-4-bromopyrrole;
- 1-(3-amidinophenyl)-2-[[5-(2'-aminosulfonylphen-1-yl) pyridin-2-yl]-aminocarbonyl]pyrrole;
- 40 1-benzyl-3-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole;

- 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole;
- 5 1-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 1-(3-amidinophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 10 1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylamino]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-(2'-(5''-CF<sub>3</sub>-tetrazolyl)-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-chloro-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethyl-pyrazole;
- 25 1-(3-amidinophenyl)-4-methoxy-5-((2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethyl-pyrazole;
- 30 1-(3-amidinophenyl)-3-methyl-5-(4'-(imidazol-1-yl-phenyl)aminocarbonyl)pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(2''-sulfonylmethyl)phenoxyphenyl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methylcarbonyl]pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 40 1-(3-amidinophenyl)-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)tetrazole;
- 45 1-(3-amidinophenyl)-5-((2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylsulfoxide]tetrazole;
- 50 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylsulfonyl]tetrazole;
- 55 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;



- 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-aminosulfonylphenyl-1-yl)pyrimidin-2-yl]-aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-2-chloro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-2-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-4'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-chloro-2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 25 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-trifluoromethylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-methyl-5-[(3-chloro-2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl) (N'-methyl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-n-butyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-n-butyl-5-[(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-3-n-butyl-5-[(2'-trifluoromethylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 55

- 1-(3-amidinophenyl)-4-methoxy-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethyl-pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[(4'-trifluoromethylphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-4-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 10 1-(3-amidinophenyl)-5-[(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl)-aminocarbonyl]-1,2,3-triazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 15 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-1,2,4-triazole;
- 3-methyl-1-(3-amidinophenyl)-5-(4'-(4''-chlorophenyl)thiazol-2'-yl)aminocarbonyl]pyrazole;
- 20 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfide-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 25 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfoxide-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-(3-amidino)phenyl-3-methyl-5-[4'-(carboxymethyl)phenylaminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminocarbonyl)phenylaminocarbonyl]pyrazole;
- 35 1-(3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminosulfonyl)phenylaminocarbonyl]pyrazole;
- 40 1-(3-amidino)phenyl-3-methyl-5-[(4'-tert-butylaminosulfonylphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(4'-aminosulfonylphenyl)aminocarbonyl]pyrazole;
- 45 1-(3-amidino)phenyl-3-methyl-5-[(4'-trifluoromethylphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(4'-benzylsulfonylpiperidyl)aminocarbonyl]pyrazole;
- 50 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-N-methylaminocarbonyl]-3-methyl-pyrazole;
- 55 1-(3-amidinophenyl)-5-[(4'-fluoro-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methyl-pyrazole;

- 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 5 1-(3-cyanophenyl)-5-[[5-(2'-aminosulfonylphenyl) pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 10 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl)-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 15 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl)-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-n-butylpyrazole;
- 20 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-n-butylpyrazole;
- 25 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-n-butylpyrazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-4-methoxypyrazole;
- 30 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 35 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 40 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl)-[1,1']-biphen-4-yl)methylcarbonyl]pyrazole;
- 45 1-(3-aminocarbonylphenyl)-5-[5-[(2'-aminosulfonylphen-1-yl)pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 55

- 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 5 1-(3-cyanophenyl)-5-[[4'-(imidazol-1-yl)phenyl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[[4'-(morpholin-1-yl)phenyl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 10 1-(3-aminocarbonylphenyl)-5-[[4'-(morpholin-1-yl)phenyl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 15 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 20 1-(3-amidinophenyl)-5-[[4'-(3-methyltetrazol-1-yl)phenyl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-(2'-naphthylaminosulfonyl)-3-methyl-pyrazole;
- 25 1-(3-amidinophenyl)-5-[[4-bromophenyl]aminosulfonyl]-3-methyl-pyrazole;
- 30 1-(3-aminomethylphenyl)-5-[[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[[2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[[2'-trifluoromethylphenyl]pyrid-2-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[2'-aminosulfonyl-1-yl]pyrimid-5-yl]aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[[2'-fluoro-[1,1']-biphen-4-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[3-chloro-(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-3-methyl-5-[[3-fluoro-2'-fluoro-[1,1']-biphen-4-yl]aminocarbonyl]pyrazole;
- 50 1-(3-amidinophenyl)-3-methyl-5-[[3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl]aminocarbonyl]pyrazole;
- 55 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-fluorophen-1-yl)pyrid-2-yl]aminocarbonyl]pyrazole;

- 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-tertbutylaminosulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-aminosulfonylphenyl)-[1,6]-dihydropyrimid-2-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4-(pyrid-3'-yl)phen-1-yl)aminocarbonyl]pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-[[2-(2'-pyridyl)ethyl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-phenylpropyl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[4-(pyrid-2'-yl)phen-1-ylaminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(4-(isopropoxy)phenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-trifluoromethylphenyl)-pyrimidin-2-yl]aminocarbonyl]pyrazole;
- 25 1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidinosulfonyl)phenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidinocarbonyl)phenyl)aminocarbonyl]pyrazole;
- 30 1-(3-amidino-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-(3-aminocarbonyl-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-methyl-3-(3-amidino)phenyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[[4-(pyrazol-4'-yl)phen-1-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-methylsulfonylphenyl)pyrid-2-yl]aminocarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole;
- 50 1-(3-cyanophenyl)-3-methyl-5-[[5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole,;
- 55 1-(3-aminocarbonylphenyl)-3-methyl-5-[[5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole;

- 1-(3-(N-aminoamidino)phenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-(3-(N-aminoamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 10 1-(3-(N-methyl-N-hydroxyamidino)phenyl)-3-methyl-5-[(4'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 15 1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 25 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 30 1-(3-amidinophenyl)-5-[[5-(2'-trifluoromethylphen-1-yl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 35 1-(3-amidinophenyl)-5-[(4'-bromophen-1-yl)aminocarbonyl]tetrazole;
- 1-(3-aminocarbonylphenyl)-5-[[5-(2'-trifluoromethylphen-1-yl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 40 5-(3-amidinophenyl)-1-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole;
- 1-[(3-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 45 1-[(4-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 50 1-(3-amidinophenyl)-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 1-(3-amidinophenyl)-5-chloro-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 55 5-(3-amidinophenyl)-2-methyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole;

- 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[(4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-[(4'-(1,2,4-triazol-2-yl)-phenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-cyclohexylphenyl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[[1,1']-biphen-4-ylaminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(4'-morpholinophenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-((2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 25 1-(3-aminomethylphenyl)-3-methyl-5-[(4'-((2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-dimethylamino)carbonylamino)phen-1'-yl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-diethylamino)phenyl)aminocarbonyl]pyrazole;
- 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-N,N-diethylamino)phenyl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole;
- 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-acetylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole;
- 50 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole,;
- 55 1-(3-amidinophenyl)-3-methyl-5-[(4'-piperizin-1-yl-phenyl)aminocarbonyl]pyrazole;

- 1-(3-amidinophenyl)-3-trifluoromethyl-5-((4'-cyclohexylphenyl)aminocarbonyl)pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholino)-3'-chlorophenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 10 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfinyl)pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole;
- 15 1-(3-aminocarbonylphenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole;
- 20 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methyl]tetrazole;
- 1-(3-amidinophenyl)-5-[(4'-cyclopentylloxyphenyl)aminocarbonyl]-3-methyl-pyrazole;
- 25 1-(3-amidinophenyl)-5-[(3-((pyrid-2-yl)methylamino)phenyl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-(N-morpholino)-3-chlorophenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-pyrrolidinocarbonyl)-3'-chlorophenyl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholinocarbonyl)-3-chlorophenyl)aminocarbonyl]pyrazole;
- 40 1-(3-cyanophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 45 1-(3-amidinophenyl)-5-[(4'-(N-methyltetrazolon-1-yl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3'-aminocarbonylphenyl)-5-[(2'-aminosulfonylphenyl-[1,1']-biphen-4-yl)methylcarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[4'-(pyrrolidinomethyl)phenyl)aminocarbonyl]-3-methyl-pyrazole;
- 55



- 1-(3-aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-(2'-aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amino-4'-chlorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 10 1-(3-amino-4'-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amino-4'-methoxyphenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 15 1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 20 1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 1-(3-amino-4'-methoxyphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 25 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonylphenyl)pyrid-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethyl-4'-methylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 30 1-(3-aminomethyl-4'-fluorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 35 1-(3-aminomethylphenyl)-5-[(4'-(N-pyrrolidino-carbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-Ethylcarboxyamidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methyl-pyrazole;
- 40 1-(3-(1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 45 1-(3-(1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 50 1-[3-[N-((5-methyl-2-oxo-1,3-dioxol-4-yl)methoxycarbonyl)amidino]phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 55 1-(pyrid-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;

- 1-(6-bromopyridin-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-(3-amino-4-chlorophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 1-(3-amino-4-chlorophenyl)-5-[(4'-(1-pyrrolidinocarbonyl)phenyl)aminocarbonyl]tetrazole;
- 10 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 15 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 20 1-(3-aminomethylphenyl)-5-[(2'-methylsulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 25 1-[3-(methylaminomethyl)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-[3-(methylaminomethyl)phenyl]-5-[(2'-methylsulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 30 1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-methoxy-3-trifluoromethyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 35 1-(3-aminomethylphenyl)-5-[(3-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 40 1-(3-aminomethylphenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 45 1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)-[1,6-dihydro]pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 55 1-[3-(2'-ethylaminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

- 1-[3-(1-(N-morpholino)imino)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 5 1-(3-aminomethylphenyl)-5-[2-(2'-aminosulfonyl-[1,1']-biphen-4-yl)-1-hydroxyethyl]-3-trifluoromethyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 10 1-(3-aminomethylphenyl)-5-[(5-(2'-methylsulfonyl-phenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 15 1-[3-amidinophenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 20 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylmethyl]-3-trifluoromethyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonylmethyl)pyrazole;
- 25 1-(3-amidino)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylaminosulfonylmethyl)pyrazole;
- 30 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylaminosulfonylmethyl)pyrazole;
- 1-(3-(N-carboxymethyl)amidinophenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-methyl-pyrazole;
- 35 1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 40 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-methyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 45 1-(3-aminomethyl-4-methylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 50 1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethyl-4-chloro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 55

- 1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 5 1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 10 1-(3-aminomethyl)phenyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 15 1-(3-aminomethylphenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-[(3-fluoro-4-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 20 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole;
- 1-(3-amidinophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole;
- 25 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole;
- 1-(3-amidinophenyl)-3-trifluoromethyl-5-(2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 30 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-(2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 35 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-(3-trifluoromethyl-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 40 1-(3-aminomethylphenyl)-3-ethyl-5-[(3-fluoro-2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-aminomethylphenyl)-3-ethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 45 1-(3-aminomethylphenyl)-3-ethyl-5-[(2-fluoro-4-(2-methylsulfonylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 50 1-[(6-(aminomethyl)pyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 55 1-[(6-(N-hydroxyamidino)pyrid-2-yl)]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;

- 1-[(6-amidinopyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-[6-amidinopyrid-2-yl]-3-methyl-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-aminomethylphenyl)-3-methyl-5-[(2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 10 1-(3-aminomethylphenyl)-3-methyl-5-[4'-(3"-methyl-5"-oxo-3"-pyrazolin-2"-yl)-phenyl]aminocarbonyl]pyrazole;
- 1-[3-(aminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 15 1-(3-aminomethyl-4-fluorophenyl)-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 ethyl 1-[3-(aminomethyl)-phenyl]-5-[3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate;
- 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylic
- 25 acid;
- 1-[3-(aminomethyl)phenyl]-3-[aminocarbonyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 ethyl 1-[3-(aminomethyl)-phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-4-carboxylate;
- 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 35 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole;
- 40 1-[3-(aminomethyl)phenyl]-5-[(4-(5-(methoxyaminocarbonyl)imidazol-1-yl)phen-1-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole; and,
- 45 1-(3-aminomethylphenyl)-5-[(4-(5-methyl-1,2,3-triazol-1-yl)phen-1-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- and pharmaceutically acceptable salts thereof.

50

In a second embodiment, the present invention provides novel pharmaceutical compositions, comprising: a pharmaceutically acceptable carrier and a therapeutically

effective amount of a compound of formula (I) or a pharmaceutically acceptable salt form thereof.

In a third embodiment, the present invention provides a novel method for treating or preventing a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable salt form thereof.

#### DEFINITIONS

The compounds herein described may have asymmetric centers. Compounds of the present invention containing an asymmetrically substituted atom may be isolated in optically active or racemic forms. It is well known in the art how to prepare optically active forms, such as by resolution of racemic forms or by synthesis from optically active starting materials. Many geometric isomers of olefins, C=N double bonds, and the like can also be present in the compounds described herein, and all such stable isomers are contemplated in the present invention. Cis and trans geometric isomers of the compounds of the present invention are described and may be isolated as a mixture of isomers or as separated isomeric forms. All chiral, diastereomeric, racemic forms and all geometric isomeric forms of a structure are intended, unless the specific stereochemistry or isomeric form is specifically indicated.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. When a substituent is keto (i.e., =O), then 2 hydrogens on the atom are replaced.

When any variable (e.g., R<sup>6</sup>) occurs more than one time in any constituent or formula for a compound, its definition at each occurrence is independent of its definition at every other occurrence. Thus, for example, if a group is shown to be substituted with 0-2 R<sup>6</sup>, then said group may optionally be

substituted with up to two R<sup>6</sup> groups and R<sup>6</sup> at each occurrence is selected independently from the definition of R<sup>6</sup>. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

5       When a bond to a substituent is shown to cross a bond connecting two atoms in a ring, then such substituent may be bonded to any atom on the ring. When a substituent is listed without indicating the atom via which such substituent is bonded to the rest of the compound of a given formula, then  
10       such substituent may be bonded via any atom in such substituent. Combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

As used herein, "C<sub>1-6</sub> alkyl" is intended to include both  
15       branched and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, examples of which include, but are not limited to, methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, sec-butyl, t-butyl, pentyl, and hexyl; "Alkenyl" is intended to include  
20       hydrocarbon chains of either a straight or branched configuration and one or more unsaturated carbon-carbon bonds which may occur in any stable point along the chain, such as ethenyl, propenyl, and the like.

"Halo" or "halogen" as used herein refers to fluoro,  
25       chloro, bromo, and iodo; and "counterion" is used to represent a small, negatively charged species such as chloride, bromide, hydroxide, acetate, sulfate, and the like.

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable 3- to 7-membered monocyclic or  
30       bicyclic or 7- to 13-membered bicyclic or tricyclic, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl,; [3.3.0]bicyclooctane,  
35       [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin).

As used herein, the term "heterocycle" or "heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic ring which is saturated partially unsaturated or unsaturated (aromatic), and which consists of carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O and S and including any bicyclic group in which any of the above-defined heterocyclic rings is fused to a benzene ring. The nitrogen and sulfur heteroatoms may optionally be oxidized. The heterocyclic ring may be attached to its pendant group at any heteroatom or carbon atom which results in a stable structure. The heterocyclic rings described herein may be substituted on carbon or on a nitrogen atom if the resulting compound is stable. If specifically noted, a nitrogen in the heterocycle may optionally be quaternized. It is preferred that when the total number of S and O atoms in the heterocycle exceeds 1, then these heteroatoms are not adjacent to one another. It is preferred that the total number of S and O atoms in the heterocycle is not more than 1. As used herein, the term "aromatic heterocyclic system" is intended to mean a stable 5- to 7- membered monocyclic or bicyclic or 7- to 10-membered bicyclic heterocyclic aromatic ring which consists of carbon atoms and from 1 to 4 heterotams independently selected from the group consisting of N, O and S. It is preferred that the total number of S and O atoms in the aromatic heterocycle is not more than 1.

Examples of heterocycles include, but are not limited to, 1H-indazole, 2-pyrrolidonyl, 2H,6H-1,5,2-dithiazinyl, 2H-pyrrolyl, 3H-indolyl, 4-piperidonyl, 4aH-carbazole, 4H-quinolizinyll, 6H-1,2,5-thiadiazinyl, acridinyl, azocinyl, benzimidazolyl, benzofuranyl, benzothiofuranyl, benzothiophenyl, benzoxazolyl, benzthiazolyl, benztriazolyl, benztetrazolyl, benzisoxazolyl, benzisothiazolyl, benzimidazalonyl, carbazolyl, 4aH-carbazolyl,  $\beta$ -carbolinyl, chromanyl, chromenyl, cinnolinyl, decahydroquinolinyl, 2H,6H-1,5,2-dithiazinyl, dihydrofuro[2,3-b]tetrahydrofuran, furanyl, furazanyl, imidazolidinyl, imidazolinyl, imidazolyl, 1H-



indazolyl, indolenyl, indolinyl, indoliziny, indolyl,  
isobenzofuranyl, isochromanyl, isoindazolyl, isoindolinyl,  
isoindolyl, isoquinolinyl, isothiazolyl, isoxazolyl,  
morpholinyl, naphthyridinyl, octahydroisoquinolinyl,  
5 oxadiazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-  
oxadiazolyl, 1,3,4-oxadiazolyl, oxazolidinyl., oxazolyl,  
oxazolidinylperimidinyl, phenanthridinyl, phenanthrolinyl,  
phenarsazinyl, phenazinyl, phenothiazinyl, phenoxathiinyl,  
phenoxazinyl, phthalazinyl, piperazinyl, piperidinyl,  
10 pteridinyl, piperidonyl, 4-piperidonyl, pteridinyl, purinyl,  
pyranyl, pyrazinyl, pyrazolidinyl, pyrazolinyl, pyrazolyl,  
pyridazinyl, pyridooxazole, pyridoimidazole, pyridothiazole,  
pyridinyl, pyridyl, pyrimidinyl, pyrrolidinyl, pyrrolinyl,  
pyrrolyl, quinazolinyl, quinolinyl, 4H-quinoliziny,  
15 quinoxalinyl, quinuclidinyl, carbolinyl, tetrahydrofuranyl,  
tetrahydroisoquinolinyl, tetrahydroquinolinyl, 6H-1,2,5-  
thiadiazinyl, 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-  
thiadiazolyl, 1,3,4-thiadiazolyl, thianthrenyl, thiazolyl,  
thienyl, thienothiazolyl, thienooxazolyl, thienoimidazolyl,  
20 thiophenyl, triazinyl, 1,2,3-triazolyl, 1,2,4-triazolyl,  
1,2,5-triazolyl, 1,3,4-triazolyl, xanthenyl. Preferred  
heterocycles include, but are not limited to, pyridinyl,  
furanyl, thienyl, pyrrolyl, pyrazolyl, imidazolyl, indolyl,  
benzimidazolyl, 1H-indazolyl, oxazolidinyl, benzotriazolyl,  
25 benzisoxazolyl, oxindolyl, benzoxazoliny, or isatinoyl. Also  
included are fused ring and spiro compounds containing, for  
example, the above heterocycles.

The phrase "pharmaceutically acceptable" is employed  
herein to refer to those compounds, materials, compositions,  
30 and/or dosage forms which are, within the scope of sound  
medical judgment, suitable for use in contact with the tissues  
of human beings and animals without excessive toxicity,  
irritation, allergic response, or other problem or  
complication, commensurate with a reasonable benefit/risk  
35 ratio.

As used herein, "pharmaceutically acceptable salts" refer  
to derivatives of the disclosed compounds wherein the parent  
compound is modified by making acid or base salts thereof.

Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of basic residues such as amines; alkali or organic salts of acidic residues such as carboxylic acids; and the like. The pharmaceutically acceptable salts include the conventional non-toxic salts or the quaternary ammonium salts of the parent compound formed, for example, from non-toxic inorganic or organic acids. For example, such conventional non-toxic salts include those derived from inorganic acids such as hydrochloric, hydrobromic, sulfuric, sulfamic, phosphoric, nitric and the like; and the salts prepared from organic acids such as acetic, propionic, succinic, glycolic, stearic, lactic, malic, tartaric, citric, ascorbic, pantoic, maleic, hydroxymaleic, phenylacetic, glutamic, benzoic, salicylic, sulfanilic, 2-acetoxybenzoic, fumaric, toluenesulfonic, methanesulfonic, ethane disulfonic, oxalic, isethionic, and the like.

The pharmaceutically acceptable salts of the present invention can be synthesized from the parent compound which contains a basic or acidic moiety by conventional chemical methods. Generally, such salts can be prepared by reacting the free acid or base forms of these compounds with a stoichiometric amount of the appropriate base or acid in water or in an organic solvent, or in a mixture of the two; generally, nonaqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile are preferred. Lists of suitable salts are found in *Remington's Pharmaceutical Sciences*, 17th ed., Mack Publishing Company, Easton, PA, 1985, p. 1418, the disclosure of which is hereby incorporated by reference.

"Prodrugs" are intended to include any covalently bonded carriers which release the active parent drug according to formula (I) *in vivo* when such prodrug is administered to a mammalian subject. Prodrugs of a compound of formula (I) are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or *in vivo*, to the parent compound. Prodrugs include compounds of formula (I) wherein a

hydroxy, amino, or sulfhydryl group is bonded to any group that, when the prodrug or compound of formula (I) is administered to a mammalian subject, cleaves to form a free hydroxyl, free amino, or free sulfhydryl group, respectively.

5 Examples of prodrugs include, but are not limited to, acetate, formate and benzoate derivatives of alcohol and amine functional groups in the compounds of formula (I), and the like. Preferred prodrugs are amidine prodrugs wherein D is  $C(=NR^7)NH_2$  or its tautomer  $C(=NH)NHR^7$  and  $R^7$  is selected from

10 OH,  $C_{1-4}$  alkoxy,  $C_{6-10}$  aryloxy,  $C_{1-4}$  alkoxycarbonyl,  $C_{6-10}$  aryloxycarbonyl,  $C_{6-10}$  arylmethylcarbonyl,  $C_{1-4}$  alkylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl, and  $C_{6-10}$  arylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl. More preferred prodrugs are where  $R^7$  is

15 OH, methoxy, ethoxy, benzyloxycarbonyl, methoxycarbonyl, and methylcarbonyloxymethoxycarbonyl.

"Stable compound" and "stable structure" are meant to indicate a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into an efficacious therapeutic

20 agent.

### SYNTHESIS

The compounds of the present invention can be prepared in a number of ways known to one skilled in the art of organic

25 synthesis. The compounds of the present invention can be synthesized using the methods described below, together with synthetic methods known in the art of synthetic organic chemistry, or by variations thereon as appreciated by those skilled in the art. Preferred methods include, but are not


30 limited to, those described below. The reactions are performed in a solvent appropriate to the reagents and materials employed and suitable for the transformations being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the

35 molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired

compound of the invention. It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene and Wuts (*Protective Groups In Organic Synthesis*, Wiley and Sons, 1991). All references cited herein are hereby incorporated in their entirety herein by reference.

The compounds of Formula I in which ring M is pyrrole can be prepared by the procedures described in Schemes 1-9. In Scheme 1 is shown how to prepare pyrroles in which the group Q-E is attached to the pyrrole nitrogen, wherein Q is a functionality that can be converted into D of Formula I, R<sup>e</sup> is functionality that can be converted into Z-A-B of Formula I and R<sup>f</sup> is or can be converted into R<sup>1a</sup> of Formula I. Oxidation of a furan with bromine in acetic acid can afford a 2,5-diacetoxidyhydrofuran which can react with amine Q-E-NH<sub>2</sub> to afford a pyrrole. Vilsmeier-Haack formylation with phosphorous oxychloride and DMF preferentially can acylate the pyrrole ring at C-2. Oxidation of the resulting aldehyde can give a carboxylic acid. The carboxylic acid can then be converted into amine derivatives using either the Hofmann degradation of the derived primary amide (Huisgen et. al. *Chem. Ber.* 1960, 93, 65) or the Curtius rearrangement of the derived acyl azide (*J. Prakt. Chem.* 1909, 42, 477). Derivatives which contain a sulfur atom attached to the pyrrole ring can be obtained by direct sulfonation with pyridine sulfur trioxide complex to give the sulfonic acids or treatment with copper (II) thiocyanate (*J. Het. Chem.* 1988, 25, 431) followed by the reduction of the intermediate thiocyanate with sodium borohydride to give a mercaptan.


Scheme 1



- 5 In Scheme 2 is shown how to prepare pyrroles in which Q-E is attached to the 2-position, wherein  $\text{R}^f$  and  $\text{R}^g$  collectively are hydrogen or a group that can be converted into  $\text{R}^{1a}$  and  $\text{R}^{1b}$  of Formula I. The Hantzsch pyrrole synthesis is a versatile reaction involving the cyclization of an appropriate  $\beta$ -
- 10 ketoester with an  $\alpha$ -halo ketone or aldehyde in the presence of a primary amine (*Ber. Dtsch. Chem. Ges.* 1890, 23, 1474). The  $\beta$ -ketoesters can be prepared from acid chlorides ( $\text{X} = \text{Cl}$ ) by the addition of the magnesium anion of potassium alkylmalonate followed by decarboxylation (*Synthesis* 1993, 290).
- 15 Alternatively,  $\beta$ -ketoesters can be prepared from an appropriate aldehyde ( $\text{R} = \text{H}$ ) by Reformatsky reaction with an  $\alpha$ -bromoacetate followed by oxidation. Cyclization with an  $\alpha$ -halo ketone or aldehyde in the presence of a primary amine can afford pyrroles. Acidic hydrolysis of the 3-carboalkoxy
- 20 pyrrole can afford the carboxylic acids. Pyrroles which contain a 3-amino substituent can be prepared from the acids by treatment with phosphoryl azide and triethylamine to effect a Curtius rearrangement to afford the isocyanates (*J. Med.*

Chem. 1981, 24, 33) which upon hydrolysis can yield 3-aminopyrroles. Pyrroles which contain a sulfur atom at C-3 can be prepared from the acids by employing the Hunsdiecker procedure to give the 3-bromo derivatives. Halogen-metal exchange at low temperature with an alkyl lithium reagent can afford the 3-lithio derivative which can be quenched with a variety of electrophiles, such as  $S_8$  to afford thiols directly or  $Cu(SCN)_2$  to afford a thiocyanate which can be reduced with sodium borohydride. The thiols can further be oxidized to the sulfonic acid derivatives by an oxidant such as  $KMnO_4$ .

Scheme 2



15 In Scheme 3 is shown how to prepare pyrroles in which Q-E is attached to the 3-position. This scheme relies upon the extremely versatile Knorr pyrrole synthesis, which involves

condensation of  $\alpha$ -aminoketones with  $\beta$ -ketoesters. The  $\alpha$ -aminoketones can be prepared from  $\beta$ -ketoesters (Scheme 2) by nitrosation followed by reduction with zinc/acetic acid. Condensation of  $\alpha$ -aminoketones with appropriate  $\beta$ -ketoesters


5 can afford good yields of pyrroles. These intermediates are very versatile and can be converted into pyrroles with a wide variety of substituents with varying substitution patterns. For cases wherein  $R^e$  (Z-A-B precursor) is at the 2-position, acidic hydrolysis can selectively hydrolyze the C-3 ester.

10 Heating should then effect decarboxylation. Hydrolysis of the 2-carboxylic acid can be achieved under basic conditions. Curtius rearrangement of the acid as described previously can afford the amino derivatives. To prepare compounds with a sulfur atom attached to C-2, basic hydrolysis and

15 decarboxylation can afford the C-2 unsubstituted pyrroles. These pyrroles can undergo electrophilic substitution to afford thiols ( $\text{Cu}(\text{SCN})_2$ , then  $\text{NaBH}_4$ ) and sulfonic acids (pyridine  $\text{SO}_3$  complex or chlorosulfonic acid). The  $R^{1a}$  group contained in Formula I can be derived either from the

20 remaining ester or from  $R^f$ . Alternatively, the thiol and sulfonic acid derivatives can also be derived from the C-2 acids by manipulation of the carboxylic acid group as described previously.

Scheme 3




- 5 In Scheme 4 is shown how to prepare pyrroles in which Q-E is attached to the 3-position. Cyclization of  $\alpha$ -aminoketones as described previously with  $\beta$ -ketoesters can afford pyrroles. Hydrolysis under basic conditions can selectively hydrolyze the C-2 ester which upon heating should undergo
- 10 decarboxylation to afford 2-unsubstituted pyrroles. The C-3 ester can then be hydrolyzed under acidic conditions to afford the 3-carboxypyrroles. Curtius rearrangement under conditions described previously can afford the 3-aminopyrroles. The



carboxylic acids can be used to prepare the 3-mercapto and 3-sulfonic acid derivatives. The Hunsdiecker procedure can be used to prepare the 3-bromopyrroles. Halogen metal exchange with *t*-BuLi at low temperature followed by quenching with  
5 copper isocyanate should introduce an isocyanate group at C-3. This intermediate can be reduced with sodium borohydride to afford the 3-mercaptopyrroles. Alternatively, the carboxylic acids can be decarboxylated to afford pyrroles which can be N-protected with a bulky protecting group such as  
10 triisopropylsilyl (TIPS). This bulky group directs electrophilic substitution to C-3 of the pyrrole ring. Thus, reaction with copper isocyanate followed by sodium borohydride reduction and then fluoride induced TIPS deprotection can afford 3-mercaptopyrroles. Sulfonation of N-protected pyrrole  
15 with pyridine sulfur trioxide complex can again be directed to C-3 of the pyrrole to afford, after TIPS deprotection, the 3-sulfonic acids.

Scheme 4




5 Another general method of pyrrole synthesis that can be used to prepare compounds of the present invention is shown in Scheme 5. This approach (Cushman et. al. *J. Org. Chem.* **1996**, 61, 4999) uses N-protected  $\alpha$ -aminoketones and N-protected  $\alpha$ -aminoaldehydes which are readily available from  $\alpha$ -amino acids

10 by initial preparation of the N-methoxy-N-methylamides followed by addition of an alkyl Grignard reagent (to produce ketones) or by reduction with a hydride reducing agent such as lithium aluminum hydride or diisobutylaluminum hydride. These

aldehydes and ketones can be allowed to react with the enolates of additional ketones to afford intermediate aldol addition products which under acidic conditions cyclize to form pyrroles. The reacting partners in this approach can be of wide scope and can be chosen so that one skilled in the art will be able to prepare varied pyrroles.

Scheme 5



10




15

20

Another very general method of pyrrole synthesis useful for preparing compounds of the present invention is the Paal-Knorr reaction shown in Scheme 6. This reaction involves the reacting 1,4-diketones or 1,4-ketoaldehydes with primary amines to afford pyrroles. The starting 1,4-diketones and 1,4-ketoaldehydes can be prepared using standard enolate chemistry or by other procedures which are familiar to those skilled in the art of organic synthesis. The reaction is of wide scope and the starting materials can be chosen so that a variety of pyrroles can be prepared.

Scheme 6



25

In Scheme 7 is shown how the compounds of Schemes 1-6 wherein  $R^e$  is a carboxylic ester group can be converted into compounds containing the Z-A-B residue. For the amide linker (Formula I,  $Z = -CONH-$ ), when  $R^e =$  carboalkoxy, it can be hydrolyzed to the acid under either basic or acidic conditions depending on the substitution pattern, as described previously. Formation of the acid chloride with thionyl chloride followed by the addition of an appropriate amine  $H_2N$ -A-B can afford the amide-linked compounds. Alternatively, the acid can be combined with amine  $H_2N$ -A-B in the presence of a suitable peptide coupling agent, such as BOP-Cl, HBTU or DCC. In another method the ester can be directly coupled with an aluminum reagent, prepared by the addition of trimethylaluminum to the amine  $H_2N$ -A-B.


To form ether- or thioether-linked compounds of Formula I ( $Z = -CH_2O-$ ,  $-CH_2S-$ ) the acid can be reduced to the alcohol. Preferred procedures for this transformation are reduction with borane THF complex, or a procedure involving the reduction of the mixed anhydride with sodium borohydride (IBCF=isobutyl chloroformate and NMM=N-methylmorpholine). Completion of the ether and thioether linked compounds of Formula I can readily be accomplished by the Mitsunobu protocol with an appropriate phenol, thiophenol or hydroxy- or mercaptoheterocycle  $HX$ -A-B ( $X = O, S$ ) (Formula I,  $A =$  aryl or heteroaryl). Other ethers or thioethers ( $X = O, S$ ) can be prepared following initial conversion of the alcohol to a suitable leaving group, such as tosylate. Where  $X = S$ , thioethers can be further oxidized to prepare the sulfones (Formula I,  $Z = -CH_2SO_2-$ ).

To prepare the amine-linked compounds of Formula I ( $Z = -CH_2NH-$ ) the alcohol can be oxidized to the aldehyde by a number of procedures, two preferred methods of which are the Swern oxidation and oxidation with pyridinium chlorochromate (PCC). Alternatively, the aldehyde may be directly prepared by direct formylation of the pyrrole ring by the Vilsmeier-Haack procedure in certain cases, as described in previous schemes. Reductive amination of the aldehyde.

with an appropriate amine  $\text{H}_2\text{N-A-B}$  and sodium cyanoborohydride can then afford the amine linked compounds.

- The aldehyde also can be used to prepare the ketone-linked compounds of Formula I ( $\text{Z} = -\text{COCH}_2-$ ). Treatment with an
- 5 organometallic species can afford the alcohol. The organometallic species (wherein  $\text{M} = \text{magnesium or zinc}$ ) can preferably be prepared from the corresponding halide by treatment with metallic magnesium or zinc. These reagents should readily react with aldehydes to afford alcohols.
- 10 Oxidation of the alcohol by any of a number of procedures, such as the Swern oxidation or PCC oxidation, can afford the ketones-linked compounds.

Scheme 7




- 5 Additional compounds of Formula I in which the linking group m/z contains a nitrogen atom attached to ring M can be prepared by the procedures described in Scheme 8. The amines can be converted to sulfonamides (Formula I, m/z-NHSO<sub>2</sub>-) by treatment with an appropriate sulfonyl chloride B-A-SO<sub>2</sub>Cl in the presence of a base such as triethylamine. The amines can be converted into amides (Formula I, Z = -NHCO-) by treatment with an appropriate acid chloride Cl-CO-A-B in the presence of a base or by treatment with an appropriate carboxylic acid HO-
- 10

CO-A-B in the presence of a suitable peptide coupling agent, such as DCC, HBTU or BOP. The amines can also be converted into amine-linked compounds (Formula I,  $Z = -NHCH_2-$ ) by reductive amination with an appropriate aldehyde OHC-A-B.


5

Scheme 8




- 10 Additional compounds of Formula I in which the linking group Z contains a sulfur atom attached to ring M can be prepared by the procedures described in Scheme 9. Treatment of sulfonic acids with phosphorous pentachloride followed by treatment with an appropriate amine  $H_2N-A-B$  can afford
- 15 sulfonamide-linked compounds (Formula I,  $Z = -SO_2NH-$ ). The thiols can be alkylated with a suitable alkylating reagent in the presence of a base to afford thioethers (Formula I,  $Z = -SCH_2-$ ). These compounds can be further oxidized by a variety of reagents to afford the sulfone-linked compounds (Formula I,
- 20  $Z = -SO_2CH_2-$ ).

Scheme 9



- 5 Compounds of Formula I wherein ring M is an imidazole can be formed using procedures described in Schemes 10-16. N-Substituted imidazole derivatives can be made by the general procedure shown in Scheme 10, wherein V' is either V or a precursor of  $(\text{CH}_2)_n\text{V}$ , V is nitro, amino, thio, hydroxy, sulfonic acid, sulfonic ester, sulfonyl chloride, ester, acid, or
- 10 halide, n is 0 and 1, and PG is either a hydrogen or a protecting group. Substitution can be achieved by coupling an imidazole with a halogen containing fragment  $\text{Q-E-G-Hal}$  in the presence of a catalyst, such as base,  $\text{Cu/CuBr/base}$ , or
- 15  $\text{Pd/base}$ , followed by conversion of V' to  $(\text{CH}_2)_n\text{V}$ . Then, Q can be converted to D, and finally V can be converted to  $-\text{Z-A-B}$  following the procedures outlined in Schemes 7-9. Alternatively, V can be converted to  $\text{Z-A-B}$  followed by deprotection of N. This product can then be coupled as before
- 20 to obtain the desired imidazole.


Scheme 10





One way to make amidino-phenyl-imidazole derivatives is shown in Scheme 11. 4-Imidazole carboxylic acid can be treated with thionyl chloride and then coupled with H<sub>2</sub>N-A-B in the presence of a base and then be heated with 3-fluorobenzonitrile in the presence of a base. The Pinner reaction using standard procedures known to those of skill in the art can be used to form the amidino group.


Scheme 11



1,2-Disubstituted and 1,5-disubstituted imidazole derivatives can be made by the general procedures described in Scheme 12, wherein R<sup>1b</sup> is either a hydrogen or an alkyl group and U is aldehyde, ester, acid, amide, amino, thiol, hydroxy, sulfonic acid, sulfonic ester, sulfonyl chloride, or methylene halide. Step a involves coupling in the presence of a catalyst, such as base, Cu/CuBr/base, or Pd/base. When R<sup>1b</sup> is a hydrogen, it can be deprotonated with a lithium base and trapped by formate, formamide, carbon dioxide, sulfonyl chloride (sulfur dioxide and then chlorine), or isocyanate to give 1,2-disubstituted imidazoles (Route b1). Also, in Route b1 when R<sup>1b</sup> is CH<sub>3</sub>, it can be oxidized with SeO<sub>2</sub>, MnO<sub>2</sub>, NaIO<sub>4</sub>/cat. RhCl<sub>3</sub>, or NBS to form U. When R<sup>1b</sup> is hydrogen, sequential deprotonation and quenching with a lithium base and trimethylsilyl chloride, followed by a second deprotonation with a lithium base and quenching with formate, formamide,


carbon dioxide, sulfonyl chloride (sulfur dioxide and then chlorine), or isocyanate can afford 1,5-disubstituted imidazoles (Route b2). When R<sup>1b</sup> is not hydrogen, the procedure of Route b2 can again be used to form 1,5-disubstituted imidazoles (Route b3).

Scheme 12



A preferred way of making 1,2-disubstituted and 1,5-disubstituted imidazole derivatives is shown in Scheme 13. Imidazole can be heated with 3-fluorobenzonitrile in the presence of a base. The coupled product can then be treated with an alkyl lithium base and quenched with ClCO<sub>2</sub>Me to give the 1,2-disubstituted compound. Further treatment with a solution prepared of H<sub>2</sub>N-A-B in trimethylaluminum can give the amide, which can be further modified via the Pinner reaction to form the desired compound. The 1,5-disubstituted compounds can be made using the same procedure, except that the initial anion is protected and a second anion is formed which is then quenched as noted above. Further modifications can follow the same procedures as the 1,2-disubstituted compounds.

Scheme 13




5

Another way of making 1,2-disubstituted imidazole derivatives is described in Scheme 14. By reacting an N-substituted imidazole with a cyanate, the amide can be obtained. This amide can then be coupled with group B as will be described later.

10

Scheme 14




15

Another means of making 1,5-disubstituted imidazole derivatives is described in Scheme 15. Alkylation with 2-bromoethylacetate and subsequent reaction with Gold's reagent in the presence of a base, such as  $NaOMe$ , or  $LDA$ , can form

ester substituted imidazoles which can be further modified as previously described.


Scheme 15



A general procedure to make 2,4,5-trisubstituted or 4,5-disubstituted imidazole derivatives is shown in Scheme 16.


10 After metal halogen exchange of the Q-E-G fragment, it can be reacted with the amide shown, brominated with NBS and cyclized with excess  $\text{NH}_3$  and  $\text{R}^1\text{aCO}_2\text{H}$  to afford an imidazole. This can then be modified as before.

Scheme 16



20 A general procedure to make 4,5-disubstituted triazole derivatives is described in Scheme 17. Ethyl propiolate can be substituted in the presence of  $\text{CuI/Pd}$  and then reacted with  $\text{NaN}_3$  to form a triazole. The triazole can be converted as described previously.


Scheme 17



- 5        The tetrazole compounds of the present invention where Z is -CONH- can be prepared as exemplified in Scheme 18. An appropriately substituted amine can be acylated with ethyl oxalyl chloride. The resulting amide can be converted to the tetrazole either by the methods described by Duncia (*J. Org. Chem.* 1991, 2395-2400) or Thomas (*Synthesis* 1993, 767-768).
- 10      The amide can be converted to the iminoyl chloride first and then reacted with  $NaN_3$  to form the 5-carboethoxytetrazole (*J. Org. Chem.* 1993, 58, 32-35 and *Bioorg. & Med. Chem. Lett.* 1996, 6, 1015-1020). The 5-carboethoxytetrazole can then be
- 15      further modified as described in Scheme 7.

- The tetrazole compounds of the present invention where Z is -CO- can also be prepared via iminoyl chloride (*Chem. Ber.* 1961, 94, 1116 and *J. Org. Chem.* 1976, 41, 1073) using an appropriately substituted acyl chloride as starting material.
- 20      The ketone-linker can be reduced to compounds wherein Z is alkyl.


Scheme 18



5 The methods described in Scheme 18 can also be used to synthesize compounds where the E-Q is linked to the carbon atom of the tetrazole as shown in Scheme 19. The 5-substituted tetrazole can then be alkylated or acylated to give the desired products.

10

Scheme 19



15 The tetrazole compounds of the present invention wherein Z is -SO<sub>2</sub>NH-, -S-, -S(O)-, SO<sub>2</sub>- can be prepared from the thiol prepared as shown in Scheme 20. Appropriately substituted thioisocyanate can be reacted with sodium azide to give the 5-thiotetrazole (*J. Org. Chem.* 1967, 32, 3580-3592). The thio-


20 compound can be modified as described in Scheme 9.

The tetrazole compounds of the present invention wherein Z is -O- can be prepared via the same method described in

Scheme 20 by using appropriately substituted isocyanate as the starting material. The hydroxy compound can be modified similarly to the thiols described in Scheme 9.

5


Scheme 20



The tetrazole compounds of the present invention wherein Z is -NH-, -NHCO-, -NHSO<sub>2</sub>- can be prepared from 5-aminotetrazole, which can be prepared by Smiles Rearrangement as shown in Scheme 21. The thio-compound prepared as described in Scheme 20 can be alkylated with 2-chloroacetamide. The resulting compound can then be refluxed in ethanolic sodium hydroxide to give the corresponding 5-amino-tetrazole (Chem. Pharm. Bull. 1991, 39, 3331-3334). The resulting 5-amino-tetrazole can then be alkylated or acylated to form the desired products.

20


Scheme 21



Pyrazoles of Formula I (such as those described in Scheme 22) can be prepared by the condensation of an appropriately substituted hydrazine with a variety of diketo esters. Condensations of this type typically afford a mixture of pyrazole regioisomers which can be effectively separated via silica gel column chromatography. The esters can be converted to Z-A-B as previously described.

Alternatively, if in Scheme 22, the starting diketone contains CH<sub>3</sub> in place of CO<sub>2</sub>Et, then the resulting methyl pyrazole can be separated and oxidized as in Route b1 in Scheme 12 to form the pyrazole carboxylic acid.

Scheme 22




5

When ketoimides are used for condensations with hydrazines the corresponding pyrazole amino esters are obtained (Scheme 23). Conversion of these intermediates to the final compounds of formula I can then be accomplished by the protection of the amino functionality with a suitable protecting group or by derivatization (e.g. sulfonamide) and then modifying the ester as previously noted.

10

Scheme 23



15


As shown in Scheme 24, pyrazoles wherein the 4-position is substituted can be prepared by bromination (bromine or NBS in either dichloromethane or acetic acid) of the initial pyrazole. Conversion of 4-bromo-pyrazole to 4-carboxylic acid pyrazole can be accomplished by a number of methods commonly known to those in the art of organic synthesis. Further manipulations as previously described can afford pyrazoles of the present invention.

20

25




Scheme 24



Pyrazoles can also be prepared according to method described in Scheme 25. The bromo-pyrazoles are formed as in Scheme 24. QE can then be coupled using palladium catalysed Suzuki cross-coupling methodology. Further modification is achieved as previously described.

Scheme 25




5-substituted phenylpyrazoles can be prepared by the method shown in Scheme 26. Conversion of the 5-hydroxy pyrazole to its triflate (triflic anhydride, lutidine in dichloromethane) or bromide (POBr<sub>3</sub>) followed by palladium Suzuki cross-coupling with an appropriately substituted phenylboronic acid should then afford 5-substituted pyrazoles. Conversion of this intermediate to the 4-bromo derivative

followed by its carbonylation as described in Scheme 24 should then afford the appropriate ester which can be further afford the compounds of formula I.

5


Scheme 26



1-Substituted-1,2,3-triazoles of the present invention can be prepared by the treatment of an appropriately substituted azide with a variety of dipolarophiles (*Tetrahedron* 1971, 27, 845 and *J. Amer. Chem. Soc.* 1951, 73, 1207) as shown in Scheme 27. Typically a mixture of regioisomers are obtained which can be easily separated and elaborated to the triazole carboxylic acids. Further transformations as previously described can then afford the compounds of the present invention.


20

Scheme 27




1,2,4-Triazoles of the present invention can be obtained by the methodology of Huisgen et al (*Liebigs Ann. Chem.* 1962, 653, 105) by the cycloaddition of nitriliminium species (derived from the treatment of triethylamine and chloro  
 5 hydrazone) and an appropriate nitrile dipolarophile (Scheme 28). This methodology provides a wide variety of 1,2,4 triazoles with a varied substitution pattern at the 1, 3, and 5 positions.

Scheme 28




1,2,4-Triazoles can also be prepared by the methodology  
 15 of Zecchi et al (*Synthesis* 1986, 9, 772) by an aza Wittig condensation (Scheme 29).

Scheme 29




1,2,4-Triazoles wherein the -E-D(Q) substituent is at the 5-position of the triazole can be obtained as shown in Scheme  
 30.

Scheme 30



1,3,4-Triazoles of the present invention can be obtained via the methodology of Moderhack et al (*J. Prakt. Chem.* **1996**, 338, 169). As shown in Scheme 31, this reaction involves the condensation of a carbazide with an appropriately substituted commercially available thioisocyanate to form the cyclic thiourea derivative. Alkylation or nucleophilic displacement reactions on the thiono-urea intermediate can then afford a thio-alkyl or aryl intermediate which can be hydrolysed, oxidized and decarboxylated to the 5-H 2-thio-triazole intermediate which can be converted to the compounds of the present invention. Alternatively the thiono-urea intermediate can be oxidized directly to the 2-H triazole which can then be converted to the ester and modified as previously described. The thiono-urea intermediate can also be oxidized to the sulfonyl chloride by methods shown previously.

Scheme 31




The imidazole core shown in Scheme 32 can be prepared by the condensation of 3-cyanoaniline with n-butyglyoxylate to afford the imine which can then be treated with TosylMIC in basic methanol to afford the desired imidazole compound.

- 5 Coupling of the ester under standard conditions then affords a variety of analogs which then can be further manipulated to afford e.g. the benzylamine or the benzamidines.

Scheme 32

10




- Compounds of the present invention wherein AB is a biphenylamine or similar amine may be prepared as shown in Scheme 33. 4-Bromoaniline can be protected as Boc-derivative and coupled to a phenylboronic acid under Suzuki conditions (Bioorg. Med. Chem. Lett. 1994, 189). Deprotection with TFA provides the aminobiphenyl compound. Other similar amines wherein A and/or B are heterocycles can be prepared by the same method using appropriately substituted boronic acids and arylbromide. The bromoaniline can also be linked to the core

ring structures first as described above, and then undergo a Suzuki reaction to give the desired product.

Scheme 33


5



Compounds of the present invention wherein A-B is A-X-Y can be prepared like the piperazine derivative shown in Scheme

10 34.

Scheme 34




15

Scheme 35 shows how one can couple cyclic groups wherein X=NH, O, or S.

Scheme 35

20



When B is defined as X-Y, the following description applies. Groups A and B are available either through commercial sources, known in the literature or readily

synthesized by the adaptation of standard procedures known to practitioners skilled in the art of organic synthesis. The required reactive functional groups appended to analogs of A and B are also available either through commercial sources, known in the literature or readily synthesized by the adaptation of standard procedures known to practitioners skilled in the art of organic synthesis. In the tables that follow the chemistry required to effect the coupling of A to B is outlined.

10

**Table A: Preparation of Amide, Ester, Urea, Sulfonamide and Sulfamide linkages between A and B.**

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	A-NHR <sup>2</sup> as a substituent	ClC(O)-Y	A-NR <sup>2</sup> -C(O)-Y
2	a secondary NH as part of a ring or chain	ClC(O)-Y	A-C(O)-Y
3	A-OH as a substituent	ClC(O)-Y	A-O-C(O)-Y
4	A-NHR <sup>2</sup> as a substituent	ClC(O)-CR <sup>2</sup> R <sup>2a</sup> -Y	A-NR <sup>2</sup> -C(O)-CR <sup>2</sup> R <sup>2a</sup> -Y
5	a secondary NH as part of a ring or chain	ClC(O)-CR <sup>2</sup> R <sup>2a</sup> -Y	A-C(O)-CR <sup>2</sup> R <sup>2a</sup> -Y
6	A-OH as a substituent	ClC(O)-CR <sup>2</sup> R <sup>2a</sup> -Y	A-O-C(O)-CR <sup>2</sup> R <sup>2a</sup> -Y
7	A-NHR <sup>3</sup> as a substituent	ClC(O)NR <sup>2</sup> -Y	A-NR <sup>2</sup> -C(O)NR <sup>2</sup> -Y
8	a secondary NH as part of a ring or chain	ClC(O)NR <sup>2</sup> -Y	A-C(O)NR <sup>2</sup> -Y
9	A-OH as a substituent	ClC(O)NR <sup>2</sup> -Y	A-O-C(O)NR <sup>2</sup> -Y

10.	A-NHR <sup>2</sup> as a substituent	ClSO <sub>2</sub> -Y	A-NR <sup>2</sup> -SO <sub>2</sub> -Y
11	a secondary NH as part of a ring or chain	ClSO <sub>2</sub> -Y	A-SO <sub>2</sub> -Y
12	A-NHR <sup>2</sup> as a substituent	ClSO <sub>2</sub> -CR <sup>2</sup> R <sup>2a</sup> -Y	A-NR <sup>2</sup> -SO <sub>2</sub> -CR <sup>2</sup> R <sup>2a</sup> -Y
13	a secondary NH as part of a ring or chain	ClSO <sub>2</sub> -CR <sup>2</sup> R <sup>2a</sup> -Y	A-SO <sub>2</sub> -CR <sup>2</sup> R <sup>2a</sup> -Y
14	A-NHR <sup>2</sup> as a substituent	ClSO <sub>2</sub> -NR <sup>2</sup> -Y	A-NR <sup>2</sup> -SO <sub>2</sub> -NR <sup>2</sup> -Y
15	a secondary NH as part of a ring or chain	ClSO <sub>2</sub> -NR <sup>2</sup> -Y	A-SO <sub>2</sub> -NR <sup>2</sup> -Y
16	A-C(O)Cl	HO-Y as a substituent	A-C(O)-O-Y
17	A-C(O)Cl	NHR <sup>2</sup> -Y as a substituent	A-C(O)-NR <sup>2</sup> -Y
18	A-C(O)Cl	a secondary NH as part of a ring or chain	A-C(O)-Y
19	A-CR <sup>2</sup> R <sup>2a</sup> C(O)Cl	HO-Y as a substituent	A-CR <sup>2</sup> R <sup>2a</sup> C(O)-O-Y
20	A-CR <sup>2</sup> R <sup>2a</sup> C(O)Cl	NHR <sup>2</sup> -Y as a substituent	A-CR <sup>2</sup> R <sup>2a</sup> C(O)-NR <sup>2</sup> -Y
21	A-CR <sup>2</sup> R <sup>2a</sup> C(O)Cl	a secondary NH as part of a ring or chain	A-CR <sup>2</sup> R <sup>2a</sup> C(O)-Y
22	A-SO <sub>2</sub> Cl	NHR <sup>2</sup> -Y as a substituent	A-SO <sub>2</sub> -NR <sup>2</sup> -Y
23	A-SO <sub>2</sub> Cl	a secondary NH as part of a ring or chain	A-SO <sub>2</sub> -Y
24	A-CR <sup>2</sup> R <sup>2a</sup> SO <sub>2</sub> Cl	NHR <sup>2</sup> -Y as a substituent	A-CR <sup>2</sup> R <sup>2a</sup> SO <sub>2</sub> -NR <sup>2</sup> -Y



25	$A-CR^2R^{2a}SO_2Cl$	a secondary NH as part of a ring or chain	$A-CR^2R^{2a}SO_2-Y$
----	----------------------	-------------------------------------------------	----------------------

The chemistry of Table A can be carried out in aprotic solvents such as a chlorocarbon, pyridine, benzene or toluene, at temperatures ranging from  $-20^\circ C$  to the reflux point of the solvent and with or without a trialkylamine base.

Table B: Preparation of ketone linkages between A and B.

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	$A-C(O)Cl$	$BrMg-Y$	$A-C(O)-Y$
2	$A-CR^2R^{2a}C(O)Cl$	$BrMg-Y$	$A-CR^2R^{2a}C(O)-Y$
3	$A-C(O)Cl$	$BrMgCR^2R^{2a}-Y$	$A-C(O)CR^2R^{2a}-Y$
4	$A-CR^2R^{2a}C(O)Cl$	$BrMgCR^2R^{2a}-Y$	$A-CR^2R^{2a}C(O)CR^2R^{2a}-Y$

The coupling chemistry of Table B can be carried out by a variety of methods. The Grignard reagent required for Y is prepared from a halogen analog of Y in dry ether, dimethoxyethane or tetrahydrofuran at  $0^\circ C$  to the reflux point of the solvent. This Grignard reagent can be reacted directly under very controlled conditions, that is low temperature ( $-20^\circ C$  or lower) and with a large excess of acid chloride or with catalytic or stoichiometric copper bromide-dimethyl sulfide complex in dimethyl sulfide as a solvent or with a variant thereof. Other methods available include transforming the Grignard reagent to the cadmium reagent and coupling according to the procedure of Carson and Prout (Org. Syn. Col. Vol. 3 (1955) 601) or a coupling mediated by  $Fe(acac)_3$  according to Fiandanese et al. (Tetrahedron Lett., (1984) 4805), or a coupling mediated by manganese (II) catalysis (Cahiez and Laboue, Tetrahedron Lett., 33(31), (1992) 4437).

**Table C: Preparation of ether and thioether linkages between A and B**

Rxn. No.	if A contains :	then the reactive substituent of Y is :	to give the following product A-X-Y :
1	A-OH	Br-Y	A-O-Y
2	A-CR <sup>2</sup> R <sup>2a</sup> -OH	Br-Y	A-CR <sup>2</sup> R <sup>2a</sup> O-Y
3	A-OH	Br-CR <sup>2</sup> R <sup>2a</sup> -Y	A-OCR <sup>2</sup> R <sup>2a</sup> -Y
4	A-SH	Br-Y	A-S-Y
5	A-CR <sup>2</sup> R <sup>2a</sup> -SH	Br-Y	A-CR <sup>2</sup> R <sup>2a</sup> S-Y
6	A-SH	Br-CR <sup>2</sup> R <sup>2a</sup> -Y	A-SCR <sup>2</sup> R <sup>2a</sup> -Y

5 The ether and thioether linkages of Table C can be prepared by reacting the two components in a polar aprotic solvent such as acetone, dimethylformamide or dimethylsulfoxide in the presence of a base such as potassium carbonate, sodium hydride or potassium t-butoxide at temperature ranging from ambient temperature to the reflux point of the solvent used.

**Table D: Preparation of -SO- and -SO<sub>2</sub>- linkages from thioethers of Table 3.**

Rxn. No.	if the starting material is :	and it is oxidized with Alumina (wet)/ Oxone (Greenhalgh, Synlett, (1992) 235) the product is :	and it is oxidized with m-chloroperbenzoic acid (Sato et al., Chem. Lett. (1992) 381), the product is :
1	A-S-Y	A-S(O)-Y	A-SO <sub>2</sub> -Y
2	A-CR <sup>2</sup> R <sup>2a</sup> S-Y	A-CR <sup>2</sup> R <sup>2a</sup> S(O)-Y	A-CR <sup>2</sup> R <sup>2a</sup> SO <sub>2</sub> -Y
3	A-SCR <sup>2</sup> R <sup>2a</sup> -Y	A-S(O)CR <sup>2</sup> R <sup>2a</sup> -Y	A-SO <sub>2</sub> CR <sup>2</sup> R <sup>2a</sup> -Y

15 The thioethers of Table C serve as a convenient starting material for the preparation of the sulfoxide and sulfone analogs of Table D. A combination of wet alumina and oxone can provide a reliable reagent for the oxidation of the

thioether to the sulfoxide while m-chloroperbenzoic acid oxidation will give the sulfone.

Table E: Methods of Preparing Group E

5

Rxn	Q	D is to be	then a transformation that may be used is :
1	-CN	-C(=NH)NH <sub>2</sub>	$\text{E}-\text{C}\equiv\text{N} \xrightarrow[\text{ii) NH}_3\text{OAc, MeOH}]{\text{i) HCl MeOH}} \text{E}-\text{C} \begin{matrix} \text{NH}_2 \\ \text{=NH} \end{matrix}$
2	-CN	-CH <sub>2</sub> NH <sub>2</sub>	$\text{E}-\text{C}\equiv\text{N} \xrightarrow[\text{Et}_2\text{O}]{\text{LiAlH}_4} \text{E}-\text{CH}_2\text{NH}_2$
3	-CO <sub>2</sub> H	-CH <sub>2</sub> NH <sub>2</sub>	$\text{E}-\text{C} \begin{matrix} \text{O} \\ \parallel \\ \text{OH} \end{matrix} \xrightarrow[\text{iv) SnCl}_2, \text{MeOH}]{\begin{matrix} \text{i) iBuOC(O)Cl} \\ \text{NMM, THF} \\ \text{then NaBH}_4, \text{H}_2\text{O/THF} \\ \text{ii) MsCl, Et}_3\text{N, CH}_2\text{Cl}_2 \\ \text{iii) NaN}_3, \text{DMF} \end{matrix}} \text{E}-\text{CH}_2\text{NH}_2$
4	-CO <sub>2</sub> H	-NH <sub>2</sub>	$\text{E}-\text{C} \begin{matrix} \text{O} \\ \parallel \\ \text{OH} \end{matrix} \xrightarrow[\text{iii) HCl, Et}_2\text{O}]{\begin{matrix} \text{i) iBuOC(O)Cl} \\ \text{NMM, THF} \\ \text{then NaN}_3 \text{ and heat} \\ \text{ii) tBuOH, reflux} \end{matrix}} \text{E}-\text{NH}_2$

In Table E several methods of transforming a functional group Q into group D of Formula 1 are shown. While not all possible functional groups for Q and D are listed and the synthetic methods suggested are not comprehensive, Table E is meant to illustrate strategies and transformations available to a practitioner skilled in the art of organic synthesis for preparing compounds of Formula 1. In reaction 1 of Table E the transformation of a nitrile into an amidine by the Pinner methodology is shown; in reaction 2 the direct reduction of a nitrile by a hydride reducing agent to a methylene amine is illustrated. In reaction 3, the utility of a carboxylic acid, which may be readily derived from its ester or a nitrile if necessary, in the preparation of a methylene amine is shown. This synthetic route is exceptionally flexible because of the

several stable intermediates prepared *en route* to the final product. As outlined, formation of an activated analog, such as the mixed anhydride, allows for the mild reduction of the acid to the methylene alcohol, this may in turn be transformed  
5 into a leaving group by sulfonylation or halogenation or protected with a suitable protecting group to be transformed later in the synthesis as the chemistry demands. Once the methylene alcohol is so activated, displacement by an efficient nitrogen nucleophile, such as azide anion, can again  
10 provide another suitably stable analog, -the methylene azide- which may be used as a protected form of the methylene amine or transformed directly into the methylene amine group by reduction. Reaction 4 addresses the problem of appending the amine functionality directly through a bond to group E of  
15 Formula 1. Once again, the carboxylic acid provides a convenient *entre* into this selection for group D. The well-known Curtius rearrangement is illustrated here; an activated acid analog can be used to form an acyl azide which upon thermal decomposition is rearranged to the corresponding  
20 isocyanate. The isocyanate intermediate may then be captured as a stable carbamate by the addition of a suitable alcohol and further heating. This carbamate can be used as a stable protecting group for the amine or cleaved directly to the desired D. Alternatively, it may be convenient to quench the  
25 isocyanate intermediate with water to give the amine directly.

Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

30

#### EXAMPLES

##### Fluoro-methylsulfone Intermediate

4-amino-3-fluoro-2'-methylsulfonyl-[1,1']-biphenyl,  
hydrochloride

35

Part A: Preparation of 4-bromo-N-t-butoxycarbonyl-2-fluoroaniline.

Sodium hydride (1.16 g, 60%, 29 mmol) was added to a 0°C solution of 4-bromo-2-fluoro aniline (5.01 g, 26 mmol) in dry DMF (75 mL). The ice bath was removed and the reaction was stirred at room temperature for 1 h. Di-*t*-butyl dicarbonate (6.33 g, 29 mmol) was added, and the reaction was heated at 65°C for 17 h. The reaction was quenched dropwise with H<sub>2</sub>O, then extracted 4 times with H<sub>2</sub>O. The first two aqueous extracts were combined and extracted twice with EtOAc. The organic extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and evaporated. The crude product was taken up in a mixture of CH<sub>2</sub>Cl<sub>2</sub>, CHCl<sub>3</sub>, and EtOAc and filtered to remove a purple impurity, then concentrated and chromatographed on silica gel (30% CH<sub>2</sub>Cl<sub>2</sub> / hexanes) to yield an orange solid (4.76 g, 62%).  
<sup>1</sup>HNMR(DMSO)δ: 9.07 (bs, 1H), 7.57 (td, 1H, J = 8.7, J' = 2.2), 7.49 (dd, 1H, J = 10.2, J' = 2.2), 7.30 (dt, 1H, J = 8.8, J' = 1.1), 1.42 (s, 9H)ppm.

Part B: Preparation of 4-(*t*-butoxycarbonylamino)-3-fluoro-2'-methylthio-[1,1']-biphenyl.

A flask containing a mixture of 4-bromo-*N*-*t*-butoxycarbonyl-2-fluoroaniline (6.44 g, 22 mmol), 2-(methylthio)phenylboronic acid (6.00 g, 36 mmol), aq. sodium carbonate (2.0 M, 36 mL, 72 mmol), tetrabutylammonium bromide (360 mg, 1.1 mmol), and bis(triphenylphosphine)palladium(II) chloride in benzene (180 mL) was evacuated twice under brief high vacuum, filled with argon, and heated at reflux for 5 h. After cooling to room temperature, the layers were separated, and the aqueous layer was extracted with EtOAc. The organic extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (0-30% EtOAc / hexanes) to yield the desired product (6.50 g, 88%).  
<sup>1</sup>HNMR(CHCl<sub>3</sub>)δ: 8.14 (bt, 1H, J = 8.1), 7.30 (m, 2H), 7.17 (m, 4H), 6.75 (bs, 1H), 2.37 (s, 3H), 1.54 (s, 9H)ppm.

Part C: Preparation of 4-(*t*-butoxycarbonylamino)-3-fluoro-2'-methylsulfonyl-[1,1']-biphenyl.

4-(*t*-Butoxycarbonylamino)-3-fluoro-2'-methylthio-[1,1']-biphenyl (6.50 g, 19.5 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (150 mL) and cooled to 0°C. *m*-CPBA (14.8 g, 57-86%) was added and the reaction stirred at room temperature for 3 h. The reaction was extracted with sat. sodium sulfite, and the aqueous layer was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (20-30% EtOAc/hexanes) to yield the desired product (6.92 g, 97%). <sup>1</sup>HNMR(CHCl<sub>3</sub>)δ: 8.22 (dd, 2H, J = 7.7, J' = 1.5), 7.64 (td, 1H, J = 7.3, J' = 1.5), 7.56 (td, 1H, J = 7.7, J' = 1.5), 7.35 (dd, 1H, J = 7.3, J' = 1.5), 7.30 (dd, J = 11.7, J' = 2.2), 7.17 (d, 1H, J = 8.8), 6.82 (bs, 1H), 2.69 (s, 3H), 1.55 (s, 9H)ppm.

Part D: Preparation of 4-amino-3-fluoro-2'-methylsulfonyl-[1,1']-biphenyl, hydrochloride.

4-(*t*-Butoxycarbonylamino)-3-fluoro-2'-methylsulfonyl-[1,1']-biphenyl (1.04 g, 2.8 mmol) was dissolved in HCl/dioxane (4.0 M, 10 mL) and stirred 19 h. A solid was triturated with Et<sub>2</sub>O and filtered to yield a white solid (813 mg, 95%). <sup>1</sup>HNMR(DMSO)δ: 8.03 (dd, 1H, J = 8.0, J' = 1.4), 7.69 (td, 1H, J = 7.7, J' = 1.1), 7.59 (t, 1H, J = 7.4), 7.36 (d, 1H, J = 7.3), 7.12 (d, 1H, J = 12.4), 6.94 (m, 2H), 2.78 (s, 3H)ppm.

#### Examples 1 and 2

1-(3-amidinophenyl)-2-[[[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]pyrrole, trifluoroacetic acid salt (Example 1) and 1-(3-amidinophenyl)-2-[[[(2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]pyrrole, trifluoroacetic acid salt (Example 2)

Part A. Preparation of 1-(3-cyanophenyl) pyrrole.

3-aminobenzonitrile (47.45 g, 0.401 mol) and 58.4 g (.441 mol, 59.5 ml) of 2,5-dimethoxytetrahydrofuran were dissolved

in 200 ml of acetic acid and heated to reflux over night. The solution was allowed to cool to room temperature and diluted with 250 ml of ethylacetate and was washed 3 times with brine (200ml) and then by a solution of saturated aq sodium carbonate (200 ml). The organics were dried over magnesium sulfate and filtered through a plug of silica gel. The volatiles were removed in vacuo and the residue was recrystallized from methanol to yield the title compound as a beige solid (62.82 g, 93%) MS (H<sub>2</sub>O-CI) 169 (M+H)<sup>+</sup>.

Part B. Preparation of 1-(3-cyanophenyl) pyrrole-2-carboxaldehyde.

Phosphorous oxychloride, over the course of 15 minutes, was added to dimethylformamide (14.02 g, 191.84 mmol, 14.1 ml) at 0°C. The mixture was warmed to room temperature and stirred for 15 minutes; the solution was again cooled to 0°C followed by the addition of 100 ml of 1,2 dichloroethane. A solution of 1-(3-cyanophenyl) pyrrole (29.33 g, 191.84 mmol) in 250 ml of 1,2 dichloroethane was added slowly via an addition funnel and the mixture heated to reflux for 15 minutes. The solution was cooled to room temperature and 86.55 g (1.05 mol) of sodium acetate was added and the solution heated to reflux for 15 minutes. The solution was diluted with 250 ml of ethyl acetate and the organics washed with brine then saturated aq sodium carbonate (250 ml). The organics were dried over magnesium sulfate, filtered through a plug of silica gel and the volatiles removed in vacuo. The product was recrystallized from ethyl acetate to yield 28.4 g (83%) of the title compound. MS (NH<sub>3</sub>-CI) 214 (M+NH<sub>4</sub>)<sup>+</sup>.

Part C. Preparation of 1-(3-cyanophenyl) pyrrole-2-carboxylic acid.

To a cooled (0°C) solution of 1-(3-cyanophenyl) pyrrole-2-carboxaldehyde (5.14g, 26.20 mmol) in 300 ml of 1:1 acetone/water was added potassium permanganate (12.42 g, 78.60 mmol) over a period of 15 minutes and the reaction was allowed

to warm to room temperature. After consumption of the starting material, 10.90 g (104.8 mmol) of sodium bisulfite was added and the solution made acidic with 10% HCl. The solution was filtered through a plug of celite, diluted with ethyl acetate and washed with 200 ml of brine. The organics were dried over magnesium sulfate, filtered and dried *in vacuo*. The organics were recrystallized from methanol to yield the title compound (4.11 g, 74%) as a pale white solid. MS (ESI) 211.2 (M-H)<sup>-</sup>.

10

Part D. Preparation of 1-(3-cyanophenyl)-2-[[2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] pyrrole.

To a solution of 1-(3-cyanophenyl) pyrrole-2-carboxylic acid (2.77 g, 13.05 mmol) in 50 ml of anhydrous DMF was added triethylamine (1.98 ml, 19.58 mmol), benzotriazole-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (8.66 g, 19.58 mmol) and (2'-*N*-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)-amine (6.03 g, 19.84 mmol) and heated to 50° C overnight. The solution was diluted with ethyl acetate and washed repeatedly with brine. The organic layer was dried over magnesium sulfate and the volatiles removed *in vacuo*. The residue was subjected to flash chromatography purification with 3:2 hexane/ethyl acetate and the volatiles removed *in vacuo* to yield 1.9 g (29%) of the title compound. MS (NH<sub>3</sub>-CI) 516 (M+NH<sub>4</sub>)<sup>+</sup>.

Part E. Preparation of 1-(3-amidinophenyl)-2-[[2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] pyrrole, trifluoroacetic acid salt (Example 1) and 1-(3-amidinophenyl)-2-[[2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] pyrrole, trifluoroacetic acid salt (Example 2).

1-(3-Cyanophenyl)-2-[[2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] (0.37 g, 0.74 mmol) of pyrrole was added to a solution of 60 ml of anhydrous methyl acetate and anhydrous methanol (0.30 ml, 7.4 mmol) and cooled in an ice water bath. Gaseous HCl was bubbled in for 15 minutes, the



solution stoppered and allowed to stir overnight at room temperature. The volatiles were removed in vacuo. The residue was dried under high vacuum for 1 hr. The residue was then dissolved in 100 ml of anhydrous methanol and combined with .43 g (4.45 mmol) of ammonium carbonate and stirred overnight at room temperature. The volatiles were removed in vacuo and the residue was purified by preparative HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) to yield 1-(3-amidinophenyl)-2-[[[2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] pyrrole, trifluoroacetic acid salt (Example 1) as a white solid following lyophilization. MS (ESI) 460.3 (M+H)<sup>+</sup>; also isolated was 1-(3-amidinophenyl)-2-[[[2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl] pyrrole, trifluoroacetic acid salt (Example 2). MS (ESI) 516.4 (M+H)<sup>+</sup>.

### Example 3

1-(3-amidinophenyl)-2-[[[2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]-4-bromopyrrole, trifluoroacetic acid salt

Part A. Preparation of 1-(3-cyanophenyl)-2-formyl-4-bromopyrrole.

1-(3-Cyanophenyl) pyrrole-2-carboxaldehyde from Example 1, Part B (6.06 g, 30.89 mmol) was combined with 6.60 g (37.06 mmol) of N-bromosuccinimide in 150 ml of anhydrous THF and stirred at room temperature overnight. The residue was heated in CCl<sub>4</sub> and filtered. The residue was then dissolved in CHCl<sub>3</sub>/EtOAc, filtered through a silica gel plug and the volatiles removed. The residue was recrystallized from ethyl acetate to yield the title compound as a light brown solid (4.49 g, 53%). MS (NH<sub>3</sub>-CI) 292 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B. Preparation of 1-(3-amidinophenyl)-2-[[2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]-4-bromopyrrole, trifluoroacetic acid salt.

5        Following the procedures described in Example 1, Parts C-E, 1-(3-cyanophenyl)-2-formyl-4-bromo-pyrrole was converted into the title compound as a white powder following HPLC purification. MS (ESI) 538.2 (M+H)<sup>+</sup>.

10

#### Example 4

1-(3-amidinophenyl)-2-[[5-(2'-aminosulfonylphenyl)-1-yl)pyridin-2-yl]-aminocarbonyl]pyrrole, trifluoroacetic acid salt

15        Part A. Preparation of 1-(3-cyanophenyl)-2-[[5-(2'-tert-butylaminosulfonylphenyl)-1-yl)pyridin-2-yl]aminocarbonyl]pyrrole.

1-(3-Cyanophenyl) pyrrole-2-carboxylic acid from Example 1, Part C (1.00 g, 4.7 mmol), oxalyl chloride (.61 ml, 7.06 mmol) and 3 drops of DMF were combined at room temperature in 50 ml of anhydrous CH<sub>2</sub>Cl<sub>2</sub> and stirred for 4 hours. The volatiles were removed in vacuo and the residue was dried under high vacuum for 1 hour. The residue was then dissolved in 50 ml of CH<sub>2</sub>Cl<sub>2</sub> followed by the addition of 4-dimethylaminopyridine (1.15 g, 9.4 mmol), the solution stirred at room temperature for 5 minutes followed by the addition of [5-(2'-aminosulfonylphenyl)-1-yl)pyridin-2-yl]-amine (1.44 g, 4.7 mmol) and stirred at room temperature overnight. The solution was filtered through a silica gel plug and the volatiles removed. The residue was purified by flash chromatography (1:2 hexane/EtOAc) to yield 0.84 g (36%) of the title compound as a tan solid. MS (ESI) 500.3 (M+H)<sup>+</sup>.

Part B. Preparation of 1-(3-amidinophenyl)-2-[[5-(2'-aminosulfonylphenyl)-1-yl]pyridin-2-yl]-aminocarbonylpyrrole, trifluoroacetic acid salt.

5        Following the procedures described in Example 1, Part E, 1-(3-cyanophenyl)-2-[[5-(2'-tert-butylaminosulfonylphenyl)-1-yl]pyridin-2-yl]aminocarbonylpyrrole was converted into the title compound as a white powder following HPLC purification. MS (ESI) 461.3 (M+H)<sup>+</sup>.

10

#### Examples 5 and 6

1-Benzyl-3-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example 5) and 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example 6)

15

Part A: Preparation of ethyl 3-(3-cyanophenyl)propiolate.

20        To a solution of ethyl propiolate (25.0 g, 0.25 mol) in 750 mL of tetrahydrofuran at -78° C was added *n*-butyllithium (102 mL of a 2.5 M solution in hexane, 0.25 mol) dropwise. After stirring at the same temperature for 1 h, zinc chloride (104.2 g, 0.76 mol) was added in 900 mL of tetrahydrofuran.

25        The mixture was allowed to gradually warm to room temperature over 1 h. To this solution was added 3-iodobenzonitrile (29.2 g, 0.13 mol) and bis triphenylphosphine palladium (II) chloride (4.56 g, 6.5 mmol) and the resulting mixture was stirred at 50° C overnight. To the mixture was added 150 mL of

30        water and 150 mL of ether and the mixture was filtered through a celite pad. The filtrate was extracted 3 times with ether and the combined extracts were washed with brine, dried (MgSO<sub>4</sub>) and filtered through a thick pad of silica gel. The solvents were removed in vacuo and the residue was recrystallized from

35        hexane ethyl acetate to afford 8.8 g (35%) of the title compound as a tan solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.85 (s, 1H), 7.8 (d, 1H), 7.72 (d, 1H), 7.52 (t, 1H), 4.30 (q, 2H), 1.37 (t, 3H).

Part B: Preparation of 1-benzyl-3-carboethoxy-4-(3-cyanophenyl)- $\Delta^3$ -pyrroline.

5 To a solution of N-benzyl-N-(trimethylsilylmethyl)-aminomethyl methyl ether (12.25 g, 51.2 mmol) in 400 mL of methylene chloride at 0° C was added ethyl 3-(3-cyanophenyl)propiolate (6.79 g, 34.1 mmol) followed by trifluoroacetic acid (0.20 mL, 2.6 mmol). The mixture was  
10 allowed to warm to room temperature and was stirred for 16 h. The reaction mixture was washed with saturated aqueous NaHCO<sub>3</sub> and brine, dried over K<sub>2</sub>CO<sub>3</sub>, filtered through a large pad of silica gel and concentrated in vacuo. The residue was  
15 purified by flash chromatography (elution with 5:1 hexanes/ethyl acetate) to afford 3.2 g (28%) of the title compound. MS (ESI) 333.4 (M+H)+.

Part C: Preparation of 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-cyanophenyl)- $\Delta^3$ -pyrroline.  
20

To a solution of (2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)-amine (1.10 g, 3.6 mmol) in 50 mL of methylene chloride at room temperature was added trimethylaluminum (6.6  
25 mL of a 2.0 M solution in toluene, 13.2 mmol) dropwise. The solution was stirred (30 min) until gas evolution had ceased followed by the addition of 1-benzyl-3-carboethoxy-4-(3-cyanophenyl)- $\Delta^3$ -pyrroline (1.0 g, 3.0 mmol) in 5 mL of methylene chloride. The resulting solution was stirred at 40°C  
30 for 2h, cooled to room temperature and quenched with saturated aq NH<sub>4</sub>Cl. The mixture was diluted with ethyl acetate, washed with water and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 4:1 hexane/ethyl acetate) to afford 0.58 g (34%) of the  
35 title compound. MS (ESI) 591.5 (M+H)+.

Part D: Preparation of 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-cyanophenyl)pyrrole.

5 To a solution of 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-cyanophenyl)- $\Delta^3$ -pyrroline (0.47 g, 0.8 mmol) in 20 mL of benzene was added 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) (0.27 g, 1.2 mmol) and the resulting mixture was stirred at 70° C for 16 h.  
10 The mixture was cooled and filtered and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 5:1 hexane/ethyl acetate) to afford 0.25 g (53%) of the title compound. MS (ESI) 589.6 (M+H)+.

15 Part E: Preparation of 1-benzyl-3-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example 5) and 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example  
20 6).

A solution of 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-cyanophenyl)pyrrole (0.25 g, 0.42 mmol) in 50 mL of anhydrous methanol was cooled  
25 to 0° C. Anhydrous HCl gas was bubbled through the solution for about 30 min (until solution saturated). The flask was sealed and allowed to stand for 16 h at 0° C. The reaction mixture was concentrated *in vacuo*. The resulting solid was dissolved in 20 mL of anhydrous methanol, ammonium carbonate  
30 (0.20 g, 2.1 mmol) was added, and the mixture was allowed to stir at room temperature for 24 h. The reaction mixture was concentrated *in vacuo* and purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) to afford 120 mg (40%) of 1-benzyl-3-[(2'-aminosulfonyl-  
35 [1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example 5) as a white powder following lyophilization. MS (ESI) 550.3 (M+H)+. The preparation also afforded 40 mg (13%) of 1-benzyl-3-[(2'-tert-

butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole, trifluoroacetic acid salt (Example 6) as a white powder following lyophilization. MS (ESI) 606.5 (M+H)+.

5

#### Examples 7 and 8

1-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole (Example 7) and 1-(3-amidinophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole (Example 8)

10

Part A: Preparation of 4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole

15 To a suspension of 4-imidazolecarboxylic acid (168 mg, 1.5 mmol) in CH<sub>3</sub>CN (30 mL) was added thionyl chloride (714 mg, 6 mmol), and the resulting mixture was heated at 80°C for 2 hours. After removal of volatiles, a yellow residue reacted with 4-[(o-SO<sub>2</sub>-t-Bu)-phenyl]aniline (304 mg, 1 mmol) in  
20 pyridine (10 mL) at room temperature for 24 hours.

Evaporation of the pyridine gave a residue which was dissolved in EtOAc and washed with water, brine, and dried over MgSO<sub>4</sub>. Concentration and purification by column chromatography of the crude material provided the title compound (378 mg, 95%  
25 yield). <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 8.10 (dd, J = 7.7 Hz, 1.1 Hz, 1H), 7.79 (d, J = 3.7 Hz, 2H), 7.76 (d, J = 8.4 Hz, 2H), 7.58 (td, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.49 (dd, J = 8.1 Hz, J = 1.5 Hz, 1H), 7.45 (d, J = 8.4 Hz, 2H), 7.34 (dd, J = 7.7 Hz, J = 1.5 Hz, 1H), 1.06 (s, 9H); LRMS: 399.3 (M+H)+.

30

Part B: Preparation of 1-(3-cyanophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole

4-[(2'-tert-Butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole was heated with 3-fluorobenzonitrile (121 mg, 1 mmol) in the presence of K<sub>2</sub>CO<sub>3</sub> in DMF at 100°C for 8 hours to give the title compound in almost  
35 quantitative yield. <sup>1</sup>HNMR(acetone-d<sub>6</sub>) δ: 9.47 (s, 1H), 8.39 (d,

J = 1.5 Hz, 1H), 8.34 (d, J = 1.5 Hz, 1H), 8.25 (d, J = 1.5 Hz, 1H), 8.15-8.10 (m, 1H), 8.02 (d, J = 8.4 Hz, 2H), 7.88-7.79 (m, 2H), 7.65 (td, J = 7.3 Hz, J = 1.5 Hz, 1H), 7.56 (dd, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.50 (d, J = 8.4 Hz, 2H), 7.38 (dd, J = 8.4 Hz, J = 1.5 Hz, 1H), 2.80 (s, 1H), 1.03 (s, 9H); LRMS: 500.1 (M+H)<sup>+</sup>.

Part C: Preparation of 1-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole (Example 7) and 1-(3-amidinophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole (Example 8)

1-(3-Cyanophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole was further subjected to a Pinner reaction by standard procedures to give Examples 7 (309 mg, 62% yield) and 8 (67 mg, 12% yield).

For Example 7: <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ: 8.32 (d, J = 1.4 Hz, 2H), 8.12 (s, 1H), 8.10 (d, J = 8.1 Hz, 1H), 8.08 (dd, J = 7.7 Hz, J = 1.4 Hz, 1H), 7.88-7.81 (m, 2H), 7.78 (d, J = 8.4 Hz, 2H), 7.61 (td, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.52 (td, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.44 (d, J = 8.4 Hz, 2H), 7.35 (dd, J = 7.3 Hz, J = 1.1 Hz, 1H); <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ: 167.59, 162.59, 143.08, 141.63, 139.32, 138.97, 138.68, 137.58, 137.33, 133.72, 132.94, 132.44, 131.62, 131.28, 128.72, 128.66, 128.49, 127.66, 122.94, 122.12, 121.00; ESMS: 461.3 (M+H)<sup>+</sup>; HRMS: 461.1387 (obs.), 461.1396 (calcd.).

For Example 8: <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ: 8.33 (s, 2H), 8.12 (s, 1H), 8.10 (dd, J = 8.4 Hz, J = 1.5 Hz, 1H), 8.05 (dd, J = 8.1 Hz, J = 2.2 Hz, 1H), 7.88-7.80 (m, 3H), 7.79 (d, J = 8.4 Hz, 2H), 7.61 (td, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.53 (td, J = 7.7 Hz, J = 1.2 Hz, 1H), 7.47 (d, J = 8.4 Hz, 2H); 7.35 (dd, J = 7.3 Hz, J = 1.5 Hz, 1H), 1.02 (s, 9H); <sup>13</sup>C NMR (CD<sub>3</sub>OD) δ: 167.58, 162.57, 143.51, 141.65, 139.02, 138.68, 138.68, 137.30, 133.89, 133.05, 132.44, 131.64, 131.52, 128.72, 129.53, 128.77, 128.50, 127.65, 122.96, 122.12, 120.99, 55.06, 30.11; ESMS: 517.4 (M+H)<sup>+</sup>; HRMS: 517.2025 (obs.), 517.2022 (calcd.);

Anal.: (C<sub>27</sub>H<sub>28</sub>N<sub>6</sub>O<sub>3</sub>S<sub>1</sub> + 1.35TFA + 0.17HCl + 0.6H<sub>2</sub>O) C, H, N, S, F, Cl.

### Example 9

#### 5 1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole

##### Part A: Preparation of 1-(3-cyanophenyl)imidazole

10 3-Fluorobenzonitrile (4.84 g, 40 mmol) was heated with imidazole (2.72 g, 40 mmol) in the presence of K<sub>2</sub>CO<sub>3</sub> in DMF at 100°C for 8 hours to give the title compound as a white solid in quantitative yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.89 (s, 1H), 7.70 (d, J = 0.8 Hz, 1H), 7.68-7.58 (m, 3H), 7.30 (d, J = 1.0 Hz, 1H),  
15 7.26 (s, 1H); LRMS: 170 (M+H)<sup>+</sup>.

##### Part B: Preparation of methyl 1-(3-cyanophenyl)imidazol-2-yl carboxylate

20 1-(3-Cyanophenyl)imidazole (1.52 g, 9 mmol) was slowly treated with n-BuLi (1.6 M, 6.3 mL) in THF (60 mL) at -78°C for 40 minutes and was then slowly quenched with chloromethylformate (942 mg, 10 mmol) at this temperature. The resulting mixture was stirred at -78°C and warmed to room  
25 temperature over 2 hours and then poured into water and ethyl acetate. The organic layer was separated and washed with water, brine, and dried over MgSO<sub>4</sub>. After removal of the ethyl acetate the residue was purified by column chromatography with ethyl acetate and methylene chloride (1:1) to afford the title  
30 compound (1.33g, 65%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.80-7.77 (m, 1H), 7.65-7.61 (m, 3H), 7.33 (s, 1H), 7.20 (s, 1H); LRMS: 228 (M+H)<sup>+</sup>.

##### 35 Part C: Preparation of 1-(3-cyanophenyl)-2-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole

To a stirred solution of 4-[(o-SO<sub>2</sub>tBu)-phenyl]aniline (304 mg, 1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was slowly added



trimethylaluminum (2M in hexane, 1 mL) at 0°C and the resulting mixture was warmed to room temperature over 15 minutes. After addition a solution of methyl 1-(3-cyanophenyl)imidazol-2-yl carboxylate in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and the resulting mixture was  
5 refluxed for 2 hours. The mixture was quenched with water, diluted with ethyl acetate and filtered through Celite. The organic layer was separated, washed with water, and brine and dried over MgSO<sub>4</sub>. After removal of the ethyl acetate, a residue was purified by column chromatography with ethyl  
10 acetate and methylene chloride (1:1) to afford the title compound (260 mg, 52%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 9.41 (s, 1H), 8.15 (dd, J = 7.7 Hz, J = 1.5 Hz, 1H), 7.78 (dd, J = 7.3 Hz, J = 1.1 Hz, 1H), 7.74-7.57 (m, 6H), 7.55 (td, J = 7.7 Hz, J = 1.1 Hz, 1H), 7.49 (dd, J = 8.8 Hz, J = 1.8 Hz, 2H),  
15 7.29 (dd, J = 8.1 Hz, J = 1.5 Hz, 1H), 7.28 (d, J = 0.8 Hz, 1H), 7.22 (d, J = 0.8 Hz, 1H), 3.64 (s, 1H), 0.99 (s, 9H); LRMS: 500.1 (M+H)<sup>+</sup>.

Part D: Preparation of 1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole  
20

1-(3-Cyanophenyl)-2-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole was subjected to the Pinner reaction to form the title compound (120 mg, 50%):  
25 <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 8.08 (dd, J = 7.7 Hz, J = 1.1 Hz, 1H), 7.91-7.88 (m, 2H), 7.83 (dd, J = 8.4 Hz, J = 1.5 Hz, 1H), 7.74 (t, J = 8.0 Hz, 1H), 7.65 (d, J = 8.4 Hz, 2H), 7.58 (dd, J = 7.3 Hz, J = 1.1 Hz, 1H), 7.50 (s, 2H), 7.39 (d, J = 8.4 Hz, 2H), 7.32 (s, 1H), 7.30 (dd, J = 7.3 Hz, J = 1.1 Hz, 1H); ESMS: 461  
30 (M+H)<sup>+</sup>.

## Example 10

1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole, trifluoroacetic acid

- 5 Part A: Preparation of ethyl 1-(3-bromophenyl)-3-methyl-pyrazol-5-yl carboxylate and ethyl 1-(3-bromophenyl)-5-methyl-pyrazol-3-yl carboxylate

2-Bromophenylhydrazine hydrochloride (6.5 g, 0.029 mol)  
10 was added in portions to a ethanolic solution of 3-methoxy-trichloroacetylcrotonate (Fischer et. al. *Synthesis* 1991, 83). The reaction mixture was refluxed for 48h cooled and concentrated. The residue was dissolved in ethyl acetate (100 mL) and washed with HCl (1N, 50 mL), brine (50 mL) and dried  
15 (magnesium sulfate). Evaporation afforded an oil which was subjected to silica gel column chromatography (hexane:ethylacetate, 6:1) to afford ethyl 1-(3-bromophenyl)-5-methyl-pyrazol-3-yl carboxylate (3.73 g) and ethyl ethyl 1-(3-bromophenyl)-3-methyl-pyrazol-5-yl carboxylate (3.65 g) as  
20 pure compounds. The pyrazole carboxylate obtained this way were used directly in part B.

Part B: Preparation of ethyl 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylate

25 Ethyl 1-(3-bromophenyl)-3-methyl-pyrazol-5-yl carboxylate (2.3 g) was dissolved in N-methyl-pyrrolidinone (4 mL) and to this solution was added CuCN (1 g). The reaction mixture was refluxed for 2 h then stirred at room temperature overnight.  
30 The mixture was quenched with water (100 mL) and the organics were extracted with ethylacetate (2X100 mL) and dried (magnesium sulfate). Silica gel column chromatography (hexane:ethylacetate, 3:1) then afforded the title compound (0.59 g). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.76 (t, 1H), 7.70 (dd, 1H), 7.58  
35 (t, 1H), 6.86 (s, 1H), 4.3 (q, 2H), 2.36 (s, 3H), 1.31 (t, 3H) ppm; IR (neat), 2230, 1728, 1586, 1540, 1494, 1438, 1298, 1242, 1106, 1046, 760, 682 cm<sup>-1</sup>. Chemical Ionization mass spectrum m/z (rel. intensity) 256 (M+H, 100).

Part C: Preparation of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid

5 Ethyl 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylate (0.55 g) was dissolved in THF (20 mL) and to this was added LiOH (0.5M, 5,6mL). The reaction mixture was stirred at room temperature for 18h then quenched with water (50 mL). The unreacted organics were extracted with ethylacetate (2X50 mL).  
10 The aqueous layer was acidified and extracted with ethylacetate (2X50 mL) dried (magnesium sulfate) and evaporated to afford pure acid.  $^1\text{H NMR}$  (DMSO  $d_6$ )  $\delta$ : 8.02 (t, 1H), 7.91 (d, 1H), 7.82 (dd, 1H), 7.09 (t, 1H), 6.89 (s, 1H), 2.27 (s, 3H) ppm; IR (PEC) 2930, 2232, 1724, 1710, 1540, 1496, 1458, 1276, 1230, 1186, 1146, 1112, 900, 768, 754, 690  $\text{cm}^{-1}$ ;  
15 Chemical ionization mass spectrum  $m/z$  (rel. intensity) 228 (M+H, 100).

20 Part D: Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole

To a dichloromethane solution (20 mL) of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid (0.2 g) was  
25 added oxalyl chloride (0.11 mL). The reaction mixture was stirred at room temperature for 2h then to this solution was added 2-tert-butylsulfonamide-1-biphenyl aniline (0.27 g) and triethylamine (0.5 mL). The reaction mixture was stirred at room temperature for 24h then quenched with water (50 mL) and  
30 the organics were extracted with ethylacetate (2X50 mL), washed with brine (50 mL) and dried (magnesium sulfate). Evaporation afforded an oil which was chromatographed on silica gel column (dichloromethane:MeOH, 9:1) to afford the title compound (0.45g).  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 8.16 (d, 1H), 8.05 (s, 1H), 7.8 (d, 1H), 7.76 (d, 1H), 7.68 (d, 3H), 7.58 (m, 2H), 7.50 (md, 3H), 7.30 (d, 1H), 6.76 (s, 1H), 3.64 (s, 1H), 2.42 (s, 3H), 1.03 (s, 9H) ppm; IR (PEC), 3320, 2976, 2232, 1682, 1592, 1540, 1522, 1488, 1464, 1438, 1368, 1320, 1242, 1152, 1128, 758,

682, 608cm<sup>-1</sup>; Chemical ionization mass spectrum m/z (rel intensity) 458 (M=H, 100).

Part E: Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole, trifluoroacetic acid

1-(3-Cyanophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole (0.39 g) was dissolved in a saturated HCl solution of anhydrous MeOH (20 mL). The reaction mixture was stirred at room temperature for 24h then MeOH was evaporated. The residue was redissolved in MeOH (20 mL) and excess ammonium carbonate added. The reaction mixture was stirred at room temperature for 18 h. MeOH was evaporated and the residue was purified via HPLC to afford the desired compound as its TFA salt (0.15 g). <sup>1</sup>HNMR(DMSO d<sub>6</sub>)δ: 10.66 (s, 1H), 9.44 (s, 1.5H), 9.09 (s, 1.5H), 8.03 (d, 1H), 7.97 (s, 1H), 7.83 (t, 1H), 7.75 (d, 1H), 7.70 (d, 2H), 7.62 (m, 2H), 7.37 (d, 2H), 7.32 (d, 1H), 7.27 (s, 2H), 7.03 (s, 1H), 2.50 (s, 3H) ppm; IR (PEC) 3288, 1704, 1660, 1592, 1526, 1484, 1438, 1322, 1206, 1160, 762, 724cm<sup>-1</sup>; High resolution mass spectrum calcd. for C<sub>24</sub>H<sub>22</sub>N<sub>6</sub>O<sub>3</sub>S 475.155236, found 475.153767.

#### Example 11

1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylamino]-pyrazole, trifluoroacetic acid

Part A. Preparation of 5-amino-1-(3'-cyanophenyl)-3-methylpyrazole.

3-aminocrotonitrile (1 g, 12.2 mmol) and 3-cyanophenyl hydrazine hydrochloride (2 g, 11.8 mmol) were combined and heated to reflux in 1:1 ethanol/acetic acid (20 mL) for 4h. The reaction was concentrated and the residue basified with diluted NaOH and extracted with ethyl acetate. The crude product was purified by flash chromatography on silica gel using hexanes/ ethyl acetate (4:1) as eluent to afford 1.2 g

of a still impure amine. This amine was dissolved in dilute HCl and extracted with ethyl acetate. The aqueous layer was basified with NaOH and extracted with ethyl acetate and dried (MgSO<sub>4</sub>) to afford 0.66 g (28%) of amine; <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.97 (s, 1H), 7.92 (m, 1H), 7.57 (s+d, 2H), 5.51 (s, 1H), 3.75 (s, 2H), 2.23 (s, 3H); MS (H<sub>2</sub>O/GC) m/z 199 (M+H<sup>+</sup>).

Part B. Preparation of 1-(3'-cyanophenyl)-3-methyl-5-((4'-bromophenyl)carbonylamino)pyrazole.

10

To the product of part A (0.66 g, 3.3 mmol) in methylene chloride (20 mL) at 0°C was added 2M trimethylaluminum (8.3 mL, 16.7 mmol) in heptane. The mixture was stirred for 15 minutes and methyl-4-bromobenzoate (0.72 g, 3.3 mmol) was added. The reaction was stirred overnight. The reaction was quenched with 1N HCl and extracted with methylene chloride and dried (Na<sub>2</sub>SO<sub>4</sub>). Recrystallization from methylene chloride/hexanes yielded 0.48 g (45%) of the title compound; <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.86 (s, 1H), 7.78 (d, J=7.69Hz, 1H), 7.67 (d, J=7.69Hz, 1H), 7.63 (m, 4H), 7.60 (m, 1H), 6.52 (s, 1H), 2.36 (s, 3H); MS (ESI) m/z 381.1-383.1 (M+H<sup>+</sup>).

20

Part C. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylamino]-pyrazole, trifluoroacetic acid.

25

A mixture of the above part B amide (0.4 g, 1 mmol), 2-(t-butylsulfonamide)-phenylboronic acid (0.38 g, 1.5 mmol), 2M Na<sub>2</sub>CO<sub>3</sub> (1.3 mL), toluene (10 mL) and ethanol (10 mL) was degassed with nitrogen and then tetrakis(triphenylphosphine) palladium (10mg) was added. The reaction was heated to reflux overnight then cooled, filtered and concentrated. The residue was diluted with water and then extracted with ethyl acetate and dried (MgSO<sub>4</sub>). The crude product was purified by flash chromatography on silica gel using hexanes/ ethyl acetate (2:1) as eluent to afford 0.46 g (86%) of a foam; <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.94 (m, 5H), 7.63 (m, 7H), 7.32 (d, J=7.7Hz, 1H), 6.55 (s, 1H), 4.13 (s, 1H), 2.39 (s, 3H), 0.99 (s, 9H); MS m/z 514.3 (M+H<sup>+</sup>).

35

Part D. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylamino]-pyrazole, trifluoroacetic acid.

5

The product from part D was then subjected to the standard Pinner amidine sequence to obtain the desired benzamidine after preparative HPLC (acetonitrile/water, containing 0.05% TFA) as colorless crystals (44% yield).

10 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.57 (s, 1H), 9.43 (s, 1.5H), 9.14 (s, 1.5H), 8.07 (s, 1H), 8.05 (m, 1H), 7.94 (d, J=6.96Hz, 1H), 7.89 (d, J=8.42Hz, 2H), 7.76 (m, 2H), 7.65 (m, 2H), 7.53 (d, J=8.42Hz, 2H), 7.39 (s, 2H), 7.35 (m, 1H), 2.29 (s, 3H); MS (ESI) m/z 475.2 (M+H<sup>+</sup>); Analysis calculated for C<sub>24</sub>H<sub>22</sub>N<sub>6</sub>O<sub>3</sub>S<sub>1</sub> (TFA) 1.4 (H<sub>2</sub>O) 1: C 49.36; H 3.93; N 12.89; found C 49.69; H 3.71; N 12.77.

#### Example 12

1-(3-amidinophenyl)-3-methyl-5-(2'-(5'-(CF<sub>3</sub>-tetrazolyl)-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole

20

Part A. Preparation of 2-(5'-(CF<sub>3</sub>-tetrazolyl)biphenylaniline.

To a cold (0°C) CCl<sub>4</sub> (3 mL) solution of 2'-trifluoroacetanilide-1-nitro-biphenyl (0.15 g, 0.48 mmol) was added triphenylphosphine (0.24 g, 0.97 mmol) and the reaction stirred cold for 0.15 min, allowed to warm to room temperature and then gently refluxed overnight. Evaporation of the solvent afforded a residue which was treated with hexane (20 mL) filtered and evaporated to afford crude chloroimine which was dissolved in acetonitrile (10 mL). To this solution was added sodium azide (0.038 g, 0.58 mmol) and the reaction mixture was stirred at room temperature over night. Evaporation of the solvent followed by purification via silica gel flash chromatography (hexane/ethylacetate 4:1) afforded the desired nitro-biphenyltetrazole precursor (0.12 g) as a pale yellow solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.2 (d, 2H), 7.80 (t, 1H), 7.70 (t, 1H), 7.61 (d, 1H), 7.50 (d, 1H), 7.3 (d, 2H) ppm;

30

35

Ammonia CI mass spectrum analysis m/z (rel. intensity) 353.0 (M+NH<sub>4</sub><sup>+</sup> 100).

The above nitro biphenyl compound was then hydrogenated in ethanol (20 mL) over 10% Pd/C for 6 h to afford after filtration the title compound (0.11 g). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.70 (t, 1H), 7.59 (d, 1H), 7.50 (t, 1H), 7.40 (d, 1H), 6.8 (d, 2H), 6.55 (d, 2H), 3.75 (bd, 2H) ppm; Ammonia CI mass spectrum analysis m/z (rel. intensity) 323 (M+NH<sub>4</sub><sup>+</sup> 100).

Part B. Preparation of 1-(3-cyanophenyl)-3-methyl-5-(2'-(5''-CF<sub>3</sub>-tetrazolyl)-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole.

The 2-(5'-CF<sub>3</sub>-tetrazolyl)-[1,1']-biphenylaniline was then coupled to the 1-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid (0.09 g, 0.39 mmol) via the acid chloride methodology described previously to afford the title compound (0.12 g) as a colorless solid after silica gel column chromatography (dichloromethane:methanol, 9.6:0.4); <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.82 (s, 1H), 7.70 (m, 4H), 7.61 (m, 2H), 7.45 (m, 3H), 7.05 (d, 2H), 6.65 (s, 1H), 3.50 (d, 1H), 2.40 (s, 3H) ppm; Ammonia CI mass spectrum analysis m/z (rel. intensity) 532.0 (M+NH<sub>4</sub><sup>+</sup>, 100).

Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-(2'-(5''-CF<sub>3</sub>-tetrazolyl)-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole.

The product from part B was then subjected to the Pinner amidine reaction sequence as described previously to afford the title compound as colorless crystals after prep. HPLC (acetonitrile:water containing 0.05% TFA); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.61 (s, 1H), 9.42 (s, 2H), 9.12 (s, 2H), 7.94 (s, 1H), 7.89 (d, 1H), 7.82 (t, 2H), 7.75 (m, 4H), 7.62 (d, 2H), 7.02 (s, 2H), 6.98 (s, 1H), 2.32 (s, 3H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 532.4 (M+H, 100); High resolution mass spectrum calcd. for CHNFO 532.182116, found 532.18271

## Example 13

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-chloro-3-methyl-pyrazole, trifluoroacetic acid

5

Part A. Preparation of 4-chloro-1-(3'-cyanophenyl)-3-methyl-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

10

Chlorination of methyl-1-(3'-cyanophenyl)-3-methyl-pyrazole-5-carboxylate (255 mg, 1 mmol) with NCS (139 mg, 1.05 mmol) in refluxing acetonitrile (10 mL) for 3 hours gave the desired 4-chloropyrazole carboxylate in quantitative yield.

<sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.72-7.70 (m, 2H), 7.65-7.54 (m, 2H), 4.31 (q, J = 7.0 Hz, 2H), 2.35 (s, 3H), 1.28 (t, J = 7.0, 3H); LRMS: 290

15

(M+H). The ester in dichloromethane (5 mL) was added to a pretreated dichloromethane (20 mL) solution of 2'-t-butyl-sulfonamide-biphenylaniline and trimethylaluminum (2M in hexane, 1 mL) at 0°C and the resulting mixture was warmed to

20

room temperature over 15 minutes then refluxed for 3 hours. The mixture was quenched with water, extracted with CH<sub>2</sub>Cl<sub>2</sub> (200 mL), filtered through Celite. The organic layer was separated, washed with water, and brine and dried over MgSO<sub>4</sub>. After removal of the CH<sub>2</sub>Cl<sub>2</sub>, a residue was purified by column

25

chromatography with ethylacetate and methylene chloride (1:1) to afford the title compound (330 mg, 60.3%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.38 (s, 1H), 8.17 (dd, J = 8.7, J = 1.5 Hz,

1H), 7.78 (d, J = 1.5 Hz, 1H), 7.73-7.69 (m, 2H), 7.67 (d, J =

8.4 Hz, 2H), 7.60 (d, J = 7.7 Hz, 1H), 7.55 (dd, J = 7.7 Hz, J

= 1.5 Hz, 1H), 7.52 (d, J = 8.7 Hz, 2H), 7.51-7.48 (m, 1H),

7.29 (dd, J = 7.3 Hz, J = 1.5 Hz, 1H), 3.62 (s, 1H), 2.40 (s,

3H), 1.03 (s, 9H); LRMS: 548 (M+H).

30

Part B. Preparation of 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-chloro-3-methyl-pyrazole, trifluoroacetic acid

35



The product of part A was then subjected to the standard Pinner amidine sequence to afford after preparation HPLC and purification with CH<sub>3</sub>CN-H<sub>2</sub>O-TFA the title compound (350 mg).

<sup>1</sup>HNMR(CD<sub>3</sub>OD)  $\delta$ : 8.09 (dd, J = 8.1 Hz, J = 1.4 Hz, 1H), 8.00 (t, J = 1.8 Hz, 1H), 7.81 (td, J = 7.7 Hz, J = 1.9 Hz, 2H), 7.71 (d, J = 8.1 Hz, 1H), 7.64 (d, J = 8.4 Hz, 2H), 7.59 (dd, J = 7.3 Hz, J = 1.4 Hz, 1H), 7.52 (td, J = 7.7 Hz, J = 1.4 Hz, 1H), 7.42 (d, J = 8.4 Hz, 2H), 7.32 (dd, J = 8.4 Hz, J = 1.4 Hz, 1H), 2.36 (s, 3H); ESMS: 509.1 (M+H)<sup>+</sup>.

#### Example 14

1-(3-amidinophenyl)-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole

Part A. Preparation of 1-(3-cyanophenyl)-5-methyl-3-trifluoromethylpyrazole.

1,1,1-Trifluoro-2,4-pentanedione (1.35 mL, 11.2 mmol) was combined with 3-bromophenylhydrazine hydrochloride (3 g, 13.4 mmol) in glacial acetic acid (20 mL), 2-methoxyethanol (10 mL) and heated to reflux 2h. The solvents were removed in vacuo and the residue was dissolved in ethyl acetate. The ethyl acetate solution was washed successively with dilute HCl, sat'd NaHCO<sub>3</sub>, brine, and dried (MgSO<sub>4</sub>). The crude material was purified by flash chromatography on silica gel using hexanes/ethyl acetate (8:1) as eluent. The product was an 88/12 mixture of the two isomers with the desired 5-methylpyrazole isomer pre-dominating. This mixture was combined with 1-methyl pyrrolidine (7 mL) and copper cyanide (1.3 g, 14.5 mmol) and heated to reflux overnight. The reaction was cooled, diluted with ethyl acetate and filtered. The filtrate was washed with water and brine and dried (MgSO<sub>4</sub>). Purification by flash chromatography on silica gel afforded the desired 5-methylpyrazole isomer (0.66 g, 24%); <sup>1</sup>HNMR(CDCl<sub>3</sub>)  $\delta$ : 7.81 (d, J=1.8Hz, 1H), 7.77 (m, 2H), 7.67 (t, J=8.06Hz, 1H), 6.52 (s, 1H), 2.42 (s, 3H); MS (NH<sub>3</sub>) m/z 252.1 (M+H<sup>+</sup>), 269.2 (M+NH<sub>4</sub><sup>+</sup>).

Part B. Preparation of 1-(3-cyanophenyl)-5-hydroxymethyl-3-trifluoromethylpyrazole.

To the compound obtained in part A (0.65 g, 2.59 mmol), n-bromosuccinimide (0.48 g, 2.7 mmol), and benzoyl peroxide (20 mgs) were added and the reaction mixture was heated to reflux in carbon tetrachloride (20 mL) for 6h. The reaction was cooled, filtered, and concentrated to yield the crude bromide. The bromide was combined with 1:1 dioxane/ water (20 mL) and calcium carbonate (0.46 g, 4.6 mmol) and heated on a steam bath for 6h. The reaction was cooled, filtered and the filtrate concentrated. The aqueous residue was extracted with ethyl acetate and dried (MgSO<sub>4</sub>). The crude product was purified by flash chromatography on silica gel using hexanes/ethyl acetate (1:1) as eluent to afford a yellow solid (0.31 g, 44%); <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.07 (s, 1H), 8.01 (dd, J=2.2, 8.05Hz, 1H), 7.77 (d, J=7.7Hz, 1H), 7.68 (t, J=8.05Hz, 1H), 6.76 (s, 1H), 4.72 (d, J=5.85Hz, 2H), 2.02 (t, J=5.86Hz, 1H); MS (NH<sub>3</sub>) m/z 268.1 (M+H<sup>+</sup>), 285 (M+NH<sub>4</sub><sup>+</sup>).

Part C. Preparation of 1-(3-cyanophenyl)-3-trifluoromethylpyrazole-5-carboxylic acid.

To the above alcohol (0.18 g, 0.67 mmol) was added acetonitrile (5 mL), sodium periodate (0.3 g, 1.4 mmol) in water (5 mL), and one crystal of ruthenium(III)chloride hydrate. The reaction was stirred for 18h at room temperature. The reaction was filtered and concentrated. The aqueous residue was extracted with ethyl acetate and dried (MgSO<sub>4</sub>) to give 0.17 g (89.9%) of acid. <sup>1</sup>HNMR(CDCl<sub>3</sub>+DMSO-d<sub>6</sub>) δ: 7.82 (d, J=1.47Hz), 7.78 (dd, J= 8.0, 1.47Hz, 1H), 7.63 (t, J=7.3, 8.42, 1H), 7.29 (s, 1H); MS (ESI-) m/z 280.2 (M-H).

Part D. Preparation of 1-(3-cyanophenyl)-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole.

To the acid (0.35 g, 1.2 mmol) in methylene chloride was added oxalyl chloride (0.15 mL, 1.7 mmol) and 2 drops of DMF. The reaction was stirred for 2h at room temperature then concentrated in vacuo. The acid chloride was combined with  
5 2'-t-butylsulfonamide-biphenylaniline (0.38 g, 1.25 mmol), methylene chloride (10 mL), and N,N-dimethylaminopyridine (0.38 g, 3.1 mmol). The reaction was stirred overnight at room temperature. The reaction was washed with dilute HCl, sat'd NaHCO<sub>3</sub>, brine and dried (MgSO<sub>4</sub>). The crude product  
10 was purified by flash chromatography on silica gel using hexanes/ ethyl acetate (1:1) as eluent to afford 0.41 g (58%) of a yellow foam. <sup>1</sup>HNMR(CDCl<sub>3</sub>+DMSO-d<sub>6</sub>) δ: 9.88 (s, 1H), 8.18 (dd, J=7.69, 1.47Hz, 1H), 7.87 (d, J=1.83Hz, 1H), 7.79 (m, 4H), 7.64 (m, 3H), 7.50 (m, 3H), 7.30 (d, J=7.3Hz, 1H), 3.67  
15 (s, 1H), 1.02 (s, 9H); MS (ESI) m/z 590.14 (M+Na).

Part E. Preparation of 1-(3-amidinophenyl)-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole.

20

The product from part D was then subjected to the standard Pinner amidine sequence to obtain the title compound after preparative HPLC (acetonitrile/water, containing 0.05%TFA) as colorless crystals (46% yield). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ:  
25 10.85 (s, 1H), 9.47 (s, 1.5H), 9.20 (s, 1.5H), 8.05 (s, 1H), 8.04 (dd, J=7.69, 1.84Hz, 1H), 7.96 (m, 2H), 7.82 (d, J=7.69Hz, 1H), 7.75 (s, 1H), 7.68 (d, J=8.79Hz, 2H), 7.62 (m, 2H), 7.39 (d, J=8.43Hz, 2H), 7.32 (s+m, 3H); MS (ESI) m/z 529.03 (M+H<sup>+</sup>); Analysis calculated for C<sub>24</sub>H<sub>19</sub>F<sub>3</sub>N<sub>6</sub>O<sub>3</sub>S<sub>1</sub> (TFA) 1.2 (H<sub>2</sub>O) 1: C  
30 46.40; H 3.27; N 12.30; found C 46.11; H 3.06; N 12.05.

#### Example 15

1-(3-amidinophenyl)-4-methoxy-5-((2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole

35

Part A. Preparation of 1-(3-bromophenyl)-4-methoxy-5-methyl-3-trifluoromethylpyrazole.

3-Bromophenylhydrazine (9.4 g, 50.5 mmol) and trifluoroacetaldehyde hydrate (8.7 g, 75 mmol) were heated to 100°C for 1h. The reaction was cooled, diluted with methylene chloride, washed with brine and dried (MgSO<sub>4</sub>). To the crude  
5 hydrazone was added 40% aqueous pyruvic aldehyde (22.6 g, 126 mmol), MgSO<sub>4</sub> (13 g), butyl acetate (150 mL) and several drops of acetic acid and the reaction was heated to reflux overnight. The reaction was filtered and concentrated. The residue was dissolved in 1N NaOH and extracted with diethyl  
10 ether. The aqueous layer was acidified with HCl and extracted with ethyl acetate and dried (MgSO<sub>4</sub>). A crude orange solid (11.3 g, 70%) was collected. To the solid was added acetone (50 mL), K<sub>2</sub>CO<sub>3</sub> (7.3 g, 53 mmol), and iodomethane (8.8 mL, 140 mmol) and the mixture was heated to reflux for 2h. The  
15 reaction was filtered, concentrated and the crude product was purified by flash chromatography on silica gel using hexanes/ethyl acetate (4:1) as eluent to afford 6.9 g (60%) of yellow oil. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.65 (d, J=1.83Hz, 1H), 7.58 (dd, J=2.2, 6.96Hz, 1H), 7.39 (s+m, 2H), 3.85 (s, 3H), 2.31 (s, 3H); MS (H<sub>2</sub>O/GC) m/z 335-337 (M+H<sup>+</sup>).

Part B. Preparation of 1-(3-cyanophenyl)-4-methoxy-5-methyl-3-trifluoromethyl pyrazole.

25 1-(3-Bromophenyl)-4-methoxy-5-methyl-3-trifluoromethyl pyrazole (6.9 g, 20.6 mmol) and CuCN (2.8 g, 30.9 mmol) were combined in N-methylpyrrolidinone (12 mL) and heated to reflux for 18h. The reaction was diluted with water and extracted with ethyl acetate. The organic layers were washed with  
30 water, brine and dried (MgSO<sub>4</sub>). The product was purified by flash chromatography on silica gel using hexanes/ ethyl acetate (4:1) as eluent to afford 4.2 g (72%) of yellow solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.79 (s, 1H), 7.74 (m, 2H), 7.66 (d, J=7.3Hz, 1H), 3.86 (s, 3H), 2.35 (s, 3H); MS (H<sub>2</sub>O/GC) m/z 282 (M+H<sup>+</sup>); IR (KBr)  
35 2232, 1588, 1320, 1170, 1120, 804 cm<sup>-1</sup>; Analysis calculated for C<sub>13</sub>H<sub>10</sub>F<sub>3</sub>N<sub>3</sub>O<sub>1</sub>: C 55.52; H 3.58; N 14.94; found C 55.44; H 3.76; N 14.95.

Part C. Preparation of 5-bromomethyl-1-(3-cyanophenyl)-4-methoxy-3-trifluoromethylpyrazole.

To the product of part B (2.65 g, 9.40 mmol) was added n-bromosuccinimide (1.76 g, 9.90 mmol),  $\text{CCl}_4$  (15 mL) and benzoyl peroxide (10 mg). The reaction was heated to reflux for 4h, then cooled and filtered. The crude bromide was dissolved in 1:1 dioxane/water (20 mL) and  $\text{CaCO}_3$  (1.7 g, 16.9 mmol) was added. The reaction was stirred at room temperature overnight. The product was purified by flash chromatography on silica gel using hexanes/ethyl acetate (2:1) as eluent to afford 2.2 g (79%) solid. A sample was recrystallized from methylene chloride/hexanes.  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 8.10 (m, 1H), 8.05 (dd,  $J=8, 1.46\text{Hz}$ , 1H), 7.74 (d,  $J=7.7\text{Hz}$ , 1H), 7.66 (t,  $J=7.69\text{Hz}$ , 1H), 4.67 (d,  $J=5.13\text{Hz}$ , 2H), 3.95 (s, 3H), 2.17 (t,  $J=5.13\text{Hz}$ , 1H); MS (ESI)  $m/z$  288.2 ( $\text{M}+\text{H}^+$ ); Analysis calculated for  $\text{C}_{13}\text{H}_{10}\text{F}_3\text{N}_3\text{O}_2$ : C 52.53; H 3.39; N 14.14; found C 52.35; H 3.21; N 14.13.

Part D. Preparation of 1-(3-cyanophenyl)-4-methoxy-3-trifluoromethylpyrazole-5-carboxylic acid.

To the product of part C (0.64 g, 2.2 mmol) in  $\text{CH}_3\text{CN}$  (5 mL) at  $0^\circ\text{C}$  was added sodium periodate (0.98 g, 4.5 mmol) in water (5 mL) followed by one crystal of ruthenium(III) chloride. The reaction was stirred cold for 30 minutes, then at room temperature for 30 minutes. The reaction was concentrated and partitioned between ethyl acetate and dilute NaOH. The ethyl acetate layer was dried ( $\text{MgSO}_4$ ), filtered and concentrated to afford the aldehyde (0.42 g, 66%). The basic layer was acidified, extracted with ethyl acetate and dried ( $\text{MgSO}_4$ ) to afford the carboxylic acid (0.16 g, 23%). To the aldehyde (0.42 g, 1.40 mmol) was added ethanol (50 mL), silver nitrate (0.48 g, 2.8 mmol), and 0.5N NaOH (12 mL). The reaction was stirred 3h, then filtered through celite and concentrated. The aqueous layer was extracted with ethyl acetate and dried ( $\text{MgSO}_4$ ) to yield the title compound (0.4 g,

91%).  $^1\text{H NMR}$  ( $\text{CDCl}_3 + \text{DMSO}-d_6$ )  $\delta$ : 7.80 (m, 3H), 7.61 (m, 1H), 4.01 (s, 3H).

Part E. Preparation of 1-(3-cyanophenyl)-4-methoxy-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole-5-carboxylic acid.

To the acid of part D (0.44 g, 1.4 mmol) was added methylene chloride (15 mL), oxalyl chloride (0.17 mL, 1.9 mmol) and 2 drops of DMF. The reaction was stirred for 3h then, concentrated. To the crude acid chloride was added 2'-t-butylsulfonamide-biphenylaniline (0.43 g, 1.4 mmol), methylene chloride (15 mL), and triethylamine (0.8 mL, 5.6 mmol). The reaction was stirred 18h then, diluted with methylene chloride and washed with dilute HCl, sat'd  $\text{NaHCO}_3$ , brine and dried ( $\text{MgSO}_4$ ) to yield 0.6 g (52%) foam.

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 9.03 (s, 1H), 8.18 (m, 1H), 7.80 (s, 1H), 7.78 (m, 2H), 7.66 (d,  $J=8.79\text{Hz}$ , 2H), 7.65 (m, 1H), 7.56 (m, 2H), 7.52 (d,  $J=8.79\text{Hz}$ , 2H), 7.27 (m, 1H), 4.19 (s, 3H), 1.03 (s, 9H); MS (ESI)  $m/z$  598.4 ( $\text{M}+\text{H}^+$ ).

Part F. Preparation of 1-(3-amidinophenyl)-4-methoxy-5-((2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethylpyrazole.

The product from part D was subjected to the standard Pinner amidine sequence to obtain the desired benzamidine after preparative HPLC (acetonitrile/water, containing 0.05% TFA) as colorless crystals (46% yield).  $^1\text{H NMR}$  ( $\text{DMSO}-d_6$ )  $\delta$ : 11.05 (s, 1H), 9.49 (s, 1.5H), 9.22 (s, 1.5H), 8.03 (m, 2H), 7.89 (m, 3H), 7.65 (m+d,  $J=8.05\text{Hz}$ , 4H), 7.39 (m+d,  $J=8.40\text{Hz}$ , 5H), 3.96 (s, 3H); MS (ESI)  $m/z$  559.4 ( $\text{M}+\text{H}^+$ ); Analysis calculated for  $\text{C}_{25}\text{H}_{21}\text{F}_3\text{N}_6\text{O}_4\text{S}$  (TFA): C 48.22; H 3.31; N 12.50; found C 47.86; H 3.34; N 12.24.

## Example 16

**1-(3-amidinophenyl)-3-methyl-5-(4'-(imidazol-1-yl)-phenyl)aminocarbonylpyrazole**

## 5    Part A. Preparation of 1-(4-aminophenyl)imidazole.

1-(4-Nitrophenyl)imidazole (5.0 g) and 200 mL of methanol were combined to form a solution at ambient temperature. The addition of a catalytic amount of 10% palladium on carbon  
10    turned the solution into a suspension. Placement of the reaction mixture under a hydrogen atmosphere initiated the reduction. The reaction proceeded overnight (15h) at ambient temperature. Filtration through a celite pad separated out the catalyst. Concentration of the filtrate under reduced  
15    pressure gave the title product as a pale yellow solid (3.99 g). <sup>1</sup>HNMR(DMSO d<sub>6</sub>) δ: 7.95 (s, 1H), 7.45 (s, 1H), 7.18 (d, 2H), 6.99 (s, 1H), 6.60 (d, 2H), 5.25 (s, 2H) ppm. LRMS(GC/MS) m/z 160 (M+H, 100).

## 20    Part B. Preparation of N-(3-cyanophenyl)-3-methyl-5-[(4'-imidazol-1-yl)-phenyl]aminocarbonylpyrazole.

To 0.203 g of N-(3-cyanophenyl)-3-methyl-pyrazole 5-carboxylic acid and 10 mL dichloromethane was added oxalyl  
25    chloride and 2 drops of DMF. The reaction proceeded overnight. Concentration of the reaction mixture and placement under high vacuum gave the crude acid chloride which was then coupled to the product of part A under standard conditions to afford after standard purification techniques  
30    the title compound (0.118 g). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.73 (s, 1H) 9.35 (s, 1H) 8.13 (s, 1H) 7.95 (s, 1H) 7.90-7.60 (complex, 8H) 7.00 (s, 1H) 2.30 (s, 3H) ppm. LRMS(ESI) m/z 369.2 (M+H, 100). HRMS(NH<sub>3</sub>-CI) calc.369.146384, found369.145884.

## 35    Part C. Preparation of N-(3-amidinophenyl)-3-methyl-5-[(4'-imidazol-1-yl)-phenyl]aminocarbonylpyrazole.

Standard transformation of the benzonitrile obtained in part B to the benzamidine via the ethyl imidate converted 0.113 g of benzonitrile to 0.070 g of the benzamidine bis-TFA salt after HPLC purification.  $^1\text{H NMR}(\text{DMSO}-d_6)$ : 10.65 (s, 1H) 9.40 (s, 2H) 9.00 (s, 2H) 8.19 (s, 1H) 7.90 (s, 1H) 7.80-7.55 (complex, 8H) 7.06 (s, 1H) 7.00 (s, 1H) 2.30 (s, 3H) ppm. LRMS(ESI)  $m/z$  386.1 (M+H, 2) 193.7 (100). HRMS(FAB) calc. 386.172933, found 386.173388

10

**Example 17**

**1-(3-amidinophenyl)-3-methyl-5-[(4'-(2'-sulfonylmethyl)phenoxyphenyl)aminocarbonyl]pyrazole**

Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(2'-sulfonylmethyl)phenoxyphenyl)aminocarbonyl]pyrazole.

Coupling of 4-(2'-sulfonylmethyl)phenoxy-1-aminophenyl with 1-(3-cyano)phenyl-3-methyl-5-pyrazole carboxylic acid via standard acid chloride protocols described previously afforded the title compound;  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 8.05 (d, 1H), 7.82 (s, 1H), 7.78 (d, 1H), 7.65 (d, 2H), 7.55 (m, 4H), 7.10 (d, 2H), 6.95 (d, 2H), 6.65 (s, 1H), 3.32 (s, 3H), 2.40 (s, 3H) ppm; Ammonia mass spectrum analysis  $m/z$  (rel. intensity) 490 (M+NH<sub>4</sub><sup>+</sup>, 100).

25

Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(2'-sulfonylmethyl)phenoxyphenyl)aminocarbonyl]pyrazole

Subjecting the product obtained in part A to the Pinner amidine reaction sequence afforded after preparative HPLC (acetonitrile:water containing 0.05% TFA) the title compound as colorless crystals.  $^1\text{H NMR}(\text{DMSO } d_6)$   $\delta$ : 10.64 (s, 1H), 9.43 (s, 2H), 9.08 (s, 2H), 7.95 (m, 2H), 7.83 (d, 1H), 7.75 (d, 2H), 7.67 (m, 2H), 7.34 (t, 2H), 7.17 (d, 2H), 7.03 (s, 1H), 6.98 (d, 1H), 3.35 (s, 3H), 2.34 (s, 3H) ppm; ESI mass spectrum analysis  $m/z$  (rel. intensity) 490 (M+H, 100); high resolution mass spectrum calcd for CHNSO 490.153564, found 490.153759.



## Example 18

1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methylcarbonyl]-pyrazole

5

Part A. Preparation of 1-(3-cyanophenyl)-5-[(4'-bromophenyl)methylcarbonyl]-3-methylpyrazole.

To zinc dust (0.19 g, 2.9 mmol) in THF (3 mL) was added several drops of dibromoethane and the mixture was heated to reflux for 5 minutes, then cooled to 0°C. To the activated zinc was added 4-bromobenzyl bromide (0.59 g, 2.3 mmol) in THF (6 mL) dropwise over 5 minutes. The reaction was stirred at 0°C for 2h and then it was cannulated into a THF (5 mL) solution of LiCl (0.2 g, 4.7 mmol) and CuCN (0.21 g, 2.3 mmol) at -78°C. The mixture was warmed to -10°C for 5 minutes, then cooled to -78°C and the acid chloride of 1-(3-cyanophenyl)-5-carboxy-3-methylpyrazole (0.45 g, 1.98 mmol) in THF (5 mL) was added. The reaction was allowed to warm to room temperature overnight. The reaction was diluted with ethyl acetate and washed with sat'd NaHCO<sub>3</sub>, brine and dried (Na<sub>2</sub>SO<sub>4</sub>). The product was purified by flash chromatography on silica gel using hexanes/ethyl acetate (2:1) as eluent to afford 0.15 g (17%) solid: <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.67 (dd, J=1.83, 6.96Hz, 1H), 7.62 (s, 1H), 7.54 (m, 2H), 7.49 (d, J=8.42Hz, 2H), 7.13 (d, J=8.42Hz, 2H), 6.90 (s, 1H), 4.10 (s, 2H), 2.39 (s, 3H); MS (NH<sub>3</sub>) m/z 380-382 (M+H)<sup>+</sup>, 397-399 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B. Preparation of 1-(3-cyanophenyl)-5-[2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)methylcarbonyl]-3-methylpyrazole.

A mixture of the bromide above (0.14 g, 0.37 mmol), 2M Na<sub>2</sub>CO<sub>3</sub> (1 mL), 2-t-butylsulfonimide boronic acid (0.13 g, 0.50 mmol) and 1:1 ethanol/toluene (15 mL) was degassed with nitrogen for 15 minutes. Tetrakis(triphenylphosphine) palladium (2 mg) was added and the reaction was heated to reflux for 18h. The reaction was concentrated and the residue

was taken up in ethyl acetate, washed with water and dried (MgSO<sub>4</sub>). The product was purified by flash chromatography on silica gel using hexanes/ethyl acetate (2:1) as eluent to afford 0.19 g (100%) of a clear viscous oil: <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ:

5 8.18 (dd, J=1.46,7.69Hz, 1H), 7.68 (m, 2H), 7.58 (m, 2H), 7.52 (d, J=8.40Hz, 2H), 7.51 (m, 2H), 7.34 (d, J=8.05Hz, 2H), 7.33 (m, 1H), 6.95 (s, 1H), 4.21 (s, 2H), 3.48 (s, 1H), 2.40 (s, 3H), 0.97 (s, 9H); MS (ESI) m/z 535.19 (M+Na)<sup>+</sup>.

10 Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methylcarbonyl]-pyrazole.

The title compound was obtained in 37% yield following the standard Pinner-amidine sequence outlined previously.

15 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 9.39 (s, 1.5H), 9.03 (s, 1.5H), 8.03 (dd, J=7.32,1.83Hz, 1H), 7.85 (m, 2H), 7.68 (m, 2H), 7.59 (m, 2H), 7.44 (s, 1H), 7.36 (m, 7H), 4.34 (s, 2H), 2.34 (s, 3H); MS (ESI) m/z 474.18 (M+H)<sup>+</sup>.

20

#### Example 19

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole

The title compound was obtained as colorless crystals from N-1(meta-cyanophenyl)-1,2,3-triazole-2-carboxylic acid (Sheehan et. al. *J. Amer. Chem. Soc.* 1951, 73, 1207) following the general method described previously. <sup>1</sup>HNMR(DMSO d<sub>6</sub>)δ: 10.9 (s,1H), 9.49 (bs,1.5H), 9.20 (bs, 1.5H), 9.60 (s,1H), 8.11 (s,1H), 8.06-7.95 (m,3H), 7.88-7.80 (t, 1H), 7.69-7.56 (m,3H), 30 7.38 (d, 2H), 7.29 (bs, 3H) ppm; ESI mass spectral analysis m/z rel. intensity) 463 (M+H, 100); High resolution mass spectrum analysis calcd. for C<sub>21</sub>H<sub>19</sub>N<sub>8</sub>SO<sub>3</sub> 463.130084, found 463.129575.

## Example 20

1-(3-amidinophenyl)-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)tetrazole, trifluoroacetic acid salt

- 5 Part A. Preparation of 1-(3-cyanophenyl)-5-[(4'-bromophenyl)aminocarbonyl]tetrazole.

4-Bromoaniline was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (25 mL).

- 10 Trimethylaluminum (2 M in heptane 7.0 mL, 14 mmol) was added slowly. The mixture was stirred at room temperature under N<sub>2</sub> for 15 min. Then, a solution of 1-(3-cyanophenyl)-5-carboethoxy-tetrazole (0.77 g, 3.16 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25 mL) was added (prepared in part A of Example 24). The mixture was stirred at room temperature over the weekend. The reaction  
15 mixture was quenched carefully with 1N HCl. It was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water and brine, it was dried over MgSO<sub>4</sub>, concentrated, and chromatographed on silica gel (eluted with CH<sub>2</sub>Cl<sub>2</sub>) to give 0.30 g of the desired product.  
20 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 6.05 (q, 4H); 7.85 (t, 1H); 8.10 (t, 2H); 8.35 (s, 1H); 11.5 (s, 1H). MS (NH<sub>3</sub>-CI) 386 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B. Preparation of 1-(3-cyanophenyl)-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)tetrazole

- 25 The material from Part A (0.30 g, 0.813 mmol) and 2-trifluoromethyl phenylboronic acid (0.2 g, 1.06 mmol) were dissolved in EtOH/toluene (4.2 mL/10 mL). The mixture was stirred at room temperature and bubbled N<sub>2</sub> for 30 min. Then K<sub>2</sub>CO<sub>3</sub> (0.82 mL of 2 M, 1.63 mmol), tetrabutylammonium bromide  
30 (13 mg, 0.04 mmol) and tetrakis(triphenylphosphine)-palladium(0) (46 mg, 0.04 mmol) were added. The mixture was refluxed under N<sub>2</sub> for 4 hours. The reaction mixture was cooled and filter through celite. The solvent was removed. The residue was dissolved in EtOAc, washed with water and  
35 brine, it was dried over MgSO<sub>4</sub>, concentrated and chromatographed on silica gel (eluted with CH<sub>2</sub>Cl<sub>2</sub>) to give 0.35 g of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.15 to 7.95 (m, 12H); 9.15 (s, 1H). MS (NH<sub>3</sub>-CI) 452 (M+NH<sub>4</sub>)<sup>+</sup>.

Part C. Preparation of 1-(3-amidinophenyl)-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)tetrazole, trifluoroacetic acid salt.

5

The material from part B was dissolved in 10 mL anhydrous  $\text{CHCl}_3$  and 10 mL anhydrous  $\text{CH}_3\text{OH}$ . The mixture was cooled in an ice-bath and  $\text{HCl}$  gas was bubbled-in until the solution was saturated. The reaction mixture was sealed and kept at refrigerator for 12 h. The solvent was removed and the solid was dried under vacuum. The solid was redissolved in 20 mL of anhydrous  $\text{CH}_3\text{OH}$  and ammonium acetate (0.63 g, 10eq) was added. The mixture was sealed and stirred at room temperature for 12 h. The solvent was removed. The solid was dissolved in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}/\text{TFA}$  and purified by reversed phase HPLC to give 150.0 mg of the desired product.  $^1\text{H NMR}(\text{DMSO}-d_6)$   $\delta$ : 7.30 to 8.25 (m, 12H); 9.20 (s, 1H); 9.50 (s, 1H); 11.55 (s, 1H). MS (ESI) 452.2 (M+H) $^+$ .

20

#### Example 21

1-(3-amidinophenyl)-5-((2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole, trifluoroacetic acid salt

25

Part A. Preparation of 1-(3-cyanophenyl)-5-thio-tetrazole

m-Cyanophenylthioisocyanate (3.20 g, 20 mmol) was dissolved in 40 mL of  $\text{CHCl}_3$ . The mixture was heated to dissolve the starting material and a solution of  $\text{NaN}_3$  (2.64 g, 80 mmol) in 30 mL of  $\text{H}_2\text{O}$  was added. The mixture was refluxed under  $\text{N}_2$  for 1.5h. The mixture was cooled and the two layers were separated. The aqueous layer was acidified with conc.  $\text{HCl}$ . The white precipitate was filtered and dried to give 3.33 g of the desired product.  $^1\text{H NMR}(\text{acetone}-d_6)$   $\delta$ : 7.86 (t, 1H); 7.97 (d, 1H); 8.38 (d, 1H), 8.53 (s, 1H).

35

Part B. Preparation of 2'-t-butylaminosulfonyl-4-bromomethyl-3-chloro-[1,1']-biphenyl.

2'-t-Butylaminosulfonyl-3-chloro-4-methyl-[1,1']-biphenyl was converted to the bromo-compound by refluxing in NBS/CCl<sub>4</sub>.

Part C. Preparation of 1-(3-cyanophenyl)-5-((2'-t-butylaminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole.

1-(3-Cyanophenyl)-5-thio-tetrazole (0.22 g, 1.08 mmol) and 2'-t-Butylaminosulfonyl-4-bromomethyl-3-chloro-[1,1']-biphenyl (0.45 g, 1.08 mmol) were added together with 20 mL of THF. Triethylamine (0.15 mL, 1.08 mmol) was added and the mixture was refluxed under N<sub>2</sub> for 30 min. The solvent was removed, the residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub> and chromatographed on silica gel with 30% EtOAc in hexane to give 0.40 g white foam. <sup>1</sup>H NMR(CDCl<sub>3</sub>) δ: 1.03 (s, 9H); 3.58 (s, 1H); 4.82 (s, 2H); 7.26 (d, 1H); 7.37 (d, 1H); 7.53 (m, 3H); 7.75 (d, 2H); 7.82-7.92 (m, 3H), 8.16 (d, 1H). MS(ESI) 539.3 (M+H)<sup>+</sup>.

Part D. Preparation of 1-(3-amidinophenyl)-5-((2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole, trifluoroacetic acid salt

1-(3-cyanophenyl)-5-((2'-t-butylaminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole (0.24 g, 0.45 mmol) was dissolved in 20 mL of CHCl<sub>3</sub> and 2 mL of anhydrous CH<sub>3</sub>OH. The mixture was cooled in an ice-bath and HCl gas was bubbled-in until the solution was saturated. The reaction mixture was sealed and stirred at room temperature for 12 h. The solvent was removed and the solid was dried under vacuum. The solid was redissolved in 10 mL of anhydrous CH<sub>3</sub>OH and ammonium acetate (0.21 g, 6 eq) was added. The mixture was sealed and stirred at room temperature for 12 h. The solvent was removed. The solid was dissolved in CH<sub>3</sub>CN/H<sub>2</sub>O/TFA and purified by reversed phase HPLC to give 0.11 g of the title compound. <sup>1</sup>H NMR(DMSO-d<sub>6</sub>) δ: 4.79 (s, 2H); 7.30-7.69 (m, 8H); 7.90 (t, 1H); 8.02 (m, 3H); 8.11 (s, 1H); 9.20 (s, 2H); 9.48 (s, 2H). MS(ESI) 500.2 (M+H)<sup>+</sup>.

## Examples 22 and 23

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-  
biphen-4-yl)methylsulfoxide]tetrazole, trifluoroacetic acid  
5 salt (Example 22) and 1-(3-amidinophenyl)-5-[(2'-  
aminosulfonyl-3-chloro-[1,1']-biphen-4-  
yl)methylsulfonyl]tetrazole, trifluoroacetic acid salt  
(Example 23)

10 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-  
biphen-4-yl)methylthio]tetrazole, trifluoroacetic acid salt  
(80.0 mg, 0.13 mmol) was dissolved in 10 mL of methanol.  
Oxone (0.32 g, 0.52 mmol) was added. The mixture was stirred  
at room temperature under N<sub>2</sub> for 72 h. The mixture was  
15 filtered and the solid was washed with methanol. The filtrate  
was concentrated and then dissolved in CH<sub>3</sub>CN/H<sub>2</sub>O/TFA and  
purified by reversed phase HPLC to give 48 mg of the the  
sulfoxide and 23 mg of the sulfone. <sup>1</sup>HNMR(sulfoxide, CH<sub>3</sub>OH-  
d<sub>4</sub>) δ: 5.08 (q, 2H); 7.25-7.32 (m, 4H); 7.50-7.63 (m, 4H);  
20 7.85 (m, 2H); 8.00-8.10 (m, 3H). MS(ESI) 500.2 (M+H)<sup>+</sup>.  
<sup>1</sup>HNMR(sulfonyl, DMSO-d<sub>6</sub>) δ: 5.37 (s, 2H); 7.30-7.69 (m, 7H);  
7.82-8.10 (m, 5H); 8.20 (s, 1H); 9.18 (s, 2H); 9.52 (s, 2H).  
MS(ESI) 532.2 (M+H)<sup>+</sup>.

## 25 Example 24

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-  
yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

30 Part A. Preparation of 1-(3-cyanophenyl)-5-carboethoxy-  
tetrazole.

35 3-aminobenzonitrile (5.0 g, 42.3 mmol) was dissolved in  
CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Triethylamine (6.5 mL, 46.5 mmol) was added  
followed by ethyl oxalyl chloride (4.73 mL, 42.3 mmol). The  
mixture was stirred at room temperature under N<sub>2</sub> for 15 min.  
It was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water and brine.  
the CH<sub>2</sub>Cl<sub>2</sub> solution was dried over MgSO<sub>4</sub> and concentrated to a  
tan solid (6.33 g). The amide (3.00 g, 13.72 mmol) was then

refluxed 20 h with a solution of triphenylphosphine (5.4 g, 20.58 mmol) in 50 mL of  $\text{CCl}_4$ . The solution was stirred at  $0^\circ\text{C}$  for 15 min before the amide was added. The reaction mixture was cooled and hexane was added. The precipitate was filtered off. The filtrate was concentrated to a solid. It was then dissolved in 100 mL of  $\text{CH}_3\text{CN}$  and  $\text{NaN}_3$  (0.89 g, 1eq) was added. The mixture was stirred at room temperature under  $\text{N}_2$  for 12 h. The solvent was removed. The solid was dissolved in EtOAc and washed with water and brine. It was dried over  $\text{MgSO}_4$  and concentrated, and chromatographed on silica gel (eluted with  $\text{CH}_2\text{Cl}_2$ ) to give 2.50 g of the desired product.  $^1\text{H NMR}$  (acetone- $d_6$ )  $\delta$ : 1.24 (t, 3H); 4.38 (q, 2H); 7.90 (t, 1H); 8.11 (m, 2H); 8.24 (s, 1H). MS(DCI- $\text{NH}_3$ ) 261 ( $\text{M}+\text{NH}_4$ ) $^+$ .

Part B. Preparation of 1-(3-cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole.

2'-t-Butylaminosulfonyl-4-amino-[1,1']-biphenyl (0.25 g, 0.82 mmol) was dissolved in 10 mL of anhydrous  $\text{CH}_2\text{Cl}_2$ , and trimethylaluminum (1.64 mL of 2.0 M solution in heptane) was added slowly. The mixture was stirred at room temperature under  $\text{N}_2$  for 15 min, and 1-(3-cyanophenyl)-5-carboethoxy-tetrazole (0.20 g, 0.82 mmol) was added. The reaction mixture was stirred at room temperature under  $\text{N}_2$  for 18 h. The reaction was quenched carefully with 0.1N aqueous HCl. It was diluted with  $\text{CH}_2\text{Cl}_2$  and washed with water and brine. The organic solution was then dried over  $\text{MgSO}_4$ , concentrated, and chromatographed on silica gel (5% EtOAc/ $\text{CH}_2\text{Cl}_2$ ) to give 0.22 g of the desired product. MS(ESI) 502.3 ( $\text{M}+\text{H}$ ) $^+$ .

Part C. Preparation of 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

The material from Part B was dissolved in 20 mL of anhydrous  $\text{CHCl}_3$  and 5 mL of anhydrous  $\text{CH}_3\text{OH}$ . The mixture was cooled in an ice-bath and HCl gas was bubbled-in until the solution was saturated. The reaction mixture was sealed and

stirred at room temperature for 12 h. The solvent was removed and the solid was dried under vacuum. The solid was redissolved in 10 mL of anhydrous CH<sub>3</sub>OH and ammonium acetate (0.34 g, 10 eq) was added. The mixture was sealed and stirred at room temperature for 12 h. The solvent was removed. The solid was dissolved in CH<sub>3</sub>CN/H<sub>2</sub>O/TFA and purified by reversed phase HPLC to give 80.0 mg of the desired product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 7.28 (m, 3H); 7.37 (d, 2H); 7.60 (m, 2H); 7.78 (d, 2H); 7.89 (t, 1H); 8.02 (t, 2H); 8.15 (d, 1H); 8.20 (s, 1H), 9.14 (s, 2H); 9.50 (s, 2H); 11.52 (s, 1H). MS(ESI) 463.3 (M+H)<sup>+</sup>.

Examples 25-48, shown in Table 1a below, were prepared using the above described procedures.

15

**Example 49**

**3-Methyl-1-(3-amidinophenyl)-5-(4'-(4''-chlorophenyl)thiazol-2'-yl)aminocarbonylpyrazole**

Part A. Preparation of 3-methyl-1-(3-cyanophenyl)-5-(4'-(4''-chlorophenyl)thiazol-2'-ylaminocarbonyl)pyrazole.

1-(3-cyanophenyl)-3-methylpyrazole-5-carboxylic acid (70 mg, 0.31 mmol) was reacted with 2-amino-4-(4'-chlorophenyl)thiazole (168 mg, 0.8 mmol) in the presence of DMAP (191 mg, 1.5 mmol) and BOP reagent (benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate, 442 mg, 1 mmol) in DMF (5 mL) at 60°C for 16h to give the title compound (100 mg, 77%).

Part B. Preparation of 3-methyl-1-(3-amidinophenyl)-5-(4'-(4''-chlorophenyl)thiazol-2'-ylaminocarbonyl)pyrazole.

A Pinner reaction under standard procedures was used to form the title compound (39 mg, 17%): <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 7.93 (d, J = 1.8 Hz, 1H), 7.90 (d, J = 8.8 Hz, 2H), 7.86 (dd, J = 7.3 Hz, J = 1.8 Hz, 1H), 7.79-7.77 (m, 1H), 7.70 (t, J = 7.7 Hz, 1H), 7.44 (s, 1H), 7.38 (d, J = 8.4 Hz, 2H), 7.07 (s, 1H), 2.38 (s, 3H); HRMS: 437.0951 (M+H)<sup>+</sup>.



## Example 50

1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfide-  
[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

Part A. Preparation of 2'-trifluoromethylthio-1-aminobiphenyl.

Palladium catalysed Suzuki cross-coupling methodology of  
4-aminotrifluoromethylacetyl-phenylboronic acid with 2-bromo-  
1-trifluoromethylthio-benzene afforded 2'-trifluoromethylthio-  
1-aminotrifluoromethylacetyl-biphenyl in 72% yield;

$^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 8.53 (bs, 1H), 7.78 (d,  $J=8\text{Hz}$ , 1H), 7.62 (d,  $J=8\text{Hz}$ , 2H), 7.48-7.60 (m, 1H), 7.29-7.46 (m, 5H) ppm;  $^{19}\text{F NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : -42.5 (s, 3F) and -76.2 (s, 3F); Ammonia CI mass spectrum  $m/z$  (rel int.) 383 ( $\text{M}+\text{NH}_4^+$ , 100) 366 ( $\text{M}+\text{H}$ , 100).

Saponification (1N NaOH in methanol) then afforded the title compound in 80% yield;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 7.77 (d,  $J=8\text{Hz}$ , 1H), 7.30-7.55 (m, 4H), 7.09 (d,  $J=4\text{Hz}$ , 2H), 6.70 (d,  $J=8\text{Hz}$ , 2H), 3.69-3.80 (bs, 2H) ppm; Ammonia CI mass spectrum  $m/z$  (rel. int.) 256 ( $\text{M}+\text{H}$ , 100);  $^{19}\text{F NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : -42.5 ppm.

Part B. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(2'-trifluoromethylsulfide-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole.

Coupling of the product obtained in part A with the pyrazole acid chloride as illustrated in Example 10 then afforded the desired coupled phenylnitrile analog in 75% yield;  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 8.13 (bs, 1H), 7.70 (dd,  $J=1.8$  &  $7.4\text{Hz}$ , 1H), 7.51 (m, 2H), 7.48 (t,  $J=7.7\text{Hz}$ , 2H), 7.38 (t,  $J=7.6\text{Hz}$ , 2H), 7.28 (m, 2H), 6.67 (s, 1H), 2.36 (s, 3H) ppm; ESI mass spectrum  $m/z$  (rel. int.) 501 ( $\text{M}+\text{Na}$ , 92), 479 ( $\text{M}+\text{H}$ , 100);  $^{19}\text{F NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : -42.4 ppm.

Part C. Following the Pinner amidation reaction protocol as illustrated for Example 10 afforded the desired benzamidine compound in 50% yield after preparative HPLC (reverse phase,

CH<sub>3</sub>CN:water) as colorless crystals; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.7 (s, 1H), 9.43 (bs, 1.5H), 9.07 (bs, 1.5H), 7.98 (s, 1H), 7.89-7.65 (m, 8H), 7.58-7.49 (m, 2H), 7.35 (d, J=8Hz, 2H), 7.04 (s, 1H), 2.37 (s, 1H) ppm; ESI mass spectrum m/z (rel. int.) 496 (M+H, 100); HRMS calcd for C<sub>25</sub>H<sub>21</sub>N<sub>5</sub>F<sub>3</sub>SO 496.141892, Found 496.142995.

#### Examples 51 and 52

1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfoxide-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (Example 51) and 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (Example 52)

The product obtained in part C of Example 50 was subjected to oxidation with OXONE® (10eq.) in methanol/water 9:1 to afford a mixture of the sulfoxide and sulfonyl products. Preparative HPLC (reverse phase, CH<sub>3</sub>CN:water) afforded pure sulfoxide in 45% yield (colorless crystals after lyophilization); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 9.40 (bs, 1.5H), 9.04 (bs, 2H), 8.08 (d, J=8Hz, 1H), 7.96 (s, 1H), 7.84-7.68 (m, 8H), 7.50 (m, 3H), 7.04 (s, 1H), 2.35 (s, 3H) ppm; ESI mass spectrum m/z 512. The sulfonyl product was also obtained as colorless crystals in 15% yield (colorless crystals after lyophilization); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.43 (bs, 1.5H), 9.07 (bs, 2H), 8.23 (d, 1H), 7.99 (m, 1H), 7.98 (s, 1H), 7.89-7.69 (m, 7H), 7.55 (d, J=8Hz, 1H), 7.26 (d, J=8Hz, 1H), 7.04 (s, 1H), 2.37 (s, 2H) ppm; ESI mass spectrum m/z 528.1.

#### Example 53

1-(3-amidino)phenyl-3-methyl-5-[4'-(carboxymethyl)phenylaminocarbonyl]pyrazole

Methyl-4-amino-benzoate was coupled to the pyrazole acid chloride via the method illustrated for Example 10 to obtain the benzonitrile coupled product in quantitative yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.01 (d, J=8Hz, 2H), 7.97 (s, 1H), 7.80 (s, 1H), 7.78-7.53 (m, 4H), 6.70 (s, 1H), 3.90 (s, 2H), 2.39 (s, 3H) ppm; ESI mass spectrum m/z (rel. int.) 361 (M+H, 100); The nitrile was then subjected to the Pinner amidine reaction

sequence as illustrated for Example 10 to obtain after preparative HPLC separation the desired product in 50% yield (colorless crystals);  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 9.40 (bs, 1.5H), 9.18 (bs, 1.5H), 7.91 (m, 3H), 7.86-7.64 (m, 6H), 7.08 (s, 1H), 3.81 (s, 3H), 2.37 (s, 2H) ppm; ESI mass spectrum  $m/z$  (rel. int) 378 (M+H, 100); HRMS calcd for  $\text{C}_{20}\text{H}_{20}\text{N}_5\text{O}_3$  378.156615, Found 378.158283.

#### Example 54

1- (3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminocarbonyl)phenylaminocarbonyl]pyrazole

The coupled benzonitrile pyrazole methyl ester obtained above was subjected to saponification (LiOH, THF/water) followed by acidification (1N HCl) to obtain the corresponding carboxylic acid product which was then converted to the dimethyl amide derivative via its acid chloride. Following the Pinner amidine reaction protocols adopted for Example 10 then afforded the desired product as colorless crystals in 50% yield);  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 10.7 (s, 1H), 9.40 (bs, 2H), 9.04 (bs, 2H), 7.96 (s, 1H), 7.84-7.68 (m, 6H), 7.38 (d,  $J=8.0\text{Hz}$ , 2H), 7.03 (s, 1H), 2.95 (bs, 6H), 2.36 (s, 3H) ppm; ESI mass spectrum  $m/z$  (rel. int.) 391 (M+H, 100).

#### Example 55

1- (3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminosulfonyl)phenylaminocarbonyl]pyrazole

Coupling of 4-amino-N,N-dimethylbenzene-sulfonamide with the pyrazole acid chloride obtained for Example 10 afforded the desired benzonitrile-pyrazole coupled product in 90% yield.  $^1\text{H NMR}$  ( $\text{CDCl}_3$ )  $\delta$ : 8.09 (s, 1H), 7.80-7.65 (m, 7H), 7.54 (m, 1H), 6.77 (s, 1H), 2.71 (s, 6H), 2.40 (s, 3H) ppm; Ammonia CI mass spectrum (rel. int) 410 (M+H, 100). Subjecting the nitrile obtained above to the Pinner amidine reaction protocol as illustrated for Example 10 afforded the desired product in 70% yield as colorless crystals following preparative HPLC (reverse phase, acetonitrile:water) purification.  $^1\text{H NMR}$  (DMSO-

d<sub>6</sub>) δ: 10.8 (s, 1H), 9.39 (bs, 1.5H), 9.17 (bs, 1.5H), 7.89 (m, 3H), 7.79 (m, 1H), 7.77-7.63 (m, 4H), 7.06 (s, 1H), 2.30 (s, 3H), 2.45 (s, 3H) ppm; ESI mass spectrum m/z (rel. int.) 426 (M+H, 100).

5

#### Examples 56 and 57

1-(3-amidino)phenyl-3-methyl-5-[(4'-tert-butylaminosulfonylphenyl)aminocarbonyl]pyrazole (Example 56)

and 1-(3-amidino)phenyl-3-methyl-5-[(4'-aminosulfonylphenyl)aminocarbonyl]pyrazole (Example 57)

10

Coupling of 4-amino-N-tert-butylbenzene-sulfonamide with the pyrazole acid chloride obtained for Example 10 afforded the desired coupled benzonitrile precursor in 80% yield.

15 <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.35 (bs, 1H), 7.77 (m, 4H), 7.71 (m, 1H), 7.69-7.64 (m, 3H), 7.53 (t, 1H), 6.89 (s, 1H), 2.39 (s, 3H), 1.20 (s, 9H) ppm; ESI mass spectrum m/z (rel. int.) 460 (M+Na, 100), 438 (M+H, 20). Subjecting the nitrile obtained above to the Pinner amidine reaction protocol as illustrated for

20 Example 10 afforded the desired product in 5% yield as colorless crystals following preparative HPLC (reverse phase, acetonitrile:water) purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.8 (s, 1H), 9.41 (bs, 1.5H), 9.20 (bs, 1.5H), 7.97 (s, 1H), 7.84-7.77 (m, 9H), 7.47 (s, 1H), 7.08 (s, 1H), 3.73 (s, 1H), 2.35 (s, 3H) ppm; ESI mass spectrum m/z (rel. int.) 455 (M+H, 100).

25

The de-tertbutylated sulfonamide was obtained in 30% yield (colorless crystals); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.85 (s, 1H), 9.40 (bs, 4), 7.95 (s, 1H), 7.89-7.66 (m, 7H), 7.07 (s, 1H), 2.34 (s, 3H) ppm; ESI mass spectrum 381.3.

30

#### Example 58

1-(3-amidino)phenyl-3-methyl-5-[(4'-trifluoromethylphenyl)aminocarbonyl]pyrazole

35

Coupling of 4-amino-1-trifluoromethylbenzene with the acid chloride obtained in Example 10 afforded the desired benzonitrile precursor in 80% yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.17 (s, 1H), 7.79 (s, 1H), 7.75-7.50 (m, 7H), 6.73 (s, 1H), 2.39 (s,

3H) ppm; Ammonia CI mass spectrum 388 (M+NH<sub>4</sub>, 34), 371 (M+H, 100). Subjecting the nitrile obtained above to the Pinner amidine reaction protocol as illustrated for Example 10 afforded the desired product in 60% yield as colorless crystals following preparative HPLC (reverse phase, acetonitrile:water) purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.40 (bs, 1.5H), 9.20 (bs, 1.5H), 8.09 (s, 1H), 7.90 (s, 1H), 7.83-7.75 (dd, J=7.6 & 8.4Hz), 7.68-7.53 (m, 4H), 6.97 (s, 1H), 2.29 (s, 2H) ppm; ESI mass spectrum m/z (rel. int.) 388.1 (M+H, 100); HRMS calcd for C<sub>19</sub>H<sub>17</sub>N<sub>5</sub>F<sub>3</sub>O 388.138520, Found 388.139013.

#### Example 59

1-(3-amidino)phenyl-3-methyl-5-[(4'-benzylsulfonylpiperidyl)-aminocarbonyl]pyrazole

15

Coupling of 4-amino-1-benzylsulfonylpiperidine with the acid chloride obtained in Example 10 afforded the desired coupled product which when subjected to the Pinner amidine reaction protocol as illustrated for Example 10 afforded the desired product in 15% yield as colorless crystals following preparative HPLC (reverse phase, acetonitrile:water) purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.40 (bs, 1.5H), 9.00 (bs, 1.5H), 8.59 (d, J=8Hz, 1H), 7.86 (s, 1H), 7.77 (m, 1H), 7.75 (m, 3H), 7.38 (m, 5H), 6.79 (s, 1H), 4.40 (s, 2H), 3.50 (bd, 2H), 2.73 (m, 2H), 1.74 (m, 2H), 1.50 (m, 2H), 2.28 (s, 3H) ppm; ESI mass spectrum m/z (rel. int.) 481 (M+H, 100); HRMS calcd. for C<sub>24</sub>H<sub>29</sub>N<sub>6</sub> 481.202186. Found 481.201227.

#### Example 60

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-N-methylaminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

Part A. Synthesis of 1-(3-cyanophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-N-methylaminocarbonyl]-3-methylpyrazole.

Standard coupling of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid and 2-tert-butylsulfonamide-1-biphenyl-N-

methyl aniline afforded a yellow foam (67%),  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.16 (d,  $j=7.69\text{Hz}$ , 1H), 7.63 (m, 6H), 7.33 (m, 3H), 6.83 (brd m, 2H), 6.23 (s, 1H), 3.43 (s and m, 4H), 2.27 (s, 3H), 1.02 (s, 9H); MS (ESI)  $m/z$  528.4 ( $\text{M}+\text{H}$ ) $^+$ , 550.4 ( $\text{M}+\text{Na}$ ) $^+$ .

5

Part B: The Pinner amidine reaction protocol as illustrated for Example 10 afforded the desired product.  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$ : 9.45 (s, 1.5H), 9.12 (s, 1.5H), 8.16 (d,  $j=7.69\text{ Hz}$ , 1H), 7.81 (m, 7H), 7.30 (m, 5H), 7.15 (m, 2H), 3.10 (s, 3H), 2.12 (s, 3H) ppm; HRMS 489.170886 (calcd); 489.170289 (obs.); Analysis calcd for  $\text{C}_{25}\text{H}_{24}\text{N}_6\text{O}_3\text{S}(\text{TFA})1.1 (\text{H}_2\text{O})0.3$  C:52.74, H:4.18, N:13.57; found C:52.67, H:4.28, N:13.57.

10

#### Example 61

15 1-(3-amidinophenyl)-5-[(4'-fluoro-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

Part A. Preparation of 2-tert-butylsulfonamide-4-fluoro-1-biphenyl trifluoroacetamide.

20

Standard Suzuki coupling between 1-bromo-2-tert-butylsulfonamide-4-fluorobenzene (J. Indian Chem. Soc. Vol. 38, No.2, 1961,117) and 4-trifluoroacetamide-1-phenyl boronic acid afforded a solid (57%).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.11 (dd,  $j=2.19$ , 6.59Hz, 1H), 8.03 (s, 1H), 7.76 (m, 1H), 7.70 (d,  $j=8.79\text{Hz}$ , 2H), 7.61 (d,  $j=8.79\text{Hz}$ , 2H), 7.30 (m, 1H), 4.78 (s, 1H), 1.27 (s, 9H) ppm; MS (DCI)  $m/z$  436 ( $\text{M}+\text{NH}_4$ ) $^+$ ; Analysis calcd for  $\text{C}_{18}\text{H}_{18}\text{F}_4\text{N}_2\text{O}_3\text{S}_1$  C:51.67, H:4.34, N:6.70, found C:51.66, H:4.26, N:6.65.

25

30

Part B. Preparation of 2-tert-butylsulfonamide-4-fluoro-1-biphenyl aniline.

To the compound from part A (0.93 g, 2.2 mmol) in methanol was added 0.5 M LiOH (8 mL, 4 mmol) and heated to reflux 2h. The reaction was cooled and concentrated. The aqueous residue was extracted with  $\text{CH}_2\text{Cl}_2$ . The combined organic layers were washed with water, brine and dried ( $\text{MgSO}_4$ ) to

35

afford 0.7 g (98%) solid; mp=158-160 °C, <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.07 (dd, j=2.2, 6.96Hz, 1H), 7.66 (m, 1H), 7.40 (d, j=8.43Hz, 2H), 4.75 (s, 1H), 3.80 (s, 2H), 1.25 (s, 9H) ppm, MS (DCI) m/z 340 (M+NH<sub>4</sub>)<sup>+</sup>.

5

Part C: Standard coupling of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid and 2-tert-butylsulfonamide-4-fluoro-1-biphenyl aniline afforded a 85% yield of impure nitrile that was carried on to the next step. MS (DCI) m/z 531 (M+H)<sup>+</sup>, 549 (M+NH<sub>4</sub>)<sup>+</sup>.

10

Part D: The nitrile from part C was subjected to the standard Pinner conditions to give the title amidine, <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.7 (s, 1H), 9.43 (s, 1.5H), 9.01 (s, 1.5H), 7.99 (m, 3H), 7.81 (d, j=7.69Hz, 2H), 7.81 (m, 5H), 7.68 (d, j=8.79Hz, 2H), 7.55 (t, j=8.79Hz, 1H), 7.06 (s, 1H), 2.27 (s, 3H); HRMS 493.145814 (calcd); 493.145228 (obs.).

15

#### Example 62

20 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

Part A. Synthesis of 1-(3-cyanophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methylpyrazole.

25

Standard coupling of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid and 2-t-butylsulfonamide-1-pyridyl phenyl aniline afforded the title compound (44%), <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.59 (s, 1H), 8.37 (m, 1H), 8.23 (t, j=8.42, 2H), 7.94 (m, 7H), 6.77 (s, 1H), 3.94 (s, 1H), 2.41 (s, 3H), 1.10 (s, 9H) ppm, MS (ESI) 515.4 (M+H)<sup>+</sup>.

30

Part B: The above compound was subjected to standard Pinner reaction and HPLC purification (35%) <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.21 (s, 1H), 9.44 (s, 1.5H), 9.23 (s, 1.5H), 8.37 (t, j=1.47Hz, 1H), 8.07 (dd, j=7.30, 1.47Hz, 1H), 7.99 (d, j=7.69Hz, 2H), 7.85 (m, 1H), 7.79 (dd, j=9.52, 2.20Hz, 2H), 7.73 (d, j=7.69Hz,

35

1H), 7.69 (m, 2H), 7.44 (s, 2H), 7.40 (dd,  $j=2.20$ , 7.69Hz, 1H), 7.18 (s, 1H), 2.33 (s, 3H) ppm; HRMS 476.150485 (calcd), 476.149493 (observed); Analysis calcd for  $C_{23}H_{21}N_7O_3S(TFA)1.9$  C:46.51, H:3.33, N:14.17, found C:46.60, H:3.51, N:14.17.

5

**Example 63**

1-(3-cyanophenyl)-5-[[5-(2'-aminosulfonylphenyl) pyridin-2-yl]aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

10 1-(3-cyanophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methylpyrazole (0.18 g, 0.28 mmol) was heated to reflux in trifluoroacetic acid (6 mL) for 15 minutes. The reaction was concentrated and the residue purified by HPLC to afford 69 mg  
15 (43%) of the title compound.  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 11.15 (s, 1H), 8.37 (d,  $j=2.20\text{Hz}$ , 1H), 8.07 (m, 3H), 7.89 (d,  $j=7.69\text{Hz}$ , 1H), 7.82 (m, 2H), 7.70 (d,  $j=8.05\text{Hz}$ , 1H), 7.67 (m, 2H), 7.42 (s, 1H), 7.40 (dd,  $j=1.83$ , 6.96Hz, 2H), 7.18 (s, 1H), 2.32 (s, 3H) ppm; HRMS 459.123936 (calcd), 459.122035 (obs.); Analysis  
20 calcd for  $C_{23}H_{18}N_6O_3S_1(TFA)0.6$ : C:55.16, H:3.56, N:15.95, found C:54.89, H:3.69, N:15.67.

**Example 64**

25 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

Part A: 2-Trifluoromethylbromobenzene and 4-trifluoroacetamide phenylboronic acid were combined in standard Suzuki reaction to afford a 28% yield of 2-trifluoromethyl-1-biphenyl trifluoroacetamide, after  
30 purification by flash chromatography on silica gel using hexanes/ethyl acetate (6:1) as eluent.  $^1\text{H NMR}$  (CDCl $_3$ )  $\delta$ : 7.90 (s, 1H), 7.77 (d,  $j=7.69\text{Hz}$ , 1H), 7.64 (d,  $j=8.43\text{Hz}$ , 2H), 7.58 (d,  $j=6.59\text{Hz}$ , 1H),  
35 7.51 (m, 1H), 7.39 (d,  $j=8.42\text{Hz}$ , 2H), 7.33 (m, 1H) ppm, MS (ESI)  $m/z$  334 (M+H) $^+$ . 2-trifluoromethyl-1-biphenyl trifluoroacetamide was hydrolyzed with base as described above



to give the free aniline (90%) which was used in next step without purification. MS (DCI)  $m/z$  238.1 (M+H)<sup>+</sup>, 255.1 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B. Preparation of 1-(3-cyanophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methylpyrazole.

Standard coupling of 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid and 2-trifluoromethyl-1-biphenyl aniline afforded a yellow foam (50%) which was used in the next step without purification. MS (ESI)  $m/z$  447.3 (M+H)<sup>+</sup>.

Part C: The nitrile from part B was subjected to standard Pinner conditions, purified via HPLC and freeze-dried to yield the title compound (32%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)  $\delta$ : 10.68 (s, 1H), 9.44 (s, 1.5H), 9.10 (s, 1.5H), 7.97 (s, 1H), 7.84 (d,  $j$ =7.7Hz, 2H), 7.76 (m, 5H), 7.67 (m, 1H), 7.40 (d,  $j$ =7.33Hz, 1H), 7.31 (d,  $j$ =8.40Hz, d), 7.04 (s, 1H), 2.35 (s, 3H) ppm, HRMS : 464.169820 (calcd), 464.171171 (obs.); Analysis calcd for C<sub>25</sub>H<sub>20</sub>F<sub>3</sub>N<sub>5</sub>O(TFA) C:56.16, H:3.67, N:12.13, found C:55.77, H:3.79, N:11.85.

#### Example 65

1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole

To 1-(3-cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl-aminocarbonyl)-3-methyl pyrazole (0.18 g, 0.36 mmol) was added concentrated sulfuric acid (5 mL) and reaction stirred for 48h. Ice and water were added a solid precipitated. The mixture was extracted with ethyl acetate, washed with sat'd sodium bicarbonate and dried (MgSO<sub>4</sub>). Purification by flash chromatography on silica gel using 1-10% methanol in methylene chloride as eluent afforded 88 mg (52%) of the title compound, <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)  $\delta$ : 10.63 (s, 1H), 8.12 (s, 1H), 8.04 (m, 2H), 7.90 (m, 1H), 7.69 (d,  $j$ =8.42Hz, 2H), 7.62 (m, 5H), 7.36 (d,  $j$ =8.42Hz, 2H), 7.32 (m,

1H), 7.24 (s, 2H), 6.93 (s, 1H), 2.50 (s, 3H) ppm, HRMS 476.139251 (calcd), 476.139200 (observed).

### Example 66

5      1-(3-amidinophenyl)-5-[(2'-aminosulfonyl)-3-chloro-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-methylpyrazole

- Part A: 1-(3-cyanophenyl)-3-methyl-pyrazol-5-yl carboxylic acid and 4-bromo-2-chloroaniline were coupled via  
10 standard conditions (67%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.27 (d, j=8.79Hz, 1H), 8.17 (s, 1H), 7.82 (t, j=1.80Hz, 1H), 7.75 (m, 2H) 7.59 (m, 2H), 7.42 (dd, j=8.78, 2.2Hz, 1H), 6.72 (s, 1H), 2.41 (s, 3H) ppm.
- 15 Part B: The bromo compound from part A (0.4 g, 0.96 mmol), 2-t-butylsulfonamide phenylboronic acid (0.32 g, 1.2 mmol), 2M sodium carbonate (1 mL), and 1:1 toluene/ethanol were combined and degassed with nitrogen. Tetrakis(triphenylphosphine) palladium(0) (1 mg) was added and the reaction refluxed for  
20 18h. The reaction was filtered, concentrated and extracted with ethyl acetate and dried(MgSO<sub>4</sub>). Purification by flash chromatography on silica gel using 1:1 hexanes/ethyl acetate as eluent afforded 0.43 g (81%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.45 (d, j=8.42Hz, 1H), 8.32 (s, 1H), 8.18 (dd, j=1.47, 7.69Hz, 1H),  
25 7.85 (d, j=1.83Hz, 1H), 7.79 (d, j=8.05Hz, 1H), 7.72 (d, j=7.69Hz, 1H), 7.61 (m, 4H), 7.39 (dd, j=2.20, 8.79Hz, 1H), 7.28 (m, 1H), 6.76 (s, 1H), 3.67 (s, 1H), 2.43 (s, 3H), 1.07 (s, 9H) ppm., MS (ESI) m/z 548.3 (M+H)<sup>+</sup>, 570.3 (M+Na)<sup>+</sup>.
- 30 Part C: The nitrile from part B was subjected to the standard Pinner conditions to afford the amidine (43%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.36 (s, 1H), 9.43 (s, 1.5H), 9.09 (s, 1.5H), 8.05 (dd, j=6.96, 2.20Hz, 1H), 7.96 (s, 1H), 7.82 (d, j=7.32Hz, 2H),  
35 7.71 (m, 1H), 7.65 (m, 2H), 7.57 (d, j=6.59Hz, 1H), 7.54 (s, 1H), 7.46 (s, 2H), 7.39 (m, 2H), 7.06 (s, 1H), 2.35 (s, 3H) ppm, HRMS 509.116263 (calcd), 509.117360 (observed); Analysis calcd for C<sub>24</sub>H<sub>21</sub>ClN<sub>6</sub>O<sub>3</sub>S<sub>1</sub> (TFA) (H<sub>2</sub>O) C:48.72, H:3.77, N:13.11, found C:48.56, H:3.53, N:12.75.

## Example 67

1-(3-amidinophenyl)-5-[(2'-trifluoromethyl)-3-chloro-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-methylpyrazol, trifluoroacetic  
acid salt

Part A: N-(2-chloro-4-bromophenyl)-1-(3-cyanophenyl)-3-methylpyrazole carboxamide (0.4 g, 0.96 mmol), 2-trifluoromethylphenylboronic acid (0.24 g, 1.2 mmol), 1M sodium carbonate (1 mL) in 1:1 toluene/ethanol (10 mL) were degassed with nitrogen. Tetrakis(triphenylphosphine) palladium(0) (1 mg) was added and the reaction refluxed for 18h. The reaction was filtered, concentrated and extracted with ethyl acetate and dried (MgSO<sub>4</sub>). Purification by flash chromatography on silica gel using 1:1 hexanes/ethyl acetate as eluent afforded 0.41 g (90%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.40 (d, j=8.42Hz, 1H), 8.29 (s, 1H), 7.85 (d, j=1.83Hz, 1H), 7.77 (d, j=8.05Hz, 2H), 7.71 (d, j=7.60Hz, 1H), 7.60 (t, j=8.05Hz, 2H), 7.52 (t, j=7.69Hz, 1H), 7.42 (d, j=1.84Hz, 1H), 7.29 (m, 1H), 6.75 (s, 1H), 4.11 (s, 1H), 2.42 (s, 3H) ppm, MS (ESI) m/z 481.2 (M+H)<sup>+</sup>, 503 (M+Na)<sup>+</sup>.

Part B: The nitrile from part A was subjected to the standard Pinner conditions to afford the amidine (36%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.4 (s, 1H), 9.43 (s, 1.5H), 9.13 (s, 1.5H), 7.96 (d, j=1.83, 1H), 7.87 (m, 3H), 7.76 (m, 3H), 7.62 (d, j=8.06Hz, 1H), 7.52 (d, j=1.83Hz, 1H), 7.47 (d, j=7.69Hz, 1H), 7.34 (dd, j=8.42, 1.83Hz, 1H), 7.07 (s, 1H), 2.35 (s, 3H) ppm, HRMS 498.130848 (calcd), 498.128257 (observed); Analysis for C<sub>25</sub>H<sub>19</sub>ClF<sub>3</sub>N<sub>5</sub>O(TFA) calcd C:53.00, H:3.29, N:11.44, found C:53.33, H:3.36, N:11.55.

## Example 68

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-n-butylpyrazole, trifluoroacetic acid salt

Part A. Synthesis of ethyl 1-(3-cyanophenyl)-3-n-butylpyrazol-5-yl carboxylate.

Ethyl 2-methoxyimino-4-oxooctanoate (W.T.Aston, et al, J.Het.Chem., 30 (1993)2, 307) (0.69 g, 3.0 mmol) and 3-cyanophenyl hydrazine hydrochloride (0.66 g, 3.9 mmol) were combined in acetic acid (15 mL) and heated to reflux for 18h. The reaction was concentrated and the residue was partitioned between ethyl acetate and 1N HCl. The organic layer was washed with water and dried (MgSO<sub>4</sub>). A mixture of regioisomers (ca.9:1) was obtained and separated by flash chromatography on silica gel using 4:1 hexanes/ethyl acetate as eluent affording 0.56 g (63%) of the desired isomer as a yellow oil. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.77 (d, j=1.83Hz, 1H), 7.70 (d, j=7.69, 1.83Hz, 2H), 7.58 (t, j=7.69Hz, 1H), 6.88 (s, 1H), 4.30 (q, j=6.96Hz, 2H), 2.72 (t, j=7.69Hz, 2H), 1.71 (m, 2H), 1.45 (m, 2H), 1.32 (t, j=6.96Hz, 3H), 0.98 (t, j=7.33Hz, 3H) ppm; MS (DCI) m/z 298 (M+H)<sup>+</sup>.

Part B. Preparation of 1-(3-cyanophenyl)-3-n-butyl-pyrazol-5-yl carboxylic acid.

The ester from part A. (0.96 g, 3.2 mmol) was hydrolyzed with 1N NaOH (5 mL) in THF/water (5 mL) for 18h. Acid-base workup afforded 0.8 g (92%) acid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.79 (d, j=1.83Hz, 1H), 7.75 (dd, j=1.1, 8.05Hz, 1H), 7.66 (d, j=7.69Hz, 1H), 7.56 (t, j=7.69Hz, 1H), 6.88 (s, 1H), 2.71 (t, j=7.32Hz, 2H), 1.70 (m, 2H), 1.45 (m, 2H), 0.97 (t, j=7.32Hz, 3H) ppm; MS (DCI) m/z 270 (M+H)<sup>+</sup>.

Part C: Preparation of 1-(3-cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl) aminocarbonyl]-3-N-butylpyrazole.

Standard coupling of ethyl 1-(3-cyanophenyl)-3-n-butyl-pyrazol-5-yl carboxylate 2-t-butylsulfonamide-1-biphenyl aniline afforded a yellow solid (73%), <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.17 (dd, j=1.1, 7.69Hz, 1H), 8.03 (s, 1H), 7.82 (s, 1H), 7.77 (d, j=8.06, 1H), 7.68 (s+d, j=7.69Hz, 3H), 7.55 (m, 5H), 7.31 (dd, j=1.4, 7.7Hz, 1H), 6.76 (s, 1H), 3.64 (s, 1H), 2.77 (t,

j=7.69Hz, 2H), 1.75 (m, 2H), 1.44 (m, 2H), 1.03 (s, 9H), 1.00 (t, j=7.69Hz, 3H) ppm.

Part D: The nitrile from part A. was subjected to standard Pinner conditions to afford the title amidine (57%).

<sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.65 (s, 1H), 9.44 (s, 1.5H), 9.08 (s, 1.5H), 7.83 (m, 3H), 7.70 (d, j=9.15Hz, 2H) 7.64 (m, 2H), 7.37 (d, j=8.42Hz, 2H), 7.32 (d, j=7.32Hz, 1H), 7.28 (s, 2H), 7.06 (s, 1H), 2.72 (t, j=7.69Hz, 2H), 1.71 (m, 2H), 1.43 (m, 2H), 0.97 (t, j=7.33Hz, 3H) ppm, HRMS 517.202186 (calcd), 517.201333 (obs.); Analysis calcd for C<sub>27</sub>H<sub>28</sub>N<sub>6</sub>O<sub>3</sub>S(TFA)(H<sub>2</sub>O)0.8, C:54.00, H:4.78, N:3.03; found C:54.23, H:4.46, N:12.80.

#### Example 69

15 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-N-butylpyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-(3-cyanophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-N-butyl  
20 pyrazole.

Standard coupling of ethyl 1-(3-cyanophenyl)-3-n-butylpyrazol-5-yl carboxylate and 2-trifluoromethyl-1-biphenyl aniline afforded the nitrile. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.86 (s, 1H), 7.74 (m, 3H), 7.66 (m, 2H), 7.56 (m, 4H), 7.33 (m, 3H), 6.69 (s, 1H), 2.76 (t, j=7.96Hz, 2H), 1.75 (m, 2H), 1.44 (m, 2H), 0.98 (t, j=7.32Hz, 3H) ppm; MS (ESI) m/z 489 (M+H)<sup>+</sup>.

Part B: 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl) aminocarbonyl]-3-N-butyl pyrazole was prepared from the nitrile from part A by standard Pinner conditions. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.00 (s, 1H), 9.43 (s, 1.5H), 9.02 (s, 1.5H), 7.96 (s, 1H), 7.84-7.70 (m, 7H), 7.63 (t, j=7.69Hz, 1H), 7.40 (d, j=7.33Hz, 1H), 7.31 (d, j=8.42Hz, 2H), 7.08 (s, 1H), 2.72 (t, j=7.33Hz, 2H), 1.73 (m, 2H), 1.45 (m, 2H), 0.97 (t, j=7.33Hz, 3H) ppm; HRMS 506.216771 (calcd.), 506.214378 (obs.); Analysis for C<sub>28</sub>H<sub>26</sub>F<sub>3</sub>N<sub>5</sub>O(TFA)(H<sub>2</sub>O)0.8: C:56.84, H:4.55, N:11.05, found C:56.99, H:4.41, N:10.99.

## Example 70

1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-n-butylpyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-(3-cyanophenyl)-5-[[5-(2'-tert-butylsulfonaminocarbonylphenyl)pyridin-2-yl]-aminocarbonyl]-3-n-butylpyrazole.

Standard coupling of 1-(3-cyanophenyl)-3-n-butyl-pyrazol-5-yl carboxylic acid and 5-(2'-tert-butylsulfonaminocarbonylphenyl)pyridin-2-yl amine afforded the nitrile (25%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.59 (1H, s), 8.37 (d, j=2.20Hz, 1H), 8.24 (m, 2H), 7.85 (m, 2H), 7.78 (m, 1H), 7.76 (m, 1H), 7.70 (m, 3H), 7.30 (dd, j=1.47, 9.15Hz, 1H), 6.79 (s, 1H), 3.95 (s, 1H), 2.76 (t, j=7.33Hz, 2H), 1.73 (m, 2H), 1.47 (m, 2H), 1.10 (s, 9H), 0.98 (t, j=7.33Hz, 3H) ppm; MS (ESI) m/z 557.29 (M+H)<sup>+</sup>, 579.27 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B: 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-n-butylpyrazole, trifluoroacetic acid salt was prepared (51%) from the nitrile in part A by standard Pinner conditions. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.21 (s, 1H), 9.43 (s, 1.5H), 9.04 (s, 1.5H), 8.37 (d, j=2.20Hz, 1H), 8.07 (dd, j=1.83, 7.32Hz, 1H), 8.02 (d, j=8.79Hz, 1H), 7.96 (s, 1H), 7.84 (m, 3H), 7.73 (d, j=7.69Hz, 1H), 7.86 (m, 2H), 7.44 (s, 2H), 7.40 (dd, j=1.83, 6.96Hz, 1H), 7.24 (s, 1H), 2.70 (t, j=7.32Hz, 2H), 1.69 (m, 2H), 1.45 (m, 2H), 0.97 (t, j=7.32Hz, 3H) ppm; HRMS 518.197435 (calcd), 518.195873 (obs.); Analysis calc'd for C<sub>26</sub>H<sub>27</sub>N<sub>7</sub>O<sub>3</sub>S(TFA)1.5: C:50.58, H:4.17, N:14.24, found C:50.76, H:4.12, N:14.26.

## Example 71

1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-4-methoxypyrazole, trifluoroacetic acid salt

Part A: 1-(3-cyanophenyl)-3-trifluoromethyl-4-methoxy-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.97 (s, 1H), 7.80 (t, j=1.83Hz, 1H), 7.76 (s+m, 3H), 7.61 (d+m, j=8.70Hz, 4H), 7.50 (t, j=7.32Hz, 1H), 7.34 (d+m, j=8.0Hz, 3H), 4.17 (s, 3H) ppm; MS (DCI) m/z 531.2 (M+H)<sup>+</sup>.

Part B: 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-4-methoxy-pyrazole, trifluoroacetic acid salt was prepared from the nitrile of part A by standard Pinner conditions. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.06 (s, 1H), 9.47 (s, 1.5H), 9.15 (s, 1.5H), 8.03 (s, 1H), 7.92 (m, 4H), 7.75 (m, 1H), 7.70 (m, 3H), 7.40 (d, j=7.33Hz, 1H), 7.33 (d, j=8.42Hz, 2H), 3.96 (s, 3H) ppm; HRMS 548.152120 (calcd), 548.150458 (obs.); Analysis calcd for C<sub>26</sub>H<sub>19</sub>F<sub>6</sub>N<sub>5</sub>O<sub>2</sub> (TFA)1.3 (H<sub>2</sub>O)0.5: C:48.75, H:3.05, N:9.94, found C:49.04, H:2.70, N:9.85.

#### Example 72

1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-(4-bromobenzene)aminocarbonylpyrazole.

Standard coupling of 1-(3-cyanophenyl)-3-trifluoromethylpyrazole-5-yl carboxylic acid and 4-bromoaniline afforded the title compound in 77% yield ms (DCI) m/z 452-454 (M+H)<sup>+</sup>.

Part B: Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

Standard Suzuki coupling of the bromo compound from Part A and 2-trifluoromethyl phenyl boronic acid afforded the title compound (80.7%). <sup>1</sup>HNMR(CDC<sub>3</sub>)δ: 7.88 (m, 5H), 7.65 (d,

j=8.06Hz, 1H), 7.59 (m, 4H), 7.35 (d, j=8.79Hz, 2H), 7.29 (s, 1H), 7.15 (s, 1H) ppm; MS (ESI) m/z 501.2 (M+H)<sup>+</sup>.

Part C: 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-  
5 biphen-4-yl) aminocarbonyl]-3-trifluoromethyl-pyrazole,  
trifluoroacetic acid salt was prepared from the nitrile in  
part B by standard Pinner conditions. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.86  
(s, 1H), 9.46 (s, 1.5H), 9.11 (s, 1.5H), 8.05 (s, 1H), 7.95  
(d, j=8.06Hz, 2H), 7.84 (d, j=9.16Hz, 1H), 7.78 (m, 3H), 7.73  
10 (d, j=8.43Hz, 2H), 7.63 (m, 1H), 7.40 (d, j=7.69Hz, 1H), 7.32  
(d, j=8.43Hz, 2H) ppm; HRMS 518.141555 (calcd), 518.141456  
(obs.); Analysis calcd for C<sub>25</sub>H<sub>17</sub>F<sub>6</sub>N<sub>5</sub>O(TFA)1.1: C:50.82,  
H:2.84, N:10.89, found C:50.57, H:2.96, N:10.75.

15 **Example 73**

1-(3-amidinophenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-  
yl)aminocarbonyl]-3-trifluoromethyl-pyrazole, trifluoroacetic  
acid salt

20 Part A: 1-(3-cyanophenyl)-5-[(2'-sulfonylmethyl-[1,1']-  
biphen-4-yl) aminocarbonyl]-3-trifluoromethyl-pyrazole.

Standard coupling of 1-(3-cyanophenyl)-3-  
trifluoromethylphenyl and 2-sulfonylmethyl-1-biphenyl aniline  
25 afforded the nitrile in 65% yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 9.81 (s,  
1H), 8.24 (d, j=8.06Hz, 1H), 7.86 (d, j=1.83Hz, 1H), 7.82 (m,  
4H), 7.66 (m, 3H), 7.46 (s, 1H), 7.44 (d, j=6.23Hz, 2H), 7.37  
(dd, j=7.30, 1.46Hz, 1H), 2.68 (s, 3H) ppm; MS (ESI) 533.11  
(M+Na)<sup>+</sup>.

30 Part B: The title compound was prepared from the nitrile in  
part A by standard Pinner conditions, <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.92  
(s, 1H), 9.47 (s, 1.5H), 9.27 (s, 1.5H), 8.11 (dd, j=7.69,  
1.1Hz, 1H), 8.06 (s, 1H), 7.97 (m, 2H), 7.79 (m, 6H), 7.41  
35 (s+m, 2H), 2.85 (s, 3H) ppm; HRMS 528.131721 (calcd), 528.1331  
(obs.); Analysis calcd for C<sub>25</sub>H<sub>20</sub>F<sub>3</sub>N<sub>5</sub>O<sub>3</sub>S(TFA)(H<sub>2</sub>O)0.6: C:49.71,  
H:3.43, N:10.74, found C:49.48, H:3.35, N:10.97.



## Example 74

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt

5

Part A: Synthesis of 1-(3-cyano)phenyl-3-methyl-5-[(2'-t-butylaminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole.

10

The title compound was obtained by standard acid chloride coupling, of 1-(3-cyanophenyl)-3-methyl-pyrazole acid and 4-amino-2'-t-butylaminosulfonyl-3-bromo-[1,1']-biphen-4-yl (71%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.44 (d, j=8.79Hz, 1H), 8.34 (s, 1H), 8.18 (dd, j=1.47, 7.69Hz, 1H), 7.84 (m, 1H), 7.75 (d, j=1.83Hz, 1H), 7.69 (m, 1H), 7.61 (m, 3H), 7.43 (dd, j=1.83, 8.43Hz, 1H), 7.28 (m, 1H), 6.77 (s, 1H), 3.66 (s, 1H), 2.43 (s, 3H), 1.08 (s, 9H) ppm; MS (ESI) 614-616 (M+Na)<sup>+</sup>.

15

Part B: The title compound was prepared from the nitrile in part A by standard Pinner conditions. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.35 (s, 1H), 9.43 (s, 1.5H), 9.08 (s, 1.5H), 8.05 (m, 1H), 7.97 (s, 1H), 7.81 (m, 2H), 7.74 (d, j=7.69, 1H), 7.70 (d, j=1.83Hz, 1H), 7.65 (m, 2H), 7.53 (d, j=8.05Hz, 1H), 7.46 (m, 3H), 7.37 (m, 1H), 7.05 (s, 1H), 2.35 (s, 3H); HRMS 553.065747 (calcd), 553.066135 (obs.); Analysis calcd for C<sub>24</sub>H<sub>21</sub>BrN<sub>6</sub>O<sub>3</sub>S(TFA)(H<sub>2</sub>O)0.5: C:46.16, H:3.43, N:12.42, found C:46.06, H:3.15, N:12.14.

20

25

## Example 75

1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl pyrazole, trifluoroacetic acid salt

30

To 1-(3-cyano)phenyl-3-trifluoromethyl-5-[(2'-t-butylaminosulfonyl-3-bromo-[1,1']-biphen-4-yl) aminocarbonyl]-pyrazole (Part A Example 74) (82 mg, 0.14 mmol), cooled to 0°C was added conc. sulfuric acid (5 mL). The reaction was allowed to warm to room temperature and was stirred 18h.

35

Water was added and the reaction was extracted with methylene chloride. Purification by HPLC afforded 35 mg (46%) of the title amide,  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$ : 10.27 (s, 1H), 8.11 (s, 1H), 8.05 (m, 2H), 7.90 (d,  $j=7.32\text{Hz}$ , 1H), 7.68 (d,  $j=1.84\text{Hz}$ , 1H), 7.64 (m, 3H), 7.56 (dd,  $j=8.4$ , 2.2Hz, 2H), 7.51 (s, 1H), 7.44 (m, 3H), 7.36 (m, 1H), 6.96 (s, 1H), 2.34 (s, 3H) ppm; HRMS 554.049762 (calcd), 554.051045 (obs.).

#### Example 76

1-[(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl)-[1,1']-biphen-4-yl)methylcarbonyl]pyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-[(3-cyanophenyl)-5-(4-bromophenyl)acetyl]-3-methylpyrazole.

To zinc dust (0.56 g, 8.6 mmol) in THF (10 mL) was added 5 drops of 1,2-dibromoethane. The mixture was heated to reflux for 5 minutes, then cooled to  $0^\circ\text{C}$  and 4-bromobenzylbromide (1.85 g, 7.4 mmol) in THF (15 mL) was added dropwise. The reaction was stirred at  $0^\circ\text{C}$  for 2h, then it was cannulated into a suspension of LiCl (0.6 g, 1.4 mmol), CuCN (0.62 g, 7.0 mmol) and THF (20 mL). After warming to  $-20^\circ\text{C}$  for 5 minutes, the reaction was re-cooled to  $-78^\circ\text{C}$  and freshly prepared 1-(3-cyanophenyl)-3-methylpyrazol-5-yl carboxylic acid chloride (1.4 g, 5.7 mmol) in THF (15 mL) was added. The reaction was allowed to warm to room temperature and stirred 18h. The reaction was diluted with ethyl acetate and washed with brine and dried ( $\text{Na}_2\text{SO}_4$ ). Purification by chromatography on silica gel using 2:1 hexanes/ethyl acetate as eluent afforded 0.62 g (28%) of the title compound.  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 7.67 (dd,  $j=1.83$ , 6.96Hz, 1H), 7.62 (s, 1H), 7.54 (m, 2H), 7.49 (d,  $j=8.42\text{Hz}$ , 2H), 7.13 (d,  $j=8.42\text{Hz}$ , 2H), 6.90 (s, 1H), 4.10 (s, 2H), 2.39 (s, 3H) ppm; MS ( $\text{NH}_3\text{-CI}$ ) 380-382 ( $\text{M}+\text{H}$ )+, 397-399 ( $\text{M}+\text{NH}_4$ ) $^+$ .

Part B: To the product from part A (0.6 g, 1.6 mmol) was added 2-t-butylaminosulfonyl phenylboronic acid (0.57 g, 2.2 mmol), 2M sodium carbonate (3 mL) in 1:1 ethanol/toluene. The

reaction mixture was degassed with a stream of nitrogen for 15 minutes. Tetrakis(triphenylphosphine) palladium (0.3 g) was added and the reaction was heated to reflux for 24h. The reaction was cooled, filtered and concentrated. The aqueous residue was extracted with ethyl acetate, washed with brine and dried (MgSO<sub>4</sub>). Purification by chromatography on silica gel using 3:1 hexanes/ethyl acetate as eluent afforded 0.62 g (77%) of the title compound. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.18 (dd, j=1.46, 7.69 Hz, 1H), 7.68 (m, 2H), 7.58 (m, 2H), 7.52 (d+m, j=8.40, 4H), 7.34 (d+m, j=8.05 Hz, 3H), 6.95 (s, 1H), 4.21 (s, 2H), 3.48 (s, 1H), 2.40 (s, 3H), 0.97 (s, 9H) ppm; MS (ESI) 513.2 (M+H)<sup>+</sup>, 535.2 (M+Na)<sup>+</sup>.

Part C: Standard Pinner amidine reaction sequence then afforded the title compound as colorless crystals. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 9.39 (s, 1.5H), 9.03 (s, 1.5H), 8.03 (dd, j=7.32, 1.83 Hz, 1H), 7.85 (m, 2H), 7.68 (m, 2H), 7.59 (m, 2H), 7.44 (s, 1H), 7.36 (m, 7H), 4.34 (s, 2H), 2.34 (s, 3H) ppm; HRMS 474.159987 (calcd), 474.161536 (obs.); Analysis calcd for C<sub>25</sub>H<sub>23</sub>N<sub>5</sub>O<sub>3</sub>S(TFA)(H<sub>2</sub>O)0.5: C:54.36, H:4.22, N:11.74, found C:54.39, H:3.87, N:11.65.

#### Example 77

1-(3-aminocarbonylphenyl)-5-[5-[(2'-aminosulfonylphen-1-yl)pyridin-2-yl]aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

1-(3-cyanophenyl)-5-[5-[(2'-t-butylaminosulfonylphen-1-yl)pyridin-2-yl]aminocarbonyl]-3-methylpyrazole was converted to the title amide by the procedure described previously (Example 75); <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 11.15 (s, 1H), 8.35 (d, j=2.19 Hz, 1H), 8.12 (m, 4H), 7.90 (m, 1H), 7.81 (dd, j=2.20, 8.79 Hz, 1H), 7.66 (m, 2H), 7.55 (m, 2H), 7.48 (s, 1H), 7.41 (m, 3H), 7.08 (s, 1H), 2.32 (s, 3H) ppm; HRMS 477.134500 (calcd), 477.135223 (obs.).

## Example 78

1-(3-amidinophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole, trifluoroacetic acid salt

5

Part A: 1-(3-cyanophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl pyrazole was obtained via standard coupling protocols. <sup>1</sup>HNMR (CDCl<sub>3</sub>) δ: 9.13 (s, 1H), 8.64 (s, 2H), 8.22 (dd, j=1.47, 7.69Hz, 1H), 7.89 (m, 1H), 7.85 (m, 1H), 7.75 (dd, j=1.46, 6.59Hz, 1H), 7.65 (m, 3H), 7.30 (m, 2H), 4.60 (s, 1H), 1.13 (s, 9H) ppm; MS (ESI) 570.1 (M+H)<sup>+</sup>, 592.1 (M+Na)<sup>+</sup>.

10

Part B: Standard Pinner amidine reaction sequence then afforded the title compound as colorless crystals. <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ: 11.64 (s, 1H), 9.46 (s, 1.5H), 9.11 (s, 1.5H), 8.63 (s, 2H), 8.09 (dd, j=7.69, 1.83Hz, 1H), 8.04 (s, 1H), 7.96 (m, 2H), 7.81 (m, 2H), 7.76 (m, 2H), 7.42 (dd, j=1.46, 8.79Hz, 1H), 7.32 (s, 1H), 1.03 (s, 9H) ppm; HRMS 587.180069 (calcd), 587.177999 (obs.); Analysis calcd for C<sub>26</sub>H<sub>25</sub>F<sub>3</sub>N<sub>8</sub>O<sub>3</sub>S(TFA)1.1: C:47.57, H:3.69, N:15.74, found C:47.51, H:3.54, N:15.41.

15

20

## Example 79

1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole, trifluoroacetic acid salt

25

1-(3-cyanophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]-aminocarbonyl]-3-trifluoromethyl-pyrazole, (0.275 g, 0.39 mmol) was heated to reflux in TFA for 1h. Removal of TFA and purification by HPLC afforded 0.2 g (80%) title compound. <sup>1</sup>HNMR (DMSO-d<sub>6</sub>) δ: 11.63 (s, 1H), 9.46 (s, 1.5H), 9.42 (s, 1.5H), 8.66 (s, 2H), 8.08 (m, 2H), 7.96 (m, 2H), 7.83 (s, 1H), 7.81 (m, 1H), 7.72 (m, 2H), 7.54 (s, 2H), 7.45 (m, 1H) ppm; HRMS 531.117468 (calcd), 531.117523 (obs.); Analysis calcd for C<sub>22</sub>H<sub>17</sub>F<sub>3</sub>N<sub>8</sub>O<sub>3</sub>S(TFA)1.1 (H<sub>2</sub>O)0.5: C:43.71, H:2.90, N:16.85, found C:43.99, H:2.62, N:16.54.

30

35

## Example 80

1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethylpyrazole

- 1-(3-cyanophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl pyrazole (0.5 g, 0.8 mmol) was cooled to 0°C and con. H<sub>2</sub>SO<sub>4</sub> (5 mL) was added. The reaction was kept cold 24h. Ice water was added and the precipitated solid was collected, dissolved in ethyl acetate and dried (MgSO<sub>4</sub>). Purification, first, by chromatography on silica gel using 1-10% methanol/methylene chloride as eluent, then by HPLC afforded 72 mg (14%) of the title amide.
- <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.59 (s, 1H), 8.64 (s, 2H), 8.16 (s, 1H), 8.03 (s, 3H), 7.72 (m, 4H), 7.64 (d, j=7.33Hz, 1H), 7.58 (m, 1H), 7.51 (s, 2H), 7.43 (d, j=7.33Hz, 1H) ppm; HRMS 532.096112 (calcd), 532.098037 (obs.); Analysis calcd for C<sub>22</sub>H<sub>16</sub>F<sub>3</sub>N<sub>7</sub>O<sub>4</sub>S(TFA)0.5: C:46.99, H:2.83, N:16.66, found C:46.86, H:2.74, N:16.35.

## Example 81

1-(3-cyanophenyl)-5-[[4'-(imidazol-1-yl)phenyl]aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

- Part A: 1-(3-cyanophenyl)-3-trifluoromethyl-pyrazol-5-yl carboxylic acid (0.5 g, 1.8 mmol) was coupled with 4-imidazolyl aniline (0.3 g, 1.8 mmol) by standard conditions and purified by HPLC to afford 0.67 g (71%) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.55 (s, 1H), 8.22 (d, j=5.49Hz, 2H), 8.04 (d, j=7.69Hz, 1H), 7.96 (d, j=8.06Hz, 1H), 7.89 (s+d, j=8.79Hz, 3H), 7.80 (m, 4H) ppm; HRMS 423.118119 (calcd), found 423.116015 (obs.); Analysis calcd for C<sub>21</sub>H<sub>13</sub>F<sub>3</sub>N<sub>6</sub>O(TFA): C:51.50, H:2.63, N:15.67, found C:51.52, H:2.71, N:15.49.

Part B: 1-(3-cyanophenyl)-5-((4'-(imidazol-1-yl)phenyl)aminocarbonyl)-3-trifluoromethylpyrazole was subjected to

standard Pinner and purification conditions to afford title  
amidine (79%) as colorless crystals.  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$ : 11.02  
(s, 1H), 9.46 (s, 1.5H), 9.42 (s, 1H), 9.22 (s, 1.5H), 8.17 (s,  
1H), 8.06 (s, 1H), 7.97 (t,  $j=7.69\text{Hz}$ , 2H), 7.88 (d,  $j=8.79\text{Hz}$ ,  
5 2H), 7.80 (m, 3H), 7.79 (d,  $-j=9.0\text{Hz}$ , 2H) ppm; HRMS 440.144668  
(calcd), 440.144557 (obs.).

#### Examples 82 and 83

1- (3-amidinophenyl)-5-[(4'-(morpholin-1-yl)phenyl)-  
10 aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid  
salt (Example 82) and 1-(3-aminocarbonylphenyl)-5-[(4'-  
(morpholin-1-yl)phenyl)aminocarbonyl]-3-  
trifluoromethylpyrazole, trifluoroacetic acid salt (Example  
15 83)

Part A: 1-(3-cyanophenyl)-3-trifluoromethyl-pyrazol-5-yl  
carboxylic (0.34 g, 1.2 mmol) was coupled with 4-(4-  
morpholino) aniline (0.22 g, 1.2 mmol) by standard conditions  
to afford 0.53 g (69%) product.  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 9.63 (s, 1H),  
20 7.85 (d,  $j=1.46\text{Hz}$ , 1H), 7.79 (m, 1H), 7.74 (d,  $j=7.69\text{Hz}$ , 1H),  
7.60 (t,  $j=8.06\text{Hz}$ , 1H), 7.53 (d,  $j=8.79\text{Hz}$ , 2H), 7.37 (s, 1H),  
6.89 (d,  $j=9.15\text{Hz}$ , 2H), 3.87 (m, 4H), 3.87 (m, 4H) ppm; MS  
(ESI) 442.1 (M+H)+.

25 Part B: Synthesis of 1-(3-amidinophenyl)-5-[(4'-(morpholin-1-  
yl)phenyl) aminocarbonyl]-3-trifluoro-methylpyrazole,  
trifluoroacetic acid salt.

The nitrile from part A was subjected to standard Pinner  
30 conditions to afford 65% yield of the amidine as colorless  
crystals.  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$ : 10.56 (s, 1H), 9.45 (s, 1.5H), 9.13  
(s, 1.5H), 8.02 (s, 1H), 7.94 (m, 2H), 7.79 (t,  $j=7.69\text{Hz}$ , 1H),  
7.69 (s, 1H), 7.51 (d,  $j=9.16\text{Hz}$ , 2H), 6.94 (d,  $j=8.80\text{Hz}$ , 2H),  
3.74 (m, 4H), 3.08 (m, 4H) ppm; HRMS 459.175634 (calcd),  
35 459.173592 (obs.); Analysis calcd for  $\text{C}_{22}\text{H}_{21}\text{F}_3\text{N}_6\text{O}_2$  (TFA) 2.7  
( $\text{H}_2\text{O}$ ) 0.1: C:42.85, H:3.14, N:10.94, found C:42.87, H:2.78,  
N:10.80.

Part C: The amide was also isolated from the Pinner reaction in the part B.  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 10.54 (s, 1H), 8.15 (m, 2H), 7.68 (m, 1H), 7.60 (s, 1H), 7.55 (m, 1H), 7.50 (d,  $j=8.78\text{Hz}$ , 2H), 6.94 (d,  $j=8.78\text{Hz}$ , 2H), 3.73 (m, 4H), 3.07 (m, 4H) ppm; MS (ESI) 460.1 (M+H) $^+$ , 482 (M+Na) $^+$ .

#### Examples 84 and 85

1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethyl pyrazole, trifluoroacetic acid salt (Example 84) and 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt (Example 85)

Part A: 1-(3-cyanophenyl)-5-[[5-[(2'-t-butylaminosulfonyl)-1-yl] pyridin-2-yl]-aminocarbonyl]-3-trifluoromethyl pyrazole.  $^1\text{H NMR}$  (CDCl $_3$ )  $\delta$ : 8.75 (s, 1H), 8.35 (d,  $j=1.83\text{Hz}$ , 1H), 8.21 (m, 2H), 7.87 (m, 4H), 7.66 (t,  $j=7.69\text{Hz}$ , 1H), 7.59 (m, 2H), 7.29 (m, 2H), 4.30 (s, 1H), 1.11 (s, 9H) ppm; MS (ESI) 569.1 (M+H) $^+$ , 591.1 (M+Na) $^+$ .

Part B: Standard Pinner amidine reaction sequence then afforded the title compound as colorless crystals;  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 11.46 (s, 1H), 9.47 (s, 1.5H), 9.21 (s, 1.5H), 8.39 (d,  $j=1.84\text{Hz}$ , 1H), 8.06 (m, 2H), 7.97 (m, 4H), 7.82 (m, 2H), 7.68 (m, 2H), 7.45 (s, 2H), 7.40 (dd,  $j=2.20\text{Hz}$ , 7.69Hz, 1H) ppm; MS (ESI) 530.1 (M+H) $^+$ . Analysis calcd for C $_{23}$ H $_{18}$ F $_3$ N $_7$ O $_3$ S(TFA) $_2$ : C:42.81, H:2.66, N:12.44, found C:42.99, H:2.44, N:12.77.

Part C: The amide was also isolated from the Pinner reaction in the part B;  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 11.42 (s, 1H), 8.37 (d, 1H), 8.06 (s, 1H), 8.03 (m, 4H), 7.82 (m, 2H), 7.70 (m, 4H), 7.56 (s, 1H), 7.42 (s, 2H), 7.39 (dd,  $j=7.69$ , 2.2Hz, 1H) ppm; HRMS 531.106235 (calcd), 531.108937 (obs.).

## Example 86

1-(3-amidinophenyl)-5-[(4'-(3-methyltetrazol-1-yl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole,  
trifluoroacetic acid salt

5

Part A: Synthesis of 4-tetrazoyl nitrobenzene.

4-Nitrobenzonitrile (2 g, 13.5 mmol), sodium azide (0.92 g, 14 mmol), and tributyltin chloride (3.8 mL, 14 mmol) were  
10 combined in toluene (30 mL) and heated to reflux 18h. The reaction mixture was extracted with excess 1N NaOH. The aqueous layer was cooled, acidified with con. HCl, and the precipitated solid was filtered off and dried in vacuo. The aqueous layer was extracted with ethyl acetate, combined with  
15 the solid and dried (MgSO<sub>4</sub>) to afford 1.4 g (56%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.48 (d, j=8.79Hz, 2H), 8.34 (d, j=8.79Hz, 2H) ppm; MS (ES-) 190.0 (M-H).

Part B: To 4-tetrazoyl nitrobenzene (1.16 g, 6.1 mmol) and  
20 iodomethane (0.53 mL, 8.5 mmol) in DMF (10 mL) at 0°C was added 60% sodium hydride (0.29 g, 7.3 mmol). The reaction was allowed to warm to room temperature and stirred 24h. The reaction was quenched with water and extracted with ethyl acetate and dried (MgSO<sub>4</sub>). Purification by chromatography on  
25 silica gel using 4:1 hexanes/ethyl acetate as eluent afforded 0.9 g (72%) of the major isomer, 4-(3-methyltetrazole) nitrobenzene. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.38 (d, j=9.16Hz, 2H), 8.35 (d, j=9.52Hz, 2H), 4.45 (s, 3H) ppm; MS (NH<sub>3</sub>-CI) 206 (M+H)<sup>+</sup>, 176 (M+H-NO).

30

Part C: 4-(3-methyltetrazole) nitrobenzene (0.67 g, 3.3 mmol) was placed in ethanol (15 mL) and trifluoroacetic acid (1 mL). A catalytic amount of 10% Palladium on carbon was added and the mixture was placed under a hydrogen balloon. The reaction  
35 was stirred 4h, then filtered and concentrated. The 4-(3-methyltetrazole) aniline trifluoroacetic acid salt obtained (MS 176 (M+H)<sup>+</sup>) was used directly in the next step. 4-(3-Methyltetrazole) aniline trifluoroacetic acid salt and 1-(3-



cyanophenyl)-3-trifluoromethylpyrazol-5-yl carboxylic acid were coupled by standard conditions to give 1-(3-cyanophenyl)-5-[(4'-(3-methyltetrazol-1-yl)phenyl) aminocarbonyl]-3-trifluoromethylpyrazole. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 10.45 (s, 1H), 8.11 (d, j=8.79Hz, 2H), 7.86 (s, 1H), 7.82 (d, j= 8.79Hz, 2H), 7.77 (dd, j=7.69, 1.46Hz, 2H), 7.63 (t, j=7.69Hz, 1H), 7.50 (s, 1H), 4.40 (s, 3H) ppm; MS (ESI) 439 (M+H)<sup>+</sup>, 460.9 (M+Na)<sup>+</sup>.

Part D: The nitrile from part C was subjected to the standard Pinner conditions to give 1-(3-amidinophenyl)-5-[(4'-(3-methyltetrazol-1-yl)phenyl) aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.97 (s, 1H), 9.47 (s, 1.5H), 9.24 (s, 1.5H), 8.07 (d, j= 8.79Hz, 2H), 8.06 (m, 1H), 7.97 (m, 2H), 7.86 (d, j=8.78Hz, 2H), 7.80 (m, 2H), 4.41 (s, 3H) ppm; HRMS 456.150816 (calcd), 456.152474 (obs.); Analysis calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>N<sub>9</sub>O(TFA)1.2: C:45.43, H:2.93, N:21.29, found C:45.37, H:3.18, N:21.39.

20

#### Example 87

#### 1-(3-amidinophenyl)-5-(2'-naphthylaminosulfonyl)-3-methylpyrazole, trifluoroacetic acid salt

Part A: To 5-amino-1-(3-cyanophenyl)-3-methylpyrazole (0.5 g, 2.5 mmol) in methylene chloride (15 mL) was added 2-naphthylsulfonyl chloride (0.63 g, 2.8 mmol) and triethylamine (0.46 mL, 3.3 mmol). The reaction was stirred 18h at room temperature, but did not appear complete by TLC. A few crystals of N,N-dimethylaminopyridine were added and the reaction was heated to reflux for 5h. The reaction was cooled, diluted and washed with 1N HCl, sat'd NaHCO<sub>3</sub>, brine and dried (MgSO<sub>4</sub>). Crude NMR and Mass Spectrum indicated that the major product was the bis-sulfonamide, MS (ESI) 579 (M+H)<sup>+</sup>, 601 (M+Na)<sup>+</sup>.

35

Part B: The crude bis-sulfonamide from part A was subjected to the standard Pinner conditions and purified by HPLC to afford 0.3 g (50%) of the desired mono-sulfonamide title

compound,  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 9.36 (s, 1.5H), 9.07 (s, 1.5H), 8.29 (s, 1H), 8.14 (d,  $j=7.69\text{Hz}$ , 1H), 8.09 (t,  $j=8.79\text{Hz}$ , 2H), 7.86 (s, 1H), 7.79 (m, 6H), 7.60 (d,  $j=7.69\text{Hz}$ , 1H), 5.79 (s, 1H), 2.12 (s, 3H) ppm; HRMS 406.133772 (calcd), 406.133617 (obs.).

5

**Example 88**

**1-(3-amidinophenyl)-5-[(4-bromophenyl)aminosulfonyl]-3-methylpyrazole, trifluoroacetic acid salt**

10 Part A: To 5-amino-1-(3-cyanophenyl)-3-methylpyrazole (0.5 g, 2.5 mmol) in methylene chloride (15 mL) was added 4-bromobenzenesulfonyl chloride (0.7 g, 2.8 mmol) and triethylamine (0.46 mL, 3.3 mmol). The reaction was stirred 18h at room temperature, but did not appear complete by tlc. A  
15 few crystals of N,N-dimethylaminopyridine were added and the reaction was heated to reflux for 5h. The reaction was cooled, diluted and washed with 1N HCl, sat'd  $\text{NaHCO}_3$ , brine and dried ( $\text{MgSO}_4$ ). Crude NMR and Mass Spectrum indicated that the major product was the bis-sulfonamide, MS (ESI) 634-636.6  
20 ( $\text{M}+\text{H}$ ) $^+$ , 655-657.2 ( $\text{M}+\text{Na}$ ) $^+$ .

Part B: The crude bis-sulfonamide from part A was subjected to the standard Pinner conditions and purified by HPLC to afford 0.22 g (44%) of the desired mono-sulfonamide title  
25 compound,  $^1\text{H NMR}$  (DMSO- $d_6$ )  $\delta$ : 9.40 (s, 1.5H), 9.18 (s, 1.5H), 7.88 (s, 1H), 7.79 (m, 1H), 7.74 (d,  $j=8.40\text{Hz}$ , 2H), 7.69 (m, 2H), 7.53 (d,  $j=8.42\text{Hz}$ , 2H), 5.89 (s, 1H), 2.17 (s, 3H) ppm; HRMS 434.028633 (calcd), 434.029892 (obs.).

30

**Example 89**

**1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt**

To 1-(3-cyanophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl) aminocarbonyl]-3-methylpyrazole (0.19 g, 0.41 mmol) was  
35 added ethanol (20 mL), TFA (0.5 mL), and 10% Palladium on carbon (10 mg). The mixture was stirred under  $\text{H}_2$  (1 atmos.) for 18h. The reaction was filtered, concentrated and purified

by HPLC to afford 17 mg(9%) of the title compound. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.66 (s, 1H), 8.22 (brd, 2H), 8.03 (dd, j=1.47, 6.22Hz, 1H), 7.70 (d, j=8.79Hz, 2H), 7.67 (m, 2H), 7.64 (m, 5H), 7.37 (d, j=8.43Hz, 2H), 7.32 (m, 2H), 6.93 (s, 1H), 4.13 (d, j=4.03Hz, 2H), 2.33 (s, 3H)-ppm; HRMS 462.159987 (calcd), 462.158938 (found).

#### Example 90

1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

1-(3-cyanophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl) aminocarbonyl]-3-trifluoromethylpyrazole was reduced by hydrogenation to the title compound, <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.89 (s, 1H), 8.25 (brd s, 1H), 8.04 (d, j=7.33Hz, 1H), 7.75 (s, 1H), 7.69 (d+s, j=6.96Hz, 3H), 7.60 (m, 4H), 7.39 (d, j=8.43Hz, 2H), 7.32 (s+d, j=6.94Hz, 3H), 4.17 (d, j=5.49Hz, 2H) ppm; HRMS 516.131721 (calcd), 516.130109 (obs.); Analysis calcd for C<sub>24</sub>H<sub>20</sub>F<sub>3</sub>N<sub>5</sub>O<sub>3</sub>S(TFA)1.2: C:48.61, H:3.28, N:10.74, found C:48.35, H:3.18, N:10.69.

#### Example 91

1-(3-amidinophenyl)-3-methyl-5-[(2'-trifluoromethylphenyl)pyrid-2-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously. <sup>1</sup>HNMR(DMSO)δ: 11.21 (s, 1H); 9.39 (s, 2H); 9.11 (s, 2H); 8.31 (s, 1H); 8.00 (d, 1H); 7.93 (s, 1H); 7.86-7.63 (m, 7H); 7.45 (d, 1H); 7.16 (s, 1H); 2.29 (s, 3H) ppm; LRMS (ESI) 465.3 (M+H)+ HRMS for C<sub>24</sub>H<sub>20</sub>N<sub>6</sub>F<sub>3</sub>O<sub>1</sub> 465.165069 (calcd), 465.165566 (obs).

**Example 92**

**1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-1-yl)pyrimid-5-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

5

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 11.39 (s, 1H); 9.43 (s, 2H); 9.08 (s, 2H); 8.65 (s, 2H); 8.08-10 8.05 (m, 1H); 7.96 (s, 1H); 7.83 (m, 1H); 7.78-7.68 (m, 4H); 7.54 (s, 2H); 7.46-7.43 (m, 1H); 7.09 (s, 1H); 2.33 (s, 3H), ppm; LRMS (ESI) 477.2 (M+H) $^{+}$ ; HRMS for  $\text{C}_{22}\text{H}_{21}\text{N}_8\text{O}_3\text{S}_1$  477.148419 (calcd), 477.146755 (obs).

15

**Example 93**

**1-(3-amidinophenyl)-3-methyl-5-[(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 10.68 (s, 1H); 9.43 (s, 2H); 9.13 (s, 2H); 7.96 (s, 1H); 7.83-10 7.67 (m, 6H); 7.55 (d, 2H); 7.41 (m, 1H); 7.33-7.27 (m, 2H); 7.05 (s, 1H); 2.35 (s, 3H) ppm; LRMS (ESI) 414.3 (M+H) $^{+}$ ; HRMS for  $\text{C}_{24}\text{H}_{21}\text{N}_5\text{O}_1\text{F}_1$  414.173014 (calcd); 414.172475 (obs).

25

**Example 94**

**1-(3-amidinophenyl)-3-methyl-5-[3-chloro-(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

30

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 10.43 (s, 1H); 9.43 (s, 2H); 9.10 (s, 2H); 7.95 (s, 1H); 7.8210 7.73 (m, 2H); 7.68-7.54 (m, 3H); 7.46 (m, 1H); 7.38-7.30 (m, 2H); 7.07 (s, 1H); 2.35 (s, 3H) ppm; LRMS (ESI) 448.2 (M+H) $^{+}$ ; HRMS for  $\text{C}_{24}\text{H}_{20}\text{N}_5\text{OFC}_1$  448.134041 (calcd), 448.133737 (obs).

35

## Example 95

1-(3-amidinophenyl)-3-methyl-5-[3-fluoro-(2'-fluoro-[1,1']-  
biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

5

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO)δ:

10.47 (s, 1H); 9.43 (s, 2H); 9.09 (s, 2H); 7.96 (s, 1H);  
10 7.87-7.60 (m, 6H); 7.52 (m, 1H); 7.46 (d, 1H); 7.30 (d, 1H);  
7.18 (d, 1H); 7.07 (s, 1H); 2.34 (s, 3H) ppm; LRMS (ESI) 482.2 (M+H)+; HRMS for C<sub>25</sub>H<sub>2</sub>N<sub>5</sub>O<sub>4</sub>F<sub>4</sub> 482.160398 (calcd); 482.157655 (obs).

15

## Example 96

1-(3-amidinophenyl)-3-methyl-5-[3-fluoro-(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

20

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ:

10.45 (s, 1H); 9.43 (s, 2H); 9.09 (s, 2H); 8.04 (m, 1H); 7.96 (s, 1H); 7.80 (m, 2H); 7.73 (d, 1H); 7.65 (m, 3H); 7.43 (s, 2H);  
25 7.36-7.29 (m, 2H); 7.22 (m, 1H); 7.06 (s, 1H); 2.34 (s, 3H) ppm; LRMS (ESI) 493.2 (M+H)+; HRMS for C<sub>24</sub>H<sub>22</sub>N<sub>6</sub>O<sub>3</sub>SF 493.145814 (calcd), 493.146092 (obs).

## Example 97

30 1-(3-amidinophenyl)-3-methyl-5-[5-(2'-fluorophen-1-yl)pyrid-2-yl]aminocarbonyl]pyrazole, trifluoroacetic acid salt

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ:

35 11.25 (s, 1H); 9.43 (s, 2H); 9.09 (s, 2H); 8.59 (s, 1H); 8.10-8.07 (d, j=8.79, 1H); 8.02-7.96 (m, 2H); 7.85-7.79 (m, 2H); 7.73-7.70 (d, j=8.06, 1H); 7.64-7.59 (m, 1H); 7.49-7.44 (m,

1H); 7.39-7.31 (m, 2H); 7.21 (s, 1H); 2.33 (s, 3H) ppm; LRMS (ESI) 415.2 (M+H)+; HRMS for C<sub>23</sub>H<sub>20</sub>N<sub>6</sub>F 415.168263 (calcd); 425.168444 (obs).

5

**Example 98**

1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-tertbutylaminosulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole, trifluoroacetic acid salt

10

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO)δ: 11.40 (s, 1H); 9.43 (s, 2H); 9.08 (s, 2H); 8.62 (s, 2H); 8.09-8.06 (m, 1H); 7.95 (s, 1H); 7.83-7.65 (m, 6H); 7.43-7.40 (m, 1H); 7.08 (s, 1H); 2.32 (s, 3H); 1.04 (s, 9H) ppm; LRMS (ESI) 533.3 (M+H)+; HRMS for C<sub>26</sub>H<sub>29</sub>N<sub>8</sub>O<sub>3</sub>S 533.208334 (calcd), 533.207170 (obs).

15

**Example 99**

20 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-aminosulfonylphenyl)-[1,6]-dihydropyrimid-2-yl]aminocarbonyl]pyrazole, trifluoroacetic acid salt

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 9.95 (s, 1H); 9.38 (s, 2H); 9.29 (s, 1H); 9.25 (s, 2H); 7.95-7.92 (m, 2H); 7.84 (d, j=7.81, 1H); 7.79 (d, j=8.79, 1H); 7.70-7.66 (t, j=8.06, j=7.81, 1H); 7.58-7.56 (t, j=7.57, 1H); 7.54-7.49 (t, j=7.57, 1H); 7.48 (s, 2H); 7.40 (d, j=7.57, 1H); 6.86 (s, 1H); 6.13 (s, 1H); 4.24 (s, 2H); 2.28 (s, 3H) ppm; LRMS (ESI) 579.2 (M+H)+; HRMS for C<sub>22</sub>H<sub>23</sub>N<sub>8</sub>O<sub>3</sub>S 579.161384 (calcd), 579.161293 (obs).

30

35

**Example 100**

1-(3-amidinophenyl)-3-methyl-5-[(4-(pyrid-3'-yl)phen-1-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 10.71 (s, 1H); 9.43 (s, 2H); 9.11 (s, 2H); 8.98 (s, 1H); 8.64 (m, 1H); 8.28-8.25 (d,  $J=8.43$ , 1H); 7.97 (s, 1H); 7.84-7.06 (m, 8H); 7.06s, 1H); 2.35 (s, 3H), ppm LRMS (ESI) 379.2 (M+H) $^+$ ; HRMS for  $\text{C}_{23}\text{H}_{21}\text{N}_6\text{O}$  379.177685 (calcd), 379.176514 (obs).

#### Example 101

10 1-(3-amidinophenyl)-3-methyl-5-[[2-(2'-pyridyl)ethyl]aminocarbonyl]pyrazole, trifluoroacetic acid salt

15 The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 9.40 (s, 2H); 9.16 (s, 2H); 8.81 (m, 1H); 8.68 (m, 1H); 8.09 (m, 1H); 7.85 (s, 1H); 7.80-7.77 (d,  $j=8.06$ , 1H); 7.64-7.54 (m, 4H); 6.72 (s, 1H); 3.61-3.55 (q, 2H); 3.09-3.05 (t, 2H); 20 2.26 (s, 3H), ppm; LRMS (ESI) 349.1 (M+H) $^+$ ; HRMS for  $\text{C}_{19}\text{H}_{21}\text{N}_6\text{O}$  349.177685 (calcd); 349.175374 (obs).

#### Example 102

25 1-(3-amidinophenyl)-3-methyl-5-[(3-phenylpropyl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously;  $^1\text{H NMR}$  (DMSO)  $\delta$ : 30 9.41 (s, 2H); 9.11 (s, 2H); 8.72 (m, 1H); 7.88 (s, 1H); 7.81-7.77 (m, 1H); 7.68 (m, 2H); 7.31-7.18 (m, 5H); 6.77 (s, 1H); 3.21-3.14 (q,  $j=6.60$ , 2H); 2.62-2.57 (t,  $j=7.69$ , 2H); 2.28 (s, 3H); 1.82-1.73 (qu,  $j=7.32$ , 2H) ppm; LRMS (ESI) 362.1 (M+H) $^+$ ; HRMS for  $\text{C}_{21}\text{H}_{24}\text{N}_5\text{O}$  362.198086 (calcd); 362.197157 (obs).

35

## Example 103

1-(3-amidinophenyl)-3-methyl-5-[4-(pyrid-2'-yl)phen-1-ylaminocarbonyl]pyrazole, trifluoroacetic acid salt

5       The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 10.70 (s, 1H); 9.43 (s, 2H); 9.08 (s, 2H); 8.66 (m, 1H); 8.10 (m, 2H); 7.96-7.88 (m, 3H); 7.84--7.76 (m, 4H); 7.73 (m, 1H); 10       7.38 (m, 1H); 7.06 (s, 1H); 2.35 (s, 3H) ppm; LRMS (ESI) 397.1 (M+H)<sup>+</sup>; HRMS for C<sub>23</sub>H<sub>21</sub>N<sub>6</sub>O 397.177685 (calcd); 397.179670 (obs).

## Example 104

15       1-(3-amidinophenyl)-3-methyl-5-[(4-(isopropoxy)phenyl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

20       The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 10.40 (s, 1H); 9.42 (s, 2H); 9.06 (s, 2H); 7.94 (s, 1H); 7.82 (d, j=7.32, 1H); 7.75-7.65 (m, 2H); 7.54 (d, j=9.16, 2H); 6.97 (s, 1H); 6.89 (d, j=8.79, 2H); 4.57-4.53 (m, 1H); 2.32 (s, 25       3H); 1.25 (s, 3H); 1.23 (s, 3H), ppm LRMS (ESI) 378.1 (M+H)<sup>+</sup>; HRMS for C<sub>21</sub>H<sub>24</sub>N<sub>5</sub>O<sub>2</sub> 378.193000 (calcd); 378.194610 (obs).

## Example 105

30       1-(3-amidinophenyl)-3-methyl-5-[(5-(2'-trifluoromethylphenyl)-pyrimidin-2-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

35       The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 11.45 (s, 1H); 9.43 (s, 2H); 9.09 (s, 2H); 8.69 (s, 2H); 7.96 (s, 1H); 7.93 (d, j=8.06, 1H); 7.84-7.67 (m, 5H); 7.57 (d,



j=7.69, 1H); 7.10 (s, 1H); 2.32 (s, 3H) ppm; LRMS (ESI) 466.1 (M+H)+; HRMS for C<sub>23</sub>H<sub>19</sub>N<sub>7</sub>F<sub>3</sub>O 466.163004 (calcd), 466.161823 (obs).

5

**Example 106**

**1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidiniosulfonyl)phenyl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

10

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 10.90 (s, 1H); 9.42 (s, 2H); 9.19 (s, 2H); 7.95 (m, 3H); 7.80 (m, 2H); 7.70 (m, 3H); 7.08 (s, 1H); 2.85 (m, 4H); 2.35 (s, 3H); 1.54 (m, 4H); 1.35 (brd, 2H); ppm LRMS (ESI) (M+H)+ 467.1; HRMS for C<sub>23</sub>H<sub>27</sub>N<sub>6</sub>O<sub>3</sub>S 467.186536 (calcd); 467.185030 (obs).

15

**Example 107**

20

**1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidinocarbonyl)phenyl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 10.69 (s, 1H); 9.43 (s, 2H); 9.12 (s, 2H); 7.95 (s, 1H); 7.83 (m, 1H); 7.77-7.96 (m, 4H); 7.37 (d, j=8.79, 2H); 7.04 (s, 1H); 3.31 (brd, 2H); 3.54 (brd, 2H); 2.34 (s, 3H); 1.60 (brd, 2H); 1.50 (brd, 4H) ppm; LRMS (ESI) 431.1 (M+H)+.

30

**Examples 108 and 109**

35

**1-(3-amidino-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole, trifluoroacetic acid salt (Example 108) and 1-(3-aminocarbonyl-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole (Example 109)**

Part A: Preparation of 2-fluoro-5-aminobenzonitrile.

To a solution of 2-fluoro-5-nitrobenzonitrile (2.0 g, 12 mmol) in ethyl acetate (50 mL) was added stannous chloride (27.0 g, 120 mmol). The reaction mixture was stirred at reflux 1.5 h, then cooled to room temperature. Partitioned between ethyl acetate (150 mL) and saturated sodium bicarbonate (150 mL). Organic phase was separated and washed with water (5x75 mL), brine (75 mL); dried over sodium sulfate (anhy.); filtered and concentrated to give 2-fluoro-5-aminobenzonitrile (1.4 g) as pure compound.

Part B: Preparation of 3-cyano-4-fluorophenylhydrazine tin chloride.

To a solution of 2-fluoro-5-aminobenzonitrile (1.4 g, 10.3 mmol) in HCl (conc., 15 mL) at 0 °C was added a solution of sodium nitrite (0.71 g, 10.3 mmol) in cold water (3 mL) dropwisely. After addition, the mixture was stirred at 0 °C 0.5 h, a solution of stannous chloride (6.95 g, 30.9 mmol) in cold water (5 mL) was added dropwisely. The slurry was cooled in refrigerator overnight, the solid was filtered and washed with brine (20 mL), petroleum ether:ether (2:1, 30 mL) and air dried to leave 3-cyano-4-fluorophenylhydrazine tin chloride (2.5 g).

Part C: Preparation of ethyl 1-(3-cyano-4-fluorophenyl)-3-methyl-pyrazol-5-yl carboxylate.

To a mixture of 3-cyano-4-fluorophenylhydrazine tin chloride (0.9 g, 2.65 mmol) in acetic acid (15 mL) was added oxime. The reaction mixture was brought to reflux overnight. Acetic acid was removed on rotary evaporator under reduced pressure. Residue was partitioned between ethyl acetate (30 mL) and sodium bicarbonate (25 mL). Organic phase was separated and washed with water (3x20 mL), dried over sodium sulfate; filtered and concentrated; flash chromatography

(ethylacetate:hexane, 1:5) to give ethyl 1-(3-cyano-4'-fluorophenyl)-3-methyl-pyrazol-5-yl carboxylate (0.7 g).

Part D: Preparation of 1-(3-cyano-4-fluorophenyl)-3-methyl-5-  
5 [(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole.

To a solution of biphenyl amine (167 mg, 0.55 mmol) in methylene chloride (5 mL) was added trimethyl aluminum (2.0M in  
10 hexane, 0.55 mL, 1.1 mmol) via syringe at 0°C. The mixture was slowly warmed to room temperature and stirred for 20 minutes followed by portionwise addition of a solution of ethyl 1-(3-cyano-4'-fluorophenyl)-3-methyl-pyrazol-5-yl carboxylate (100 mg, 0.37 mmol) in methylene chloride (5 mL).  
15 The reaction mixture was stirred at 45°C under nitrogen overnight. Methylene chloride was removed, the residue quenched with HCl (10%, 5 mL), and partitioned between ethylacetate (20 mL) and HCl (10%, 15 mL). The organic phase was separated and washed with HCl (10%, 3x10 mL) and  
20 water (2x10 mL); dried over sodium sulfate; filtered and concentrated to leave 1-(3-cyano-4-fluorophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole (150 mg) as a pure compound.  
<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ: 8.21 (s, 1H), 8.17-8.14 (m, 1H), 7.75 (d, 1H),  
25 7.72 (t, 1H), 7.66 (d, 2H), 7.60-7.46 (m, 5H), 7.31-7.28 (m, 2H), 6.78 (s, 1H), 3.67 (s, 1H), 2.41 (s, 3H), 1.03 (s, 9H) ppm; ESI m/z (rel. intensity) 554 (M+Na, 100).

Part E: Preparation of 1-(3-amidino-4-fluorophenyl)-3-methyl-  
30 5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole, trifluoroacetic acid salt and 1-(3-carboxamido-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole

35 1-(3-cyano-4-fluorophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole (150 mg) was dissolved in a saturated HCl solution of anhydrous methanol (10 mL). The reaction mixture was stirred

at room temperature for 24h. Then methanol was evaporated. The residue was redissolved in methanol (10 mL) and excess ammonium carbonate added. The reaction mixture was stirred at room temperature overnight. Methanol was evaporated and the residue was purified via HPLC to afford 1-(3-amidino-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole as its TFA salt (20 mg).  
<sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 8.07-8.04 (m, 3H), 7.63 (d, 2H), 7.58 (d, 2H), 7.42-7.55 (m, 2H), 7.38 (d, 2H), 7.35 (d, 2H), 6.80 (s, 1H), 2.34 (s, 3H) ppm; ESI m/z (rel. intensity) 493 (M+H, 100) and 1-(3-aminocarbonyl-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole (10 mg).  
<sup>1</sup>HNMR(DMSO d<sub>6</sub>) δ: 10.59 (s, 1H), 7.99 (dd, 1H), 7.81 (br, 1H), 7.72-7.67 (m, 2H), 7.63 (d, 2H), 7.60-7.49 (m, 4H), 7.38-7.26 (m, 4H), 7.21 (s, 2H), 6.90 (s, 1H), 2.29 (s, 3H). High resolution mass spectrum calcd. for C<sub>24</sub>H<sub>20</sub>FN<sub>5</sub>O<sub>4</sub>S (M+H): 494.129829, found: 494.131923.

#### Example 110

20 **1-Methyl-3-(3-amidino)phenyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole**

Part A: A mixture of ethyl-3-cyanobenzoylacetate (2.01 g) and N,N-dimethyldiethylacetal (50 mL) were heated to gentle reflux overnight. Evaporation of the solvent *in vacuo* afforded a thick viscous reddish oil which was redissolved in anhydrous methanol (50 mL). To this solution was then added N-methylhydrazine (0.43 g) dropwise. The reaction mixture was stirred at room temperature overnight. Then concentrated to a viscous oil containing a regioisomeric mixture of pyrazoles. Without any further purification the mixture of pyrazoles obtained above (0.45 g, 1.79 mmol) was added to a dichloromethane (50 mL) solution of 2'-tert-butylsulfonamide-1-aminobiphenyl (0.54 g, 1.79 mmol) and trimethylaluminum (5.37 mL, 10.7 mmol). The reaction mixture was stirred at room temperature overnight followed by quenching with dil. HCl (1N). The organics were extracted with ethylacetate (2x50 mL), dried (MgSO<sub>4</sub>) and evaporated to a colorless residue.

Silica gel column chromatography (dichloromethane:MeOH, 9:1) afforded regioisomeric mixtures of coupled pyrazoles.

Preparatory HPLC reverse phase (acetonitrile:water gradient flow) allowed for the separation of pure 1-methyl-3-(3-cyano)phenyl-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole as a colorless oil (0.35 g);

<sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.14 (d, 1H), 8.01 (s, 1H), 7.83-7.65 (m, 4H), 7.60-7.41 (m, 6H), 7.27 (m, 2H), 3.78 (s, 3H), 3.63 (s, 1H), 1.00 (s, 9H) ppm; ESI mass spectrum 536 (M+Na, 45), 514 (M+H, 100).

Part B: The product from part A was then subjected to the Pinner amidine reaction sequence as outlined in Example 10 to obtain after preparative HPLC separation and lyophilization colorless crystals of the title compound (0.15 g); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.90 (s, 1H), 9.37 (bs, 1.5H), 9.29 (bs, 1.5H), 8.24 (s, 1H), 8.00 (d, 1H), 7.90 (bs, 2H), 7.84 (d, 1H), 7.73 (m, 1H), 7.69-7.50 (m, 4H), 7.37-7.27 (m, 3H), 7.17 (bs, 1H), 3.98 (s, 3H) ppm; ESI mass spectrum m/z (rel. int.) 475.3 (M+H, 100).

#### Example 111

1-(3-amidinophenyl)-5-[[4-(pyrazol-4'-yl)phen-1-yl]aminocarbonyl]pyrazole, trifluoroacetic acid salt

Part A: 4-Iodopyrazole (20 mmol) was treated with Et<sub>3</sub>N (30 mmol) and (Boc)<sub>2</sub>O (22 mmol) in THF (60 mL) at r.t. for 2 hours to form N-Boc-4-iodopyrazole (5.88 g, 100%). N-Boc-4-iodopyrazole in THF (100 mL) was reacted with hexamethylditin (20 mmol) in the presence of Pd(Ph<sub>3</sub>P)<sub>4</sub> (1.1 g, 1 mmol) under nitrogen at 78°C overnight. To it was added aqueous 10% KF and the resulting mixture was stirred for 30 minutes, and then filtered through a pad of Celite. The filtrate was extracted with EtOAc. The EtOAc layer was washed with water, and dried over MgSO<sub>4</sub>. Filtration and concentration followed by purification of the mixture by column chromatography afforded the 3-trimethyltinpyrazole derivative (5 g, 75%) as a white solid.

Part B: The product from part A (10 mmol) was treated with 4-nitrobromobenzene (10 mmol) in the presence of  $\text{Pd}(\text{Ph}_3\text{P})_4$  (0.36 g, 0.3 mmol) under nitrogen at 78°C overnight, followed by workup as described above afforded the 4-pyrazolo-nitrobenzene derivative (0.95 g, 33%). Hydrogenation (0.85 g, 2.94 mmol) in MeOH (150 mL) in the presence of Pd (5% on C, 0.09 g) at r.t. for 16 hours afforded the aniline derivative (0.76 g, 100%).

Part C: Standard coupling of the product from part B with the pyrazole acid chloride under reflux for 1.5 with  $\text{Et}_3\text{N}$  (1 mL) followed by usual workup and purification afforded the coupled amide pyrazole-benzonitrile derivative (255 mg, 55%) which was subjected to the Pinner amidine sequence to afford after purification the title compound as colorless (148 mg, 70%).  $^1\text{H NMR}(\text{CD}_3\text{OD}) \delta$ : 7.93 (bs, 2H), 7.90-7.87 (m, 1H), 7.80 (td,  $J=7.4$  Hz,  $J=1.2$  Hz, 2H), 7.70 (t,  $J=7.8$  Hz, 1H), 7.57 (d,  $J=7.4$  Hz, 2H), 7.60-7.54 (m, 2H), 6.93 (d,  $J=1.9$  Hz, 1H), 2.38 (s, 3H);  $^{13}\text{C NMR}(\text{CD}_3\text{OD}) \delta$ : 167.92, 159.84, 151.36, 142.27, 139.28, 137.30, 131.43, 131.07, 130.51, 128.33, 126.99, 125.48, 122.48, 110.77, 13.29; ESMS:  $m/z$  386.3 ( $\text{M}+\text{H}$ , 100); HRMS calcd for  $\text{C}_{21}\text{H}_{20}\text{N}_7\text{O}_1$  386.1729, found 386.1738.

#### Example 112

1-(3-amidinophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrid-2-yl]aminocarbonyl)pyrazole trifluoroacetate

Part A: Preparation of 2-(tertbutoxycarbonyl)amino-5-bromopyridine and 2-[bis(tertbutoxycarbonyl)amino]-5-bromopyridine.

Sodium hydride (1.27 g, 60%, 32 mmol) was added to 2-amino-5-bromopyridine (5.0 g, 29 mmol) in THF (75 mL) at 0°C. The ice bath was removed and the reaction stirred 25 min at room temp. Di-*t*-butyl dicarbonate (6.94 g, 32 mmol) was added and the reaction was refluxed 15 h. After cooling, the

reaction was carefully quenched with water and extracted into EtOAc. The combined organics were washed with sat.  $\text{NH}_4\text{Cl}$  and sat.  $\text{NaHCO}_3$ , dried over  $\text{Na}_2\text{SO}_4$ , filtered, and evaporated. The crude mixture was chromatographed on silica gel (5-7.5% EtOAc/hexanes, followed with 100%  $\text{CHCl}_3$ ) to yield both the mono-protected (2.85 g, 36%) and bis-protected (1.87 g, 17%) products.  $^1\text{H NMR}$ (mono,  $\text{CDCl}_3$ ) $\delta$ : 8.32 (d, 1H,  $J=2.2$ ), 8.13 (bs, 1H), 7.91 (d, 1H,  $J=8.8$ ), 7.75 (dd, 1H,  $J=8.8$ ,  $J'=2.2$ ), 1.54 (s, 9H);  $^1\text{H NMR}$ (bis,  $\text{CDCl}_3$ ) $\delta$ : 8.53 (d, 1H,  $J=2.5$ ), 7.84 (dd, 1H,  $J=8.5$ ,  $J'=2.5$ ), 7.18 (d, 1H,  $J=8.4$ ), 1.45 (s, 18H).

Part B: Preparation of 2-[bis(tertbutoxycarbonyl)amino]-5-(2'-methylthiophenyl)pyridine.

2-[Bis(tertbutoxycarbonyl)amino]-5-bromopyridine (1.87 g, 5.0 mmol) was dissolved in benzene (120 mL). 2-Methylthiophenylboronic acid (1.95 g, 11.5 mmol), aq. sodium carbonate (12 mL, 2.0 M, 24 mmol), tetrabutyl ammonium bromide (80 mg, 0.25 mmol), and bis(triphenylphosphine)palladium(II)chloride (175 mg, 0.25 mmol) were added, and the resulting mixture was purged with vacuum and argon and then refluxed 16 h. The cooled mixture was diluted with EtOAc and water. The layers were separated, and the organic phase was washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, and evaporated. The crude product was chromatographed on silica gel (10-20% EtOAc/hexanes) to yield a thick oil (1.82 g, 87.1%).  $^1\text{H NMR}$ ( $\text{CDCl}_3$ ) $\delta$ : 8.51 (d, 1H,  $J=2.2$ ), 7.83 (dd, 1H,  $J=8.1$ ,  $J'=2.2$ ), 7.30 (m, 5H), 2.35 (s, 3H), 1.47 (s, 18H).

Part C: Preparation of 2-[bis(tertbutoxycarbonyl)amino]-5-(2'-methylsulfonylphenyl)pyridine

2-[Bis(tertbutoxycarbonyl)amino]-5-(2'-methylthiophenyl)pyridine (1.69 g, 4.1 mmol) was dissolved in MeOH (20 mL). In a separate beaker, a solution of Oxone (10 g) was generated by dilution to 49 mL with water. A portion of this solution (14.5 mL, 4.8 mmol) was removed and adjusted

to pH 4 with sat.  $\text{Na}_3\text{PO}_4$  solution (4.0 mL). This mixture was added to the reaction and stirred 22 h at room temp. The resulting mixture was diluted with water, extracted with  $\text{CHCl}_3$ , dried over  $\text{Na}_2\text{SO}_4$ , filtered, and evaporated. The crude product was chromatographed on silica gel (40-75% EtOAc/hexanes) to yield the sulfone (1.19 g, 65%).  
5  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.48 (d, 1H,  $J=1.8$ ), 8.26 (dd, 1H,  $J=8.1$ ,  $J'=1.5$ ), 7.95 (dd, 1H,  $J=8.1$ ,  $J'=2.2$ ), 7.71 (td, 1H,  $J=7.4$ ,  $J'=1.5$ ), 7.64 (td, 1H,  $J=7.7$ ,  $J'=1.4$ ), 7.40 (dd, 1H,  $J=7.3$ ,  $J'=1.4$ ), 7.36 (d, 1H,  $J=8.8$ ), 2.68 (s, 3H), 1.48 (s, 18H).  
10

Part D: Preparation of 2-amino-5-(2'-methylsulfonylphenyl)pyridine hydrochloride.

15 2-[Bis(tertbutoxycarbonyl)amino-5-(2'-methylsulfonylphenyl)pyridine (1.18 g, 2.6 mmol) and 2-(tertbutoxycarbonyl)amino-5-(2'-methylsulfonylphenyl)pyridine (1.62 g, 4.6 mmol) were suspended in HCl/dioxane (30 mL, 4.0 M) and stirred 23 h at room temp. The resulting mixture was  
20 diluted with  $\text{Et}_2\text{O}$  and filtered to yield a tan solid (2.27 g, 100%).  $^1\text{H NMR}(\text{DMSO})\delta$ : 8.09 (m, 3H), 7.98 (d, 1H,  $J=1.8$ ), 7.90 (dd, 1H,  $J=9.1$ ,  $J'=2.2$ ), 7.75 (m, 2H), 7.45 (dd, 1H,  $J=7.3$ ,  $J'=1.1$ ), 6.98 (d, 1H,  $J=9.1$ ), 3.04 (s, 3H).

25 Part E: Preparation of 1-(3-cyanophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrid-2-yl]aminocarbonyl)pyrazole.

Oxalyl chloride (175  $\mu\text{L}$ , 2.0 mmol) and DMF (2 drops) were added to 1-(3-cyanophenyl)-3-methylpyrazole-5-carboxylic acid  
30 (300 mg, 1.3 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and stirred under argon 160 min. The resulting solution was evaporated and redissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL). 4-Dimethylaminopyridine (484 mg, 4.0 mmol) and 2-amino-5-(2'-methylsulfonylphenyl)pyridine hydrochloride (376 mg, 1.3 mmol) were added and stirred at  
35 room temperature under argon for days. The reaction was evaporated and chromatographed on silica gel (50-100% EtOAc/hexanes, followed by 1% MeOH/EtOAc) to yield the desired product (303 mg, 50%).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.54 (s, 1H), 8.39 (d,



1H, J=2.2), 8.25 (d, 2H, J=8.4), 7.82 (m, 2H), 7.66 (m, 5H), 7.37 (dd, 1H, J=7.7, J'=1.5), 6.76 (s, 1H), 2.75 (s, 3H), 2.41 (s, 3H).

- 5 Part F: Preparation of 1-(3-amidinophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrid-2-yl]aminocarbonyl)pyrazole trifluoroacetate.

1-(3-cyanophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrid-2-yl]aminocarbonyl)pyrazole (300 mg, 0.66 mmol) was dissolved in dry CHCl<sub>3</sub> (15 mL) and dry MeOH (5 mL) and cooled to 0°C. HCl (g) was generated by the addition of H<sub>2</sub>SO<sub>4</sub> (45 mL) to NaCl (200 g) over 90 min, and bubbled into the reaction. The generator was removed, and the reaction was sealed and placed in the refrigerator (4°C) overnight. The reaction was evaporated and redissolved in dry MeOH (10 mL). Ammonium carbonate (316 mg, 3.3 mmol) was added and the reaction was stirred 20 h at room temp, and evaporated. The crude product was purified by prep HPLC on a C-18 reverse phase column (10-70% MeCN/H<sub>2</sub>O/0.05% TFA) to yield a white powder (161 mg, 42%). <sup>1</sup>HNMR(DMSO)δ: 11.21 (s, 1H), 9.38 (s, 2H), 8.96 (s, 2H), 8.36 (s, 1H), 8.07 (d, 1H, J=7.3), 7.99 (d, 1H, J=8.5), 7.92 (s, 1H), 7.73 (m, 6H), 7.42 (d, 1H, J=7.7), 7.16 (s, 1H), 2.92 (s, 3H), 2.29 (s, 3H). HRMS calc. for C<sub>24</sub>H<sub>23</sub>N<sub>6</sub>O<sub>3</sub>S: 475.1552; found, 475.1554.

#### Examples 113, 114, and 115

1-(3-amidinophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole trifluoroacetate, (Example 113) 1-(3-cyanophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole, (Example 114) and 1-(3-aminocarbonylphenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole (Example 115)

Part A: Preparation of 2-methylthiophenylboronic acid.

2-Bromothioanisole (29.0 g, 143 mmol) was dissolved in dry THF (400 mL) and cooled to -75°C. *N*-BuLi (62.0 mL, 2.5 M in hexane, 155 mmol) was added over 50 min. After stirring 25 min, triisopropyl borate (46 mL, 199 mmol) was added over 35 min. The cold bath was removed and the reaction was stirred at room temperature for 16 h. The resulting solution was cooled in an ice bath, and 6 M HCl (100 mL) was added. This mixture was stirred at room temp 5 h and concentrated to about half of the original volume. The concentrated solution was partitioned between Et<sub>2</sub>O and water. The organic layer was extracted with 2 M NaOH, which was subsequently reacidified with 6 M HCl and extracted several times back into Et<sub>2</sub>O. These Et<sub>2</sub>O washes were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to yield a beige solid (20.4 g, 85%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.01 (dd, 1H, J=7.3, J'=1.4), 7.53 (dd, 1H, J=7.7, J'=1.1), 7.43 (td, 1H, J=7.3, J'=1.8), 7.34 (td, 1H, J=7.3, J'=1.5), 6.22 (s, 2H), 2.50 (s, 3H).

Part B: Preparation of 2-[bis(tert-butoxycarbonyl)amino]-5-bromopyrimidine.

Sodium hydride (5.06 g, 60%, 127 mmol) was added in 2 portions to 2-amino-5-bromopyrimidine (10.0 g, 57 mmol) in dry THF (500 mL) at 0°C. After stirring 30 min, di-*t*-butyl dicarbonate (27.6 g, 126 mmol) was added. The resulting mixture was refluxed 17 h, quenched carefully with water, and concentrated. The concentrated mixture was diluted with EtOAc and extracted with water. The combined aqueous layers were extracted with EtOAc. All of the organic layers were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (10-15% EtOAc/hexanes) to yield the desired product (15.48 g, 72%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.78 (s, 2H), 1.47 (s, 18H).

Part C: Preparation of 2-[bis(tert-butoxycarbonyl)amino]-5-(2'-methylthiophenyl)pyrimidine.

2-[Bis(tert-butoxycarbonyl)amino]-5-bromopyrimidine (2.00 g, 5.3 mmol) was dissolved in benzene (130 mL). 2-methylthiophenylboronic acid (2.24 g, 13.3 mmol), aq. sodium carbonate (13 mL, 2.0 M, 26 mmol), tetrabutyl ammonium bromide (86 mg, 0.26 mmol), and bis(triphenylphosphine)palladium(II)chloride (190 mg, 0.27 mmol) were added, and the resulting mixture was purged with vacuum and argon and then refluxed 17 h. The cooled mixture was diluted with EtOAc and water. The layers were separated, and the organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (50% EtOAc/hexanes), evaporated, and chromatographed a second time on silica gel (30-50% EtOAc/hexanes) to yield the desired product (2.13 g, 96%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.81 (s, 2H), 7.41 (m, 2H), 7.25 (m, 2H), 2.39 (s, 3H), 1.49 (s, 18H).

Part D: Preparation of 2-[bis(tert-butoxycarbonyl)amino]-5-(2'-methylsulfonylphenyl)pyrimidine.

2-[Bis(tert-butoxycarbonyl)amino]-5-(2'-methylthiophenyl)pyrimidine (2.13 g, 5.1 mmol) was dissolved in MeOH (20 mL) and cooled to 0°C. In a separate beaker, a solution of Oxone (5.49 g) was generated by dilution to 27 mL with water. A portion of this solution (17 mL, 5.6 mmol) was removed and adjusted to pH 4.2 with sat. Na<sub>3</sub>PO<sub>4</sub> solution (4.7 mL). This mixture was added to the reaction and stirred 23 h at room temp. The resulting mixture was diluted with water and extracted with CHCl<sub>3</sub>. The organics were combined, washed with water and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (50-100% EtOAc/hexanes) to yield the sulfone (1.28 g, 56%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.81 (s, 2H), 8.28 (dd, 1H, J=7.6, J'=1.4), 7.72 (m, 2H), 7.39 (dd, 1H, J=7.3, J'=1.4), 2.76 (s, 3H), 1.50 (s, 18H).

Part E: Preparation of 2-amino-5-(2'-methylsulfonylphenyl)pyrimidine hydrochloride.

2-[Bis(tertbutoxycarbonyl)amino]-5-(2'-methylsulfonylphenyl)pyrimidine (1.28 g, 2.8 mmol) was suspended in HCl/dioxane (10 mL, 4.0 M) and stirred 20 h at room temp. The resulting mixture was triturated with Et<sub>2</sub>O and filtered to yield a white solid (772 mg, 95%). <sup>1</sup>HNMR(CDCl<sub>3</sub> + few drops MeOD)δ: 8.53 (s, 2H), 8.22 (dd, 1H, J=7.7, J'=1.8), 7.77 (m, 2H), 7.40 (dd, 1H, J=7.4, J'=1.5), 2.94 (s, 3H).

Part F: Preparation of 1-(3-cyanophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole.

Oxalyl chloride (175 μl, 2.0 mmol) and DMF (2 drops) were added to 1-(3-cyanophenyl)-3-methylpyrazole-5-carboxylic acid (300 mg, 1.3 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and stirred under argon for 120 min. The resulting solution was evaporated and redissolved in CH<sub>2</sub>Cl<sub>2</sub> (5 mL). 4-Dimethylaminopyridine (480 mg, 3.9 mmol) and 2-amino-5-(2'-methylsulfonylphenyl)pyrimidine hydrochloride (377 mg, 1.3 mmol) were added and stirred at room temp under argon 5 days. The crude reaction was chromatographed on silica gel (2-5% MeOH/CHCl<sub>3</sub>) to yield crude product, which was redissolved in CHCl<sub>3</sub> and extracted with 1 M HCl. The organics were dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated to yield clean product (486 mg, 80%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.69 (s, 2H), 8.64 (s, 1H), 8.25 (dd, 1H, J=7.7, J'=1.5), 7.84 (m, 1H), 7.73 (m, 4H), 7.55 (t, 1H, J=7.6), 7.35 (dd, 1H, J=7.3, J'=1.4), 6.79 (s, 1H), 2.80 (s, 3H), 2.42 (s, 3H).

Part G: Preparation of 1-(3-amidinophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole trifluoroacetate, and 1-(3-aminocarbonylphenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole.

1-(3-cyanophenyl)-3-methyl-5-([5-(2'-methylsulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole (471 mg, 1.0 mmol) was dissolved in dry CHCl<sub>3</sub> (15 mL) and dry MeOH (5 mL) and cooled to 0°C. HCl (g) was generated by the

addition of H<sub>2</sub>SO<sub>4</sub> (45 mL) to NaCl (480 g) over 30 min, and bubbled into the reaction. The generator was removed, and the reaction was sealed and placed in the refrigerator (4°C) for 18 h. The reaction was evaporated and redissolved in dry MeOH (15 mL). Ammonium carbonate (487 mg, 5.1 mmol) was added and the reaction was stirred for 20 h at room temp, and evaporated. The crude product was dissolved/suspended in a mixture of MeCN, water, TFA, DMSO, and MeOH. The soluble portion was purified by prep HPLC on a C-18 reverse phase column (10-70% MeCN/H<sub>2</sub>O/0.05% TFA) to yield the desired amidine as its TFA salt (0.31 g, 51%). <sup>1</sup>HNMR(DMSO)δ: 11.38 (s, 1H), 9.39 (s, 2H), 9.00 (s, 2H), 8.67 (s, 2H), 8.10 (dd, 1H, J=8.1, J'=1.5), 7.92 (m, 1H), 7.74 (m, 5H), 7.49 (dd, 1H, J=7.3, J'=1.1), 7.06 (s, 1H), 3.03 (s, 3H), 2.29 (s, 3H). HRMS calc. for C<sub>23</sub>H<sub>22</sub>N<sub>7</sub>O<sub>3</sub>S: 476.1505; found, 476.1529. A second product was isolated from the prep HPLC and combined with the insoluble solid from above for purification on silica gel (10% MeOH/CHCl<sub>3</sub>). The crude amide was suspended in toluene and filtered. The white solid thus obtained was the desired amide (52 mg, 11%). <sup>1</sup>HNMR(DMSO)δ: 11.33 (s, 1H), 8.64 (s, 2H), 8.08 (m, 2H), 7.92 (s, 1H), 7.77 (m, 3H), 7.48 (m, 4H), 6.95 (s, 1H), 3.01 (s, 3H), 2.27 (s, 3H). HRMS calc. for C<sub>23</sub>H<sub>21</sub>N<sub>6</sub>O<sub>4</sub>S: 477.1345; found, 477.1350.

#### Example 116

**1-(3-(N-aminoamidino)phenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt**

1-(3-Cyanophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (150 mg) was dissolved in anhydrous CH<sub>3</sub>OH and cooled to 0°C. Anhydrous HCl was bubbled through the rxn for 15 minutes. The resulting solution was allowed to warm to rt over 18 hrs. The mixture was concentrated in vacuo. LRMS (M+H)<sup>+</sup>=489 C<sub>25</sub>H<sub>23</sub>N<sub>5</sub>O<sub>4</sub>S<sub>1</sub>. 50 mg was dissolved in 10 mL of anhydrous CH<sub>3</sub>OH. Hydrazine (0.10 mL) was added and the resulting mixture was

stirred at rt for 4 hours. The mixture was concentrated under vacuo. Purification was done by HPLC yielding 2.5 mg (98% purity by HPLC). HRMS for  $C_{28}H_{31}N_7O_3S_1$  (M+H)<sup>+</sup> calc. 490.162947, found 490.164868. <sup>1</sup>HNMR(CD<sub>3</sub>OD)  $\delta$ : 1.02 (s, 9H), 2.38 (s, 3H), 5 6.94 (s, 1H), 7.305 (d, 1H, -J= 7.69 Hz), 7.53 (t, 1H, 7.69 Hz), 6.64-7.85 (m 7H), 8.085 (d, 1H, J= 8.06 Hz).

#### Example 117

1- (3- (N-aminoamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-  
10 [1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

3-[4-(2-(N-butylaminosulfonyl)phenyl)aminophenyl-3-methyl-5-carboxypyrazole]cyanophenyl (1.0 g) was dissolved in  
15 anhydrous CH<sub>3</sub>OH and cooled to 0°C. Anhydrous HCl was bubbled through the rxn for 15 minutes. The resulting solution was allowed to warm to rt over 18 hrs. The mixture was concentrated in vacuo. LRMS (M+H)<sup>+</sup>=489  $C_{25}H_{23}N_5O_4S_1$ . 300 mg was dissolved in 10 mL of anhydrous CH<sub>3</sub>OH. Hydrazine (0.023  
20 mL) was added and the resulting mixture was stirred at rt for 4 hours. The mixture was concentrated under vacuum. Purification was done by HPLC yielding 23 mg (98% purity by HPLC). HRMS for  $C_{24}H_{23}N_7O_3S_1$  (M+H)<sup>+</sup> calc. 546.228735, found 546.228088. <sup>1</sup>HNMR(CD<sub>3</sub>OD)  $\delta$ : 2.38 (s, 3H), 6.94, (s, 1H), 7.31  
25 (d, 1H, J= 7.33 Hz), 7.495 (d, 2H, J= 7.33 Hz), 7.59-7.86 (m, 7H), 8.08 (d, 1H, J= 7.69 Hz).

#### Example 118

1- (3- (N-methyl-N-hydroxyamidino)phenyl)-3-methyl-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole,  
30 trifluoroacetic acid salt

3-[4-(2-(t-butylaminosulfonyl)phenyl)amino phenyl-3-methyl-5-carboxypyrazole]cyanophenyl (300 mg) was  
35 dissolved/suspended in 25 mL of CH<sub>3</sub>OH. Triethylamine (0.098 mL) added along with N,N-methylhydroxylamine hydrochloride (0.048 g). The reaction was stirred at 50°C for 15 hours. The mixture was concentrated under vacuo. Purification was done

on silica gel using 10% CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub> as the eluent yielding 360 mg. HRMS for C<sub>29</sub>H<sub>32</sub>N<sub>6</sub>O<sub>4</sub>S<sub>1</sub> (M+H)<sup>+</sup> calc. 561.228401, found 561.22987. <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 1.02 (s, 9H), 2.38 (s, 3H), 3.40 (s, 3H), 3.62 (s, 1H), 6.96 (s, 1H), 7.305 (d, 1H, J= 7.69 Hz), 7.42 (d, 2H, J=8.79 Hz), 3.53 (t, 1H, J= 8.06 Hz), 7.60 (t, 1H, J= 7.32 Hz), 7.65 (d, 2H, J= 8.06 Hz), 7.70-7.78 (m, 4H), 8.085 (d, 1H, J= 7.69 Hz).

#### Example 119

10        1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

15        1-(3-(N-Methyl-N-hydroxy-amidino)phenyl)-3-methyl-5-[(2'-n-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (300 mg) was dissolved in acetic acid (25 mL). Trifluoroacetic anhydride (0.106 mL) was added and the reaction was stirred at rt for 35 minutes. 10% Pd/C (300 mg) was added and the reaction vessel was placed on the Parr Shaker (50 psi H<sub>2</sub>) for 17 hours. The reaction was filtered through C-lite and the mixture was concentrated under vacuum. Purification was done by HPLC yielding 33 mg (97% purity by HPLC). HRMS for C<sub>29</sub>H<sub>32</sub>N<sub>6</sub>O<sub>3</sub>S<sub>1</sub> (M+H)<sup>+</sup> calc. 545.233486, found 545.233079; <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 1.02 (s, 9H), 2.38 (s, 3H), 3.09 (s, 3H), 6.94 (s, 1H), 7.30 (d, 1H, J= 7.33 Hz), 7.425 (d, 2H, J= 8.42 Hz) 7.50 (t, 1H, J= 7.69 Hz), 7.57-7.64 (m, 3H), 7.685 (d, 1H, J= 7.32 Hz), 7.73-7.77 (m, 2H), 7.87 (s, 1H), 8.085 (d, 1H, J= 7.70 Hz).

30

#### Example 120

1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

35

1-(3'-(N-Methyl-N-hydroxy-amidino)phenyl)-3-methyl-5-[(2'-n-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (347 mg) was dissolved in trifluoroacetic acid (10 mL) and stirred at 50°C for 1 hour.

The mixture was concentrated in vacuo (346 mg). LRMS for  $C_{25}H_{24}N_6O_4S_1$   $(M+H)^+=505$ . This material (346 mg) was dissolved in acetic acid (100 mL). Trifluoroacetic anhydride (0.116 mL) was added and the reaction was stirred at rt for 35 minutes.

5 10% Pd/C (300 mg) was added and the rxn was placed on the Parr shaker (50 psi  $H_2$ ) for 18 hours. The reaction was filtered through Celite and the mixture was concentrated in vacuo. Purification was done by HPLC yielding 80 mg (98% purity by HPLC). HRMS for  $C_{25}H_{24}N_6O_3S_1$   $(M+H)^+$  calc. 489.172971, found

10 489.172971;  $^1H$ NMR( $CD_3OD$ )  $\delta$ : 2.38 (s, 3H), 3.08 (s, 3H), 6.94 (s, 1H), 7.31 (d, 1H,  $J=7.33$  Hz), 7.395 (d, 2H,  $J=8.79$  Hz) 7.51 (t, 1H,  $J=7.32$  Hz), 7.57-7.68 (m, 6H), 8.085 (d, 1H,  $J=7.47$  Hz).

15

**Example 121**

1-(3-amidinophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole, trifluoroacetic acid salt.

20

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously (Example 24);  $^1H$ NMR(DMSO)  $\delta$ : 8.40-6.95 (m, 11H); 9.25 (s, 1H); 9.50 (s, 1H); 11.55 (s, 1H). MS (ESI) 464.17  $(M+H)^+$ .

25

**Example 122**

1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole

30

The title compound was prepared as colorless crystals following the standard Pinner followed by hydrolysis and purification protocols outlined previously;  $^1H$ NMR(DMSO)  $\delta$ : 8.40-7.39 (m, 11H); 11.55 (s, 1H). MS (ESI) 465.11  $(M+H)^+$ .

35

**Example 123**

1-(3-amidinophenyl)-5-[[5-(2'-trifluoromethylphen-1-yl)pyridin-2-yl]aminocarbonyl]tetrazole, trifluoroacetic acid salt



The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 8.40-7.49 (m, 11H); 9.25 (s, 1H); 9.5 (s, 1H); 11.60 (s, 1H);  
5 MS (ESI) 453.20 (M+H)<sup>+</sup>.

#### Example 124

1-(3-amidinophenyl)-5-[(4'-bromophen-1-yl)  
aminocarbonyl]tetrazole, trifluoroacetic acid salt

10

The title compound was prepared as colorless crystals following the standard Pinner amidine reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 8.20-7.55 (m, 8H); 9.20 (s, 1H); 9.5 (s, 1H); 11.55 (s, 1H);  
15 MS (ESI) 386.03 (M+H)<sup>+</sup>.

#### Example 125

1-(3-aminocarbonylphenyl)-5-([5-(2'-trifluoromethylphen-1-yl)pyridin-2-yl]aminocarbonyl)tetrazole

20

The title compound was prepared as colorless crystals following the standard Pinner followed by hydrolysis reaction sequence and purification protocols outlined previously; <sup>1</sup>HNMR(DMSO) δ: 8.40-7.50 (m, 11H); 11.60 (s, 1H). MS (ESI)  
25 454.12 (M+H)<sup>+</sup>.

#### Example 126

5-(3-amidinophenyl)-1-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole, trifluoroacetic acid salt

30

Part A. Preparation of N-(4-bromophenylmethyl)-3-cyanobenzamide.

4-Bromobenzylamine HCl (3.36 g, 15.1 mmol) was dissolved  
35 in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Triethylamine (8.4 mL, 60 mmol) was added followed by 3-cyanobenzyl chloride (2.50 g, 15.1 mmol). The mixture was stirred at room temperature under N<sub>2</sub> for 15 min. It was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water and brine.

The CH<sub>2</sub>Cl<sub>2</sub> solution was dried over MgSO<sub>4</sub> and concentrated to 3.5 g of the desired product. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 4.60 (d, 2H); 7.0 (s, 1H); 7.20 to 8.20 (m, 8H). MS (DCI-NH<sub>3</sub>) 315 (M+H)<sup>+</sup>.

5    Part B. Preparation of 1-(4-bromophenylmethyl)-5-(3-cyanophenyl)tetrazole.

10    The material from Part A (3.2 g, 10 mmol) was dissolved in CH<sub>3</sub>CN (100 mL) and NaN<sub>3</sub> (0.7 g, 10 mmol) was added. The mixture was cooled in an ice bath and triflic anhydride (1.7 mL, 10 mmol) was added. Then, the ice bath was removed and stirred at room temperature under N<sub>2</sub> overnight. The reaction mixture was diluted with EtOAc and washed with water and brine. It was dried over MgSO<sub>4</sub>, concentrated, and  
15    chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>) to give 2.0 g of the desired product. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 5.60 (s, 2H); 7.05 to 7.90 (m, 8H). MS (NH<sub>3</sub>-CI) 340, 342 (M+H)<sup>+</sup>.

20    Part C. Preparation of 5-(3-cyanophenyl)-1-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole.

25    The material from Part B (0.36 g, 1.06 mmol), and 2-trifluoromethyl phenylboronic acid (0.24 g, 1.26 mmol) were dissolved in benzene (30 mL). The mixture was stirred at room temperature and bubble N<sub>2</sub> for 30 min. Then K<sub>2</sub>CO<sub>3</sub> (2 mL of 2 M, 4 mmol), tetrabutylammonium bromide (50 mg, 0.15 mmol) and tetrakis(triphenylphosphine)-palladium(0) (200 mg, 0.17 mmol) were added. The mixture was refluxed under N<sub>2</sub> for 4 hours. The solvent was removed. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>  
30    and washed with water and brine. It was dried over MgSO<sub>4</sub>, concentrated, and chromatographed on silica gel (eluted with CH<sub>2</sub>Cl<sub>2</sub>) to give 0.41 g of the desired product. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 5.70 (s, 2H); 7.10 to 7.85 (m, 12H). MS (NH<sub>3</sub>-CI) 406.1 (M+H)<sup>+</sup>.

35

Part D. Preparation of 5-(3-amidinophenyl)-1-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole, trifluoroacetic acid salt.

The material from part C was dissolved in 10 mL anhydrous  $\text{CHCl}_3$  and 10 mL anhydrous  $\text{CH}_3\text{OH}$ . The mixture was cooled in an ice bath and  $\text{HCl}$  gas was bubbled in until the solution was saturated. The reaction mixture was sealed and kept at  $0^\circ\text{C}$  for 12 h. The solvent was removed and the solid was dried under vacuum. The resulting solid was redissolved in 20 mL of anhydrous  $\text{CH}_3\text{OH}$  and ammonium acetate (0.77 g, 10eq) was added. The mixture was sealed and stirred at room temperature for 12 h. The solvent was removed. The solid was dissolved in  $\text{CH}_3\text{CN}/\text{H}_2\text{O}/\text{TFA}$  and purified by reverse phase HPLC to give 150.0 mg of the desired product.  $^1\text{H NMR}$  ( $\text{DMSO}-d_6$ )  $\delta$ : 5.95 (s, 1H); 7.19 to 8.20 (m, 12H); 9.35 (s, 1H); 9.50 (s, 1H). (ESI) 423.17 (M+H) $^+$ .

15

#### Example 127

1-[(3-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

20

Part A. Preparation of ethyl 1-[(3-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate.

To a solution of ethyl 3-methylpyrazole-5-carboxylate (2.0 g, 13.0 mmol) in 50 mL of dimethylformamide was added 3-cyanobenzyl bromide (2.54 g, 13.0 mmol) and potassium iodide (6.46 g, 38.9 mmol). The resulting mixture was allowed to stir at  $65^\circ\text{C}$  for 16 h. The reaction mixture was cooled to room temperature, diluted with ethyl acetate, washed with saturated aq. sodium thiosulfate (2 times) and brine (2 times), dried ( $\text{MgSO}_4$ ) and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 1:1 hexanes/ethyl acetate) to give 2.5 g (71%) of the title compound. MS (ESI) 270 (M+H) $^+$ .

35

Part B. Preparation of 1-[(3-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate.

To a solution of ethyl 1-[(3-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate (2.37 g, 8.80 mmol) in 20 mL of methanol and 20 mL of water was added sodium hydroxide (0.70 g, 17.6 mmol) and the resulting solution was stirred at room temperature for 16 h. The mixture was acidified with 10% aq HCl, diluted with ethyl acetate, washed with brine (2 times), dried (MgSO<sub>4</sub>) and concentrated in vacuo to afford the title compound (1.9 g, 90%) which was used without purification. MS (ESI) 242 (M+H)+.

Part C. Preparation of 1-[(3-cyanophenyl)methyl]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

To a solution of 1-[(3-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate (1.80 g, 7.46 mmol) in 20 mL of dimethylformamide was added (2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)amine (2.50 g, 8.21 mmol), benzotriazol-1-yloxytris(dimethylamino)phosphonium hexafluorophosphate (BOP reagent, 4.95 g, 11.19 mmol) and triethylamine (1.13 g, 11.19 mmol). The resulting mixture was stirred at 60°C for 16 h. The reaction was allowed to cool to room temperature and then was diluted with ethyl acetate, washed with brine (4 times), dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 1:1 hexanes/ethyl acetate) to afford 1.9 g (49%) of the title compound. MS (ESI) 528 (M+H)+.

Part D. Preparation of 1-[(3-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

To a solution of 1-[(3-cyanophenyl)methyl]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (1.77 g, 3.35 mmol) in 40 mL of methyl acetate was added anhydrous methanol (1.36 mL, 33.5 mmol). The solution was cooled to 0°C. Then anhydrous HCl was bubbled through the solution for 15 minutes. The solution was stoppered and allowed to stir overnight

at room temperature. The volatiles were removed in vacuo. The residue was dried under high vacuum for 1 hr. The residue was then dissolved in 100 mL of anhydrous methanol. Ammonium carbonate (1.93 g, 20.21 mmol) was added and the reaction was stirred overnight at room temperature. The volatiles were removed in vacuo and the residue was purified by preparative HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) to yield the title compound as a white powder. MS (ESI) 489 (M+H)<sup>+</sup>.

#### Example 128

1-[(4-Amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

Part A. Preparation of ethyl 1-[(4-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate.

Ethyl-3-methylpyrazole-5-carboxylate (2.50 g, 16.21 mmol) was allowed to react with 4-cyanobenzyl bromide (3.18 g, 16.21 mmol) and potassium iodide (8.07 g, 48.65 mmol) to afford 3.1 g (70%) of the title compound. MS (ESI) 270 (M+H)<sup>+</sup>.

Part B. Preparation of 5-carboxy-1-[(4-cyanophenyl)methyl]-3-methylpyrazole.

Ethyl-1-[(4-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate (2.96 g, 10.99 mmol) was converted into 2.4 g (91%) of the title compound following the procedure outlined previously; MS (ESI) 242 (M+H)<sup>+</sup>.

Part C. Preparation of 1-[(4-cyanophenyl)methyl]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

5-carboxy-1-[(4-cyanophenyl)methyl]-3-methylpyrazole-5-carboxylate (2.29 g, 9.49 mmol) was converted into 2.0 g (40%)

of the title compound following the procedure outlined previously; MS (ESI) 528 (M+H)+.

Part D. Preparation of 1-[(4-amidinophenyl)methyl]-3-methyl-5-  
5 [(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

1-[(4-cyanophenyl)methyl]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (0.78  
10 g, 1.47 mmol) was converted into the title compound following methods described previously; MS (ESI) 489 (M+H)+.

#### Example 129

1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-  
15 yl)aminocarbonyl]imidazole, trifluoroacetic acid salt

Part A: 3-Fluorobenzonitrile (4.84 g, 40 mmol) was heated with imidazole (2.72 g, 40 mmol) in the presence of K<sub>2</sub>CO<sub>3</sub> in DMF at 100°C for 8 hours to afford the coupled product  
20 as a white solid in quantitative yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.89 (s, 1H), 7.70 (d, J=0.8 Hz, 1H), 7.68-7.58 (m, 3H), 7.30 (d, J=1.0 Hz, 1H), 7.26 (s, 1H); LRMS: 170 (M+H)+.

Part B: Product from part A (1.52 g, 9 mmol) was slowly  
25 treated with n-BuLi (1.6 M, 6.3 mL) in THF (60 mL) at -78°C for 40 minutes and was then slowly quenched with chloromethylformate (942 mg, 10 mmol) at this temperature. The resulting mixture was stirred at -78°C and warmed to room temperature over 2 hours. Then poured into water and  
30 ethyl acetate. The organic layer was separated and washed with water, brine, and dried over MgSO<sub>4</sub>. After removal of the ethyl acetate, a residue was purified by column chromatography with ethyl acetate and methylene chloride (1:1) to afford the 2-imidazolylphenylethylester  
35 derivative (1.33 g, 65%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.80-7.77 (m, 1H), 7.65-7.61 (m, 3H), 7.33 (s, 1H), 7.20 (s, 1H); LRMS: 228 (M+H)+.

Part C: To a stirred solution of 4-[(o-SO<sub>2</sub>tBu)-phenyl]aniline (304 mg, 1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was slowly added trimethylaluminum (2M in hexane, 1 mL) at 0°C and the resulting mixture was warmed to room temperature over 15 minutes. The product from part B in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added dropwise and the resulting mixture was refluxed for 2 hours. The mixture was quenched with water, diluted with ethyl acetate and filtered through Celite. The organic layer was separated, washed with water, brine and dried over MgSO<sub>4</sub>. After removal of the ethyl acetate, a residue was purified by column chromatography with ethyl acetate and methylene chloride (1:1) as eluent to afford the coupled product (260 mg, 52%) as a white solid.

<sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 9.41 (s, 1H), 8.15 (dd, J=7.7 Hz, J=1.5 Hz, 1H), 7.78 (dd, J=7.3 Hz, J=1.1 Hz, 1H), 7.74-7.57 (m, 6H), 7.55 (td, J=7.7 Hz, J=1.1 Hz, 1H), 7.49 (dd, J=8.8 Hz, J=1.8 Hz, 2H), 7.29 (dd, J=8.1 Hz, J=1.5 Hz, 1H), 7.28 (d, J=0.8 Hz, 1H), 7.22 (d, J=0.8 Hz, 1H), 3.64 (s, 1H), 0.99 (s, 9H); LRMS: 500.1 (M+H)<sup>+</sup>.

20

Part D: Standard Pinner amidine and purification methods then afforded the titled product (120 mg, 50%):

<sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 8.08 (dd, J=7.7 Hz, J=1.1 Hz, 1H), 7.91-7.88 (m, 2H), 7.83 (dd, J=8.4 Hz, J=1.5 Hz, 1H), 7.74 (t, J=8.0 Hz, 1H), 7.65 (d, J=8.4 Hz, 2H), 7.58 (dd, J=7.3 Hz, J=1.1 Hz, 1H), 7.50 (s, 2H), 7.39 (d, J=8.4 Hz, 2H), 7.32 (s, 1H), 7.30 (dd, J=7.3 Hz, J=1.1 Hz, 1H); ESMS: 461 (M+H)<sup>+</sup>

30

#### Example 130

1-(3-amidinophenyl)-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole

Ethyl-1-(3-cyanophenyl)-4-methyl-imidazolyl-2-carboxylate was prepared following the standard coupling procedure outlined previously. This was coupled following the standard Weinreb conditions (trimethylaluminum) and subjected to the Pinner amidine reaction protocols

followed by usual methods of purification to afford the title compound as colorless crystals.  $^1\text{H NMR}(\text{CD}_3\text{OD})\delta$ : 8.09 (dd,  $J=8.1$  Hz,  $J=1.1$  Hz, 1H), 7.89 (t,  $J=1.5$  Hz, 1H), 7.87 (d,  $J=1.8$  Hz, 1H), 7.81-7.78 (m, 1H), 7.72 (t,  $J=8.1$  Hz, 1H), 7.64 (d,  $J=8.4$  Hz, 2H), 7.57 (dd,  $J=7.7$  Hz,  $J=1.5$  Hz, 1H), 7.50 (td,  $J=7.7$  Hz,  $J=1.5$  Hz, 1H), 7.38 (d,  $J=8.4$  Hz, 2H), 7.30 (dd,  $J=7.7$  Hz,  $J=1.5$  Hz, 1H), 7.26 (s, 1H), 2.33 (s, 3H);  $^{13}\text{C NMR}(\text{CD}_3\text{OD})\delta$ : 167.73, 158.04, 143.04, 141.49, 140.47, 139.62, 138.64, 137.53, 133.65, 133.45, 132.93, 132.76, 132.35, 131.25, 130.55, 129.09, 128.74, 128.63, 126.69, 120.87, 13.27; ESMS:  $m/z$  475.19 ( $M+H$ , 100); HRMS: calcd. for  $\text{C}_{24}\text{H}_{23}\text{N}_6\text{O}_3\text{S}_1$  475.1552 found 475.1548.

#### Example 131

15        1-(3-amidinophenyl)-5-chloro-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole

Chlorination of ethyl-1-(3-cyanophenyl)-4-methyl-imidazole-2-carboxylate with NCS in refluxing carbontetrachloride afforded the 5-chloroimidazole derivative which was then subjected to the Pinner amidine reaction protocols followed by usual methods of purification to afford the title compound as colorless crystals (145 mg, 34.8%).  $^1\text{H NMR}(\text{CD}_3\text{OD})\delta$ : 8.07 (d,  $J=7.7$  Hz, 1H), 7.96 (d,  $J=7.3$  Hz, 1H), 7.82 (s, 1H), 7.78 (d,  $J=7.7$  Hz, 1H), 7.73 (d,  $J=8.1$  Hz, 1H), 7.61 (d,  $J=8.5$  Hz, 2H), 7.57 (d,  $J=8.8$  Hz, 1H), 7.49 (t,  $J=7.7$  Hz, 1H), 7.37 (d,  $J=8.4$  Hz, 2H), 7.29 (d,  $J=7.7$  Hz, 1H), 2.32 (s, 3H);  $^{13}\text{C NMR}(\text{CD}_3\text{OD})\delta$ : 167.63, 157.41, 143.05, 141.47, 139.26, 138.46, 138.32, 137.59, 135.51, 134.27, 133.63, 132.91, 131.48, 131.22, 130.84, 129.98, 128.74, 128.61, 128.43, 120.98, 12.22; ESMS:  $m/z$  509.1 ( $M+H$ , 100); HRMS: calcd. for  $\text{C}_{24}\text{H}_{22}\text{Cl}_1\text{N}_6\text{O}_3\text{S}_1$  509.1163 found 509.1172.

35

#### Example 132

5-(3-amidinophenyl)-2-methyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole



Part A: Ethyl-2-methyl-4-(3'-cyano)phenyl-5-carboxylate was prepared via the reaction of ethyl-2-bromo-(3-cyano)benzoylacetate and ammonium acetate in acetic acid in 20% yield.  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 10.03 (BS, 1H), 8.25 (bs, 1H), 8.17 (bd, 1H), 7.40 (d, 1H), 7.44 (t, 1H), 4.30 (q, 2H), 2.50 (s, 3H), 1.30 (t, 3H) ppm; Ammonia CI mass spectrum 272 (M+H, 100).

Part B: Weinreb coupling of the product from part A with the -2'-tert-butylaminosulfonyl-1-aminobiphenyl and trimethyl aluminum afforded the desired coupled product which when subjected to the standard Pinner amidine reaction and the usual purification protocols to afford the title compound as colorless crystals;  $^1\text{H NMR}(\text{CD}_3\text{OD})$   $\delta$ : 8.29 (s, 1H), 8.10 (dd, J=7.9 Hz, J=1.2 Hz, 1H), 8.06 (d, J=7.8 Hz, 1H), 7.83 (d, J=7.6 Hz, 1H), 7.73 (d, J=7.8 Hz, 1H), 7.70 (bs, 2H), 7.61 (td, J=7.6 Hz, J=1.5 Hz, 1H), 7.52 (td, J=7.6 Hz, J=1.5 Hz, 1H), 7.42 (d, J=6.8 Hz, 2H), 7.33 (dd, J=7.6 Hz, J=1.2 Hz, 1H), 2.53 (bs, 3H); ESMS: m/z 475.1 (M+H, 100) for  $\text{C}_{24}\text{H}_{22}\text{N}_6\text{O}_3\text{S}_1$ .

#### Examples 133 and 134

1-(3-amidinophenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole and 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole

Part A. Preparation of N-(4-nitrophenyl)benzimidazole.

Made a suspension of 1.26 g of 4-bromonitrobenzene and 0.74 g of benzimidazole in 50 mL of anhydrous dimethylformamide. Added 0.94 g of potassium carbonate to reaction mixture. Warmed reaction mixture to 80°C for 72H. Diluted reaction mixture with 100 mL water and extracted 3 times with 50 mL ethyl acetate portions. Combined extracts and dried. Filtered and concentrated resulting organics in vacuo to give the crude product. LRMS ( $\text{NH}_3$ -CI): 240, (M+H,

100),  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.50 (d, 2H), 8.20 (s, 1H), 7.93 (complex, 1H), 7.75 (d, 2H), 7.63 (complex, 1H), 7.42 (complex, 2H).

Part B. Preparation of N-(4-aminophenyl)benzimidazole.

5

Made a suspension of 0.6 g crude N-(4-nitrophenyl)benzimidazole and a catalytic amount of 10% palladium on carbon in 20 mL methanol. Placed reaction mixture under 1 atmosphere of hydrogen and let stir for 15H.

10 Passed reaction mixture through a 1" celite pad and concentrated filtrate in vacuo to give the crude product. LRMS ( $\text{NH}_3\text{-CI}$ ): 210 (M+H, 100),  $^1\text{H NMR}(\text{DMSO-d}_6)\delta$ : 9.25 (s, 1H), 7.83 (complex, 1H), 7.60 (complex, 1H), 7.47 (complex, 2H), 7.35 (d, 2H), 6.80 (d, 2H).

15

Part C. Preparation of N-(3-cyanophenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole.

To 0.16 g of N-(3-cyanophenyl)-3-methyl-pyrazole 5-carboxylic acid and 25 mL dichloromethane was added 0.07 mL oxalyl chloride and 2 drops DMF. The reaction proceeded overnight. Concentration of the reaction mixture and placement under high vacuum gave the crude acid chloride which was then coupled to the product of part B under standard conditions to afford crude N-(3-cyanophenyl)-3-methyl-5-[(4'-N-benzimidazol-1-yl-phenyl)aminocarbonyl]pyrazole. LRMS (ESI): 419 (M+H, 20), 210 (M+2H) $^{++}$ .

Part D. Preparation of N-(3-amidinophenyl)-3-methyl-5-[(4'-benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole.

Standard transformation of the benzonitrile obtained in part C to the benzamidine via the ethyl imidate converted 0.24 g of the crude benzonitrile to 0.02 g of the benzamidine bis-TFA salt after standard HPLC purification. LRMS (ES $^{+}$ ): 436.21 (M+H), HRMS (FAB): Calc: 436.188584 Mass: 436.191317 and 0.003 g of the benzamide LRMS (ES $^{+}$ ): 437 (M+H), 459 (M+Na), HRMS ( $\text{NH}_3\text{-CI}$ ): Calc: 437.172599 Mass: 437.173670.  $^1\text{H NMR}(\text{DMSO-}$

d6, 300MHz)  $\delta$ : 10.76 (s, 1H), 9.40 (s, 2H), 9.02 (s, 2H), 8.59 (s, 1H), 7.94 (s, 1H), 7.88 (d, 2H), 7.76 (complex, 3H), 7.64 (complex, 4H), 7.32 (complex, 2H), 7.05 (s, 1H), 2.30 (s, 3H).

5

#### Examples 135 and 136

1-(3-amidinophenyl)-3-methyl-5-[(4'-(2-methylimidazolyl)phenyl)aminocarbonyl]pyrazole and 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(2-methylimidazolyl)phenyl)aminocarbonyl]pyrazole

10

Part A. Preparation of N-(4-nitrophenyl)-2-methylimidazole.

2-Methylimidazole (1.04 g) was treated with 0.56 g 60% sodium hydride in an oil dispersion in 60 mL DMF with cooling. After 0.33H added 4-bromonitrobenzene in three portions over 0.5 H. Let reaction mixture warm to ambient temperature overnight. Diluted mixture with 100 mL of 1.0M HCl solution and extracted three times with 30 mL portions of ethyl acetate. Combined extracts and dried over magnesium sulfate. Concentrated resulting organics in vacuo. Purified crude material by standard chromatographic techniques to give the purified product as a crystalline solid. LRMS (NH<sub>3</sub>-CI): 204 (M+H, 100); <sup>1</sup>HNMR(CDCl<sub>3</sub>)  $\delta$ : 8.40 (d, 2H), 7.50 (d, 2H), 7.05 (d, 2H), 2.43 (s, 3H).

25

Part B. Preparation of 1-(4-aminophenyl)-2-methylimidazole.

N-(4-nitrophenyl)-2-methylimidazole (0.47 g) was treated with a catalytic amount of 10% palladium on carbon in 15 mL methanol. The mixture was placed under an atmosphere of hydrogen. The reaction mixture was stirred for 3H and then passed through a 1" celite pad. The resulting filtrate was concentrated under reduced pressure to give the title compound. LRMS (NH<sub>3</sub>-CI): 174 (M+H, 100), <sup>1</sup>HNMR(CDCl<sub>3</sub>)  $\delta$ : 7.05 (d, 1H), 6.97 (d, 2H), 6.77 (d, 1H), 6.60 (d, 2H), 5.34 (s, 2H), 2.13 (s, 3H).

35

Part C. Preparation of N-(3-cyanophenyl)-3-methyl-5-((4'-2-methylimidazolylphenyl)aminocarbonyl)pyrazole.

To 0.24 g of N-(3-cyanophenyl)-3-methyl-pyrazole 5-carboxylic acid and 20 mL dichloromethane was added 0.14 mL oxalyl chloride and 2 drops DMF. The reaction proceeded overnight. Concentration of the reaction mixture and placement under high vacuum gave the crude acid chloride which was then coupled to the product of part B under standard conditions to afford the title compound isolated as the hydrochloride salt. LRMS (ESI):383 (M+H, 100), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.90 (s, 1H), 7.95 (s, 1H), 7.90 (d, 2H), 7.83 (m, 2H), 7.75 (m, 2H), 7.63 (m, 1H), 7.57 (d, 2H), 7.10 (s, 1H), 2.49 (s, 3H), 2.30 (s, 3H).

Part D. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-2-methylimidazolylphenyl)aminocarbonyl]pyrazole and 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-2-methylimidazolylphenyl)aminocarbonyl]pyrazole.

The N-(3-cyanophenyl)-3-methyl-5-((4'-2-methylimidazolylphenyl)aminocarbonyl)pyrazole was converted to the corresponding benzamidine via Pinner synthesis and amidination by subsequent treatment of the imide with ammonium carbonate. The crude mixture was then purified by standard HPLC technique to give the benzamidine as a white solid after lyophilization LRMS (ES+):400 (M+H, 100); HRMS: Calc:400.188584, Mass:400.188113 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.87 (s, 1H), 9.40 (s, 2H), 9.30 (s, 2H), 7.95 (s, 1H), 7.89 (d, 2H), 7.80 (m, 2H), 7.75 (m, 2H), 7.65 (m, 1H), 7.55 (d, 2H), 7.05 (s, 1H) 2.47 (s, 3H), 2.30 (s, 3H). The corresponding benzamide was isolated as a by-product during purification. LRMS (ES+): 401 (M+H) HRMS (NH<sub>3</sub>-CI): Calc. 401.172599 Mass: 410.170225; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.77 (s, 1H), 8.78 (s, 1H), 8.09 (s, 1H), 7.94 (s, 1H), 7.87 (m, 3H), 7.77 (m, 1H), 7.65 (d, 2H), 7.63 (m, 1H), 7.50 (complex, 3H), 7.36 (m, 2H), 6.95 (s, 1H), 2.30 (s, 3H).

## Example 137

1-(3-amidinophenyl)-3-methyl-5-[[4'-(1,2,4-triazol-2-yl)-phenyl]aminocarbonyl]pyrazole

- 5 Part A. 1-(3-cyanophenyl)-3-methyl-5-((4'-(1,2,4-triazolyl)phenyl)aminocarbonyl)pyrazole.

The pyrazole acid chloride was generate by standard method and coupled to 0.18 g of commercially available 4-(1-N-  
10 1, 2, 4-triazolo)aniline using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-((4'-(1,2,4-triazol-1-yl)phenyl)aminocarbonyl)pyrazole. The crude product was recrystallized from 2:1 methylene chloride to methanol to give the product as a white solid. LRMS (NH<sub>3</sub>-CI):370 (M+H),  
15 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.57 (s, 1H), 9.20 (s, 1H), 8.19 (s, 1H), 7.97 (s, 1H), 7.80 (complex, 6H), 7.65 (t, 1H), 7.00 (s, 1H), 2.29 (s, 3H).

- 20 Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-((4'-(1,2,4-triazolyl)phenyl)aminocarbonyl)pyrazole.

Standard transformation of the benzonitrile obtained in part A to the benzamidine via the ethyl imidate converted 0.13 g of the benzonitrile to the benzamidine bis-TFA salt after  
25 standard HPLC purification. LRMS (ES<sup>+</sup>): 387 (M+H) HRMS (NH<sub>3</sub>-CI): Calc: 387.168182 Mass: 387.166790; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.70 (s, 1H), 9.39 (s, 2H), 9.20 (2, 1H), 9.02 (s, 2H), 8.19 (s, 1H), 7.91 (s, 1H), 7.79 (m, 5H), 7.70 (m, 2H), 7.02 (s, 1H), 2.31 (s, 3H).

30

## Example 138

1-(3-amidinophenyl)-3-methyl-5-((4'-cyclohexylphenyl)aminocarbonyl)pyrazole

- 35 Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-((4'-cyclohexylphenyl)aminocarbonyl)pyrazole.

The pyrazole acid chloride was generate in the standard method and coupled to 0.19 g of commercially available 4-cyclohexylaniline using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-((4'-

5 cyclohexylphenyl)aminocarbonyl)pyrazole. LRMS (NH<sub>3</sub>-CI):385 (M+H), 402 (M+NH<sub>4</sub>), <sup>1</sup>HNMR(DMSO, 300MHz)δ: 10.40 (s, 1H), 7.92 (s, 1H), 7.82 (d, 1H), 7.72 (d, 1H), 7.61 (t, 1H), 7.50 (d, 2H), 7.13 (d, 2H), 6.92 (s, 1H), 3.31 (s, 1H), 2.25 (s, 3H), 1.71 (complex, 5H), 1.13 (complex, 5H).

10

Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-((4'-cyclohexylphenyl)aminocarbonyl)pyrazole.

Standard transformation of the benzonitrile obtained in part A to the benzamidine via the ethyl imidate converted the 15 crude benzonitrile to the benzamidine TFA salt. The crude product was purified by standard HPLC purification. LRMS (ES<sup>+</sup>): 402 (M+H) HRMS (NH<sub>3</sub>-CI): Calc: 402.229386 Mass: 402.227504 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.30 (s, 1H), 9.38 (s, 20 2H), 9.07 (s, 2H), 7.90 (s, 1H), 7.77 (m, 1H), 7.69 (m, 2H), 7.50 (d, 2H), 7.12 (d, 2H), 6.93 (s, 1H), 3.31 (m, 1H), 2.28 (s, 3H), 1.71 (complex, 5H), 1.32 (complex, 5H).

#### Example 139

25 1-(3-amidinophenyl)-3-methyl-5-[[1,1']-biphen-4-ylaminocarbonyl]pyrazole

Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[[1,1']-biphen-4-ylaminocarbonyl]pyrazole.

30

The pyrazole acid chloride was generate by standard method and coupled to 0.19 g of commercially available 4-aminobiphenyl using standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-[[1,1']-biphen-4-ylaminocarbonyl]pyrazole. LRMS (NH<sub>3</sub>-CI):379 (M+H), 396 (M+NH<sub>4</sub>) 35 HRMS (NH<sub>3</sub>-CI): Calc:396.182436 Mass:396.181736. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.57 (s, 1H), 9.20 (s, 1H), 8.19 (s, 1H), 7.97 (s,

1H), 7.80 (complex, 6H), 7.65 (t, 1H), 7.00 (s, 1H), 2.29 (s, 3H).

Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-  
5 [[1,1']-biphen-4-ylaminocarbonyl]pyrazole.

Standard transformation of the benzonitrile obtained in part B to the benzamidine via the ethyl imidate converted of the crude benzonitrile to the benzamidine TFA salt. The  
10 crude product was purified by standard HPLC purification technique. LRMS (ES+): 396 (M+H) HRMS (NH3-CI): Calc: 396.181736 Mass: 396.182436; <sup>1</sup>HNMR(DMSO, 300MHz)δ: 10.60 (s, 1H), 9.40 (s, 2H), 8.99 (s, 2H), 7.91 (m, 1H), 7.80 (complex, 5H), 7.61 (m, 4H), 7.41 (m, 2H), 7.30 (m, 1H), 7.00 (s, 1H),  
15 2.29 (s, 3H).

Example 140

1-(3-amidinophenyl)-3-methyl-5-((4'-  
morpholinophenyl)aminocarbonyl)pyrazole

20

Part A. 1-(3-cyanophenyl)-3-methyl-5-((4'-  
morpholinophenyl)aminocarbonyl)pyrazole.

The pyrazole acid chloride was generate from the pyrazole  
25 acid by standard method and coupled to 0.26 g of commercially available 4-morpholinoaniline using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-((4'-  
morpholinophenyl)aminocarbonyl)pyrazole. LRMS (NH3-CI):388 (M+H), <sup>1</sup>HNMR(DMSO, 300MHz)δ: 10.30 (s, 1H), 7.90 (m, 1H), 7.82  
30 (d, 1H), 7.71 (m, 1H), 7.62 (t, 6H), 7.49 (d, 2H), 6.89 (s, 1H), 6.87 (d, 2H), 3.69 (t, 4H), 3.02 (t, 4H), 2.25 (s, 3H).

Part B. Standard transformation of the benzonitrile obtained in part A to the benzamidine via the ethyl imidate converted  
35 the crude benzonitrile to the benzamidine bis-TFA salt. The crude product was purified by standard HPLC purification. LRMS (ES+): 405 (M+H) HRMS (NH3-CI): Calc: 405.203899 Mass: 405.201545 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.38 (s, 1H), 9.40 (s,

2H), 9.12 (s, 2H), 7.90 (s, 1H), 7.78 (d, 1H), 7.68 (m, 2H), 7.49 (d, 2H), 6.92 (s, 1H), 6.90 (d, 2H), 3.80 (t, 4H), 3.01 (t, 4H), 2.29 (s, 3H).

5

**Example 141**

1-(3-amidinophenyl)-3-methyl-5-[(4'-((2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole

Part A. Preparation 4-(2-trifluoromethyltetrazolyl)nitrobenzene.

3.0 g of commercially available 4-nitroaniline was trifluoromethylacetylated in the presence of trifluoroacetic anhydride to give the crude N-trifluoroacetyl-4-nitroaniline. LRMS (NH<sub>3</sub>-CI): 252 (M+NH<sub>4</sub>); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 11.75 (s, 1H), 8.28 (d, 2H), 7.92 (d, 2H). The crude material was then treated with triphenylphosphine in carbon tetrachloride to give the chloroimine. <sup>1</sup>HNMR(CDCl<sub>3</sub>, 300MHz)δ: 8.35 (d, 2H), 7.15 (d, 2H). The crude chloroimine was cyclized to the 4-(2-trifluoromethyltetrazole)nitrobenzene with sodium azide in acetonitrile. <sup>1</sup>HNMR(CDCl<sub>3</sub>, 300MHz)δ: 8.54 (d, 2H), 7.80 (d, 2H). The crude 2-trifluoromethyltetrazoloaniline was triturated to give the semi-crude product which was catalytically reduced to the aniline with 10% palladium on carbon. LRMS (NH<sub>4</sub>-CI): 230 (M+H), 247 (M+NH<sub>4</sub>), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 7.256 (d, 2H), 6.65 (d, 2H).

Part B. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-((2-trifluoromethyl)tetrazolyl)phenyl)aminocarbonyl]pyrazole.

30

The pyrazole acid chloride was generated by standard method and coupled to 0.49 g of 4-(2-trifluoromethyltetrazolo)aniline using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-[(4'-((2-trifluoromethyltetrazol)-1-yl-phenyl)aminocarbonyl]pyrazole. LRMS (NH<sub>3</sub>-CI): 439 (M+H), 461 (M+Na<sup>+</sup>), 877 (2 M+H), 899 (2M+Na); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.87 (s, 1H), 8.00 (s, 1H), 7.91 (d,



2H), 7.84 (m, 1H), 7.77 (m, 1H), 7.69 (d, 2H), 7.63 (t, 1H), 7.02 (s, 1H), 2.29 (s, 3H).

Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-  
5 ((2-trifluoromethyl)tetrazolyl)phenyl)aminocarbonyl]pyrazole.

Standard transformation of the benzonitrile obtained in part B to the benzamidine via the ethyl imidate converted the crude benzonitrile to the benzamidine TFA salt after HPLC  
10 purification. LRMS (ES+): 456 (M+H) HRMS (NH<sub>3</sub>-CI): Calc: 456.150816 Mass: 456.150428; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.92 (s, 1H), 9.40 (s, 2H), 9.18 (s, 2H), 7.90 (complex, 3H), 7.78 (m, 2H), 7.67 (complex, 3H), 7.08 (s, 1H), 2.32 (s, 3H).

15

#### Example 142

1-(3-aminomethylphenyl)-3-methyl-5-[(4'-((2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole

0.06 g of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(2-trifluoromethyltetrazolyl)phenyl)aminocarbonyl]pyrazole was  
20 reacted with 10% palladium on carbon in TFA/methanol under a hydrogen atmosphere. After a few hours the reaction mixture was filtered through a 1 inch celite pad. The filtrate was concentrated under reduced pressure and the residue was  
25 purified by standard HPLC method to give the desired compound. LRMS (NH<sub>4</sub>-CI): 443 (M+H) HRMS (NH<sub>4</sub>-CI): calc: 443.155567 mass: 443.155567; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.90 (s, 1H), 8.20 (brd. s, 2H), 7.90 (d, 2H), 7.69 (d, 2H), 7.62 (s, 1H), 7.42 (complex, 3H), 6.97 (s, 1H), 4.09 (m, 2H), 2.29 (s, 3H).

30

#### Example 143

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-dimethylamino)carbonylamino)phen-1'-yl)aminocarbonyl]pyrazole

35 Part A. Preparation of 4-((N,N-dimethylamino)carbonylamino)-1-nitrobenzene.

1.56 g of 4-nitroaniline was treated with 0.50 g sodium hydride in 60% oil dispersion in DMF at 0°C. After 20 minutes added 1.04 mL of N,N-dimethylcarbonyl chloride dropwise. Let mixture warm to ambient temperature overnight. 5  
5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55  
60  
65  
70  
75  
80  
85  
90  
95  
100  
105  
110  
115  
120  
125  
130  
135  
140  
145  
150  
155  
160  
165  
170  
175  
180  
185  
190  
195  
200  
205  
210  
215  
220  
225  
230  
235  
240  
245  
250  
255  
260  
265  
270  
275  
280  
285  
290  
295  
300  
305  
310  
315  
320  
325  
330  
335  
340  
345  
350  
355  
360  
365  
370  
375  
380  
385  
390  
395  
400  
405  
410  
415  
420  
425  
430  
435  
440  
445  
450  
455  
460  
465  
470  
475  
480  
485  
490  
495  
500  
505  
510  
515  
520  
525  
530  
535  
540  
545  
550  
555  
560  
565  
570  
575  
580  
585  
590  
595  
600  
605  
610  
615  
620  
625  
630  
635  
640  
645  
650  
655  
660  
665  
670  
675  
680  
685  
690  
695  
700  
705  
710  
715  
720  
725  
730  
735  
740  
745  
750  
755  
760  
765  
770  
775  
780  
785  
790  
795  
800  
805  
810  
815  
820  
825  
830  
835  
840  
845  
850  
855  
860  
865  
870  
875  
880  
885  
890  
895  
900  
905  
910  
915  
920  
925  
930  
935  
940  
945  
950  
955  
960  
965  
970  
975  
980  
985  
990  
995  
1000  
1005  
1010  
1015  
1020  
1025  
1030  
1035  
1040  
1045  
1050  
1055  
1060  
1065  
1070  
1075  
1080  
1085  
1090  
1095  
1100  
1105  
1110  
1115  
1120  
1125  
1130  
1135  
1140  
1145  
1150  
1155  
1160  
1165  
1170  
1175  
1180  
1185  
1190  
1195  
1200  
1205  
1210  
1215  
1220  
1225  
1230  
1235  
1240  
1245  
1250  
1255  
1260  
1265  
1270  
1275  
1280  
1285  
1290  
1295  
1300  
1305  
1310  
1315  
1320  
1325  
1330  
1335  
1340  
1345  
1350  
1355  
1360  
1365  
1370  
1375  
1380  
1385  
1390  
1395  
1400  
1405  
1410  
1415  
1420  
1425  
1430  
1435  
1440  
1445  
1450  
1455  
1460  
1465  
1470  
1475  
1480  
1485  
1490  
1495  
1500  
1505  
1510  
1515  
1520  
1525  
1530  
1535  
1540  
1545  
1550  
1555  
1560  
1565  
1570  
1575  
1580  
1585  
1590  
1595  
1600  
1605  
1610  
1615  
1620  
1625  
1630  
1635  
1640  
1645  
1650  
1655  
1660  
1665  
1670  
1675  
1680  
1685  
1690  
1695  
1700  
1705  
1710  
1715  
1720  
1725  
1730  
1735  
1740  
1745  
1750  
1755  
1760  
1765  
1770  
1775  
1780  
1785  
1790  
1795  
1800  
1805  
1810  
1815  
1820  
1825  
1830  
1835  
1840  
1845  
1850  
1855  
1860  
1865  
1870  
1875  
1880  
1885  
1890  
1895  
1900  
1905  
1910  
1915  
1920  
1925  
1930  
1935  
1940  
1945  
1950  
1955  
1960  
1965  
1970  
1975  
1980  
1985  
1990  
1995  
2000  
2005  
2010  
2015  
2020  
2025  
2030  
2035  
2040  
2045  
2050  
2055  
2060  
2065  
2070  
2075  
2080  
2085  
2090  
2095  
2100  
2105  
2110  
2115  
2120  
2125  
2130  
2135  
2140  
2145  
2150  
2155  
2160  
2165  
2170  
2175  
2180  
2185  
2190  
2195  
2200  
2205  
2210  
2215  
2220  
2225  
2230  
2235  
2240  
2245  
2250  
2255  
2260  
2265  
2270  
2275  
2280  
2285  
2290  
2295  
2300  
2305  
2310  
2315  
2320  
2325  
2330  
2335  
2340  
2345  
2350  
2355  
2360  
2365  
2370  
2375  
2380  
2385  
2390  
2395  
2400  
2405  
2410  
2415  
2420  
2425  
2430  
2435  
2440  
2445  
2450  
2455  
2460  
2465  
2470  
2475  
2480  
2485  
2490  
2495  
2500  
2505  
2510  
2515  
2520  
2525  
2530  
2535  
2540  
2545  
2550  
2555  
2560  
2565  
2570  
2575  
2580  
2585  
2590  
2595  
2600  
2605  
2610  
2615  
2620  
2625  
2630  
2635  
2640  
2645  
2650  
2655  
2660  
2665  
2670  
2675  
2680  
2685  
2690  
2695  
2700  
2705  
2710  
2715  
2720  
2725  
2730  
2735  
2740  
2745  
2750  
2755  
2760  
2765  
2770  
2775  
2780  
2785  
2790  
2795  
2800  
2805  
2810  
2815  
2820  
2825  
2830  
2835  
2840  
2845  
2850  
2855  
2860  
2865  
2870  
2875  
2880  
2885  
2890  
2895  
2900  
2905  
2910  
2915  
2920  
2925  
2930  
2935  
2940  
2945  
2950  
2955  
2960  
2965  
2970  
2975  
2980  
2985  
2990  
2995  
3000  
3005  
3010  
3015  
3020  
3025  
3030  
3035  
3040  
3045  
3050  
3055  
3060  
3065  
3070  
3075  
3080  
3085  
3090  
3095  
3100  
3105  
3110  
3115  
3120  
3125  
3130  
3135  
3140  
3145  
3150  
3155  
3160  
3165  
3170  
3175  
3180  
3185  
3190  
3195  
3200  
3205  
3210  
3215  
3220  
3225  
3230  
3235  
3240  
3245  
3250  
3255  
3260  
3265  
3270  
3275  
3280  
3285  
3290  
3295  
3300  
3305  
3310  
3315  
3320  
3325  
3330  
3335  
3340  
3345  
3350  
3355  
3360  
3365  
3370  
3375  
3380  
3385  
3390  
3395  
3400  
3405  
3410  
3415  
3420  
3425  
3430  
3435  
3440  
3445  
3450  
3455  
3460  
3465  
3470  
3475  
3480  
3485  
3490  
3495  
3500  
3505  
3510  
3515  
3520  
3525  
3530  
3535  
3540  
3545  
3550  
3555  
3560  
3565  
3570  
3575  
3580  
3585  
3590  
3595  
3600  
3605  
3610  
3615  
3620  
3625  
3630  
3635  
3640  
3645  
3650  
3655  
3660  
3665  
3670  
3675  
3680  
3685  
3690  
3695  
3700  
3705  
3710  
3715  
3720  
3725  
3730  
3735  
3740  
3745  
3750  
3755  
3760  
3765  
3770  
3775  
3780  
3785  
3790  
3795  
3800  
3805  
3810  
3815  
3820  
3825  
3830  
3835  
3840  
3845  
3850  
3855  
3860  
3865  
3870  
3875  
3880  
3885  
3890  
3895  
3900  
3905  
3910  
3915  
3920  
3925  
3930  
3935  
3940  
3945  
3950  
3955  
3960  
3965  
3970  
3975  
3980  
3985  
3990  
3995  
4000  
4005  
4010  
4015  
4020  
4025  
4030  
4035  
4040  
4045  
4050  
4055  
4060  
4065  
4070  
4075  
4080  
4085  
4090  
4095  
4100  
4105  
4110  
4115  
4120  
4125  
4130  
4135  
4140  
4145  
4150  
4155  
4160  
4165  
4170  
4175  
4180  
4185  
4190  
4195  
4200  
4205  
4210  
4215  
4220  
4225  
4230  
4235  
4240  
4245  
4250  
4255  
4260  
4265  
4270  
4275  
4280  
4285  
4290  
4295  
4300  
4305  
4310  
4315  
4320  
4325  
4330  
4335  
4340  
4345  
4350  
4355  
4360  
4365  
4370  
4375  
4380  
4385  
4390  
4395  
4400  
4405  
4410  
4415  
4420  
4425  
4430  
4435  
4440  
4445  
4450  
4455  
4460  
4465  
4470  
4475  
4480  
4485  
4490  
4495  
4500  
4505  
4510  
4515  
4520  
4525  
4530  
4535  
4540  
4545  
4550  
4555  
4560  
4565  
4570  
4575  
4580  
4585  
4590  
4595  
4600  
4605  
4610  
4615  
4620  
4625  
4630  
4635  
4640  
4645  
4650  
4655  
4660  
4665  
4670  
4675  
4680  
4685  
4690  
4695  
4700  
4705  
4710  
4715  
4720  
4725  
4730  
4735  
4740  
4745  
4750  
4755  
4760  
4765  
4770  
4775  
4780  
4785  
4790  
4795  
4800  
4805  
4810  
4815  
4820  
4825  
4830  
4835  
4840  
4845  
4850  
4855  
4860  
4865  
4870  
4875  
4880  
4885  
4890  
4895  
4900  
4905  
4910  
4915  
4920  
4925  
4930  
4935  
4940  
4945  
4950  
4955  
4960  
4965  
4970  
4975  
4980  
4985  
4990  
4995  
5000  
5005  
5010  
5015  
5020  
5025  
5030  
5035  
5040  
5045  
5050  
5055  
5060  
5065  
5070  
5075  
5080  
5085  
5090  
5095  
5100  
5105  
5110  
5115  
5120  
5125  
5130  
5135  
5140  
5145  
5150  
5155  
5160  
5165  
5170  
5175  
5180  
5185  
5190  
5195  
5200  
5205  
5210  
5215  
5220  
5225  
5230  
5235  
5240  
5245  
5250  
5255  
5260  
5265  
5270  
5275  
5280  
5285  
5290  
5295  
5300  
5305  
5310  
5315  
5320  
5325  
5330  
5335  
5340  
5345  
5350  
5355  
5360  
5365  
5370  
5375  
5380  
5385  
5390  
5395  
5400  
5405  
5410  
5415  
5420  
5425  
5430  
5435  
5440  
5445  
5450  
5455  
5460  
5465  
5470  
5475  
5480  
5485  
5490  
5495  
5500  
5505  
5510  
5515  
5520  
5525  
5530  
5535  
5540  
5545  
5550  
5555  
5560  
5565  
5570  
5575  
5580  
5585  
5590  
5595  
5600  
5605  
5610  
5615  
5620  
5625  
5630  
5635  
5640  
5645  
5650  
5655  
5660  
5665  
5670  
5675  
5680  
5685  
5690  
5695  
5700  
5705  
5710  
5715  
5720  
5725  
5730  
5735  
5740  
5745  
5750  
5755  
5760  
5765  
5770  
5775  
5780  
5785  
5790  
5795  
5800  
5805  
5810  
5815  
5820  
5825  
5830  
5835  
5840  
5845  
5850  
5855  
5860  
5865  
5870  
5875  
5880  
5885  
5890  
5895  
5900  
5905  
5910  
5915  
5920  
5925  
5930  
5935  
5940  
5945  
5950  
5955  
5960  
5965  
5970  
5975  
5980  
5985  
5990  
5995  
6000  
6005  
6010  
6015  
6020  
6025  
6030  
6035  
6040  
6045  
6050  
6055  
6060  
6065  
6070  
6075  
6080  
6085  
6090  
6095  
6100  
6105  
6110  
6115  
6120  
6125  
6130  
6135  
6140  
6145  
6150  
6155  
6160  
6165  
6170  
6175  
6180  
6185  
6190  
6195  
6200  
6205  
6210  
6215  
6220  
6225  
6230  
6235  
6240  
6245  
6250  
6255  
6260  
6265  
6270  
6275  
6280  
6285  
6290  
6295  
6300  
6305  
6310  
6315  
6320  
6325  
6330  
6335  
6340  
6345  
6350  
6355  
6360  
6365  
6370  
6375  
6380  
6385  
6390  
6395  
6400  
6405  
6410  
6415  
6420  
6425  
6430  
6435  
6440  
6445  
6450  
6455  
6460  
6465  
6470  
6475  
6480  
6485  
6490  
6495  
6500  
6505  
6510  
6515  
6520  
6525  
6530  
6535  
6540  
6545  
6550  
6555  
6560  
6565  
6570  
6575  
6580  
6585  
6590  
6595  
6600  
6605  
6610  
6615  
6620  
6625  
6630  
6635  
6640  
6645  
6650  
6655  
6660  
6665  
6670  
6675  
6680  
6685  
6690  
6695  
6700  
6705  
6710  
6715  
6720  
6725  
6730  
6735  
6740  
6745  
6750  
6755  
6760  
6765  
6770  
6775  
6780  
6785  
6790  
6795  
6800  
6805  
6810  
6815  
6820  
6825  
6830  
6835  
6840  
6845  
6850  
6855  
6860  
6865  
6870  
6875  
6880  
6885  
6890  
6895  
6900  
6905  
6910  
6915  
6920  
6925  
6930  
6935  
6940  
6945  
6950  
6955  
6960  
6965  
6970  
6975  
6980  
6985  
6990  
6995  
7000  
7005  
7010  
7015  
7020  
7025  
7030  
7035  
7040  
7045  
7050  
7055  
7060  
7065  
7070  
7075  
7080  
7085  
7090  
7095  
7100  
7105  
7110  
7115  
7120  
7125  
7130  
7135  
7140  
7145  
7150  
7155  
7160  
7165  
7170  
7175  
7180  
7185  
7190  
7195  
7200  
7205  
7210  
7215  
7220  
7225  
7230  
7235  
7240  
7245  
7250  
7255  
7260  
7265  
7270  
7275  
7280  
7285  
7290  
7295  
7300  
7305  
7310  
7315  
7320  
7325  
7330  
7335  
7340  
7345  
7350  
7355  
7360  
7365  
7370  
7375  
7380  
7385  
7390  
7395  
7400  
7405  
7410  
7415  
7420  
7425  
7430  
7435  
7440  
7445  
7450  
7455  
7460  
7465  
7470  
7475  
7480  
7485  
7490  
7495  
7500  
7505  
7510  
7515  
7520  
7525  
7530  
7535  
7540  
7545  
7550  
7555  
7560  
7565  
7570  
7575  
7580  
7585  
7590  
7595  
7600  
7605  
7610  
7615  
7620  
7625  
7630  
7635  
7640  
7645  
7650  
7655  
7660  
7665  
7670  
7675  
7680  
7685  
7690  
7695  
7700  
7705  
7710  
7715  
7720  
7725  
7730  
7735  
7740  
7745  
7750  
7755  
7760  
7765  
7770  
7775  
7780  
7785  
7790  
7795  
7800  
7805  
7810  
7815  
7820  
7825  
7830  
7835  
7840  
7845  
7850  
7855  
7860  
7865  
7870  
7875  
7880  
7885  
7890  
7895  
7900  
7905  
7910  
7915  
7920  
7925  
7930  
7935  
7940  
7945  
7950  
7955  
7960  
7965  
7970  
7975  
7980  
7985  
7990  
7995  
8000  
8005  
8010  
8015  
8020  
8025  
8030  
8035  
8040  
8045  
8050  
8055  
8060  
8065  
8070  
8075  
8080  
8085  
8090  
8095  
8100  
8105  
8110  
8115  
8120  
8125  
8130  
8135  
8140  
8145  
8150  
8155  
8160  
8165  
8170  
8175  
8180  
8185  
8190  
8195  
8200  
8205  
8210  
8215  
8220  
8225  
8230  
8235  
8240  
8245  
8250  
8255  
8260  
8265  
8270  
8275  
8280  
8285  
8290  
8295  
8300  
8305  
8310  
8315  
8320  
8325  
8330  
8335  
8340  
8345  
8350  
8355  
8360  
8365  
8370  
8375  
8380  
8385  
8390  
8395  
8400  
8405  
8410  
8415  
8420  
8425  
8430  
8435  
8440  
8445  
8450  
8455  
8460  
8465  
8470  
8475  
8480  
8485  
8490  
8495  
8500  
8505  
8510  
8515  
8520  
8525  
8530  
8535  
8540  
8545  
8550  
8555  
8560  
8565  
8570  
8575  
8580  
8585  
8590  
8595  
8600  
8605  
8610  
8615  
8620  
8625  
8630  
8635  
8640  
8645  
8650  
8655  
8660  
8665  
8670  
8675  
8680  
8685  
8690  
8695  
8700  
8705  
8710  
8715  
8720  
8725  
8730  
8735  
8740  
8745  
8750  
8755  
8760  
8765  
8770  
8775  
8780  
8785  
8790  
8795  
8800  
8805  
8810  
8815  
8820  
8825  
8830  
8835  
8840  
8845  
8850  
8855  
8860  
8865  
8870  
8875  
8880  
8885  
8890  
8895  
8900  
8905  
8910  
8915  
8920  
8925  
8930  
8935  
8940  
8945  
8950  
8955  
8960  
8965  
8970  
8975  
8980  
8985  
8990  
8995  
9000  
9005  
9010  
9015  
9020  
9025  
9030  
9035  
9040  
9045  
9050  
9055  
9060  
9065  
9070  
9075  
9080  
9085  
9090  
9095  
9100  
9105  
9110  
9115  
9120  
9125  
9130  
9135  
9140  
9145  
9150  
9155  
9160  
9165  
9170  
9175  
9180  
9185  
9190  
9195  
9200  
9205  
9210  
9215  
9220  
9225  
9230  
9235  
9240  
9245  
9250  
9255  
9260  
9265  
9270  
9275  
9280  
9285  
9290  
9295  
9300  
9305  
9310  
9315  
9320  
9325  
9330  
9335  
9340  
9345  
9350  
9355  
9360  
9365  
9370  
9375  
9380  
9385  
9390  
9395  
9400  
9405  
9410  
9415  
9420  
9425  
9430  
9435  
9440  
9445  
9450  
9455  
9460  
9465  
9470  
9475  
9480  
9485  
9490  
9495  
9500  
9505  
9510  
9515  
9520  
9525  
9530  
9535  
9540  
9545  
9550  
9555  
9560  
9565  
9570  
9575  
9580  
9585  
9590  
9595  
9600  
9605  
9610  
9615  
9620  
9625  
9630  
9635  
9640  
9645  
9650  
9655  
9660  
9665  
9670  
9675  
9680  
9685  
9690  
9695  
9700  
9705  
9710  
9715  
9720  
9725  
9730  
9735  
9740  
9745  
9750  
9755  
9760  
9765  
9770  
9775  
9780  
9785  
9790  
9795  
9800  
9805  
9810  
9815  
9820  
9825  
9830  
9835  
9840  
9845  
9850  
9855  
9860  
9865  
9870  
9875  
9880  
9885  
9890  
9895  
9900  
9905  
9910  
9915  
9920  
9925  
9930  
9935  
9940  
9945  
9950  
9955  
9960  
9965  
9970  
9975  
9980  
9985  
9990  
9995  
10000  
10005  
10010  
10015  
10020  
10025  
10030  
10035  
10040  
10045  
10050  
10055  
10060  
10065  
10070  
10075  
10080  
10085  
10090  
10095  
10100  
10105  
10110  
10115  
10120  
10125  
10130  
10135  
10140  
10145  
10150  
10155  
10160  
10165  
10170  
10175  
10180  
10185  
10190  
10195  
10200  
10205  
10210  
10215  
10220  
10225  
10230  
10235  
10240  
10245  
10250  
10255  
10260  
10265  
10270  
10275  
10280  
10285  
10290  
10295  
10300

Standard transformation of the benzonitrile obtained in part C to the benzamidine via the ethyl imidate converted the crude benzonitrile to the benzamidine TFA salt after HPLC purification. LRMS (ES+): 406 (M+H), 811 (H+-dimer) HRMS (NH<sub>3</sub>-CI): Calc: 406.199148 Mass: 406.198887; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.37 (s, 1H), 9.40 (s, 2H), 9.02 (s, 2H), 8.23 (s, 1H), 7.91 (s, 1H), 7.78 (d, 1H), 7.68 (m, 2H), 7.43 (d, 2H), 7.38 (d, 2H), 6.95 (s, 1H), 2.87 (s, 6H), 2.29 (s, 3H).

10

#### Examples 144 and 145

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-diethylamino)phenyl)aminocarbonyl]pyrazole (Example 144) and 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-N,N-diethylamino)phenyl)aminocarbonyl]pyrazole (Example 145)

15

Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(N,N-diethylamino)phenyl)aminocarbonyl]pyrazole.

The pyrazole acid chloride was generated by the standard method and coupled to 0.24 g of commercially available N,N-diethyl-1,4-phenylenediamine using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-[(4'-N,N-diethylaminoaniline)aminocarbonyl]pyrazole. LRMS (NH<sub>3</sub>-CI): 374 (M+H), 747 (2M+H); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.16 (s, 1H), 7.90 (s, 1H), 7.81 (m, 1H), 7.71 (m, 1H), 7.60 (t, 1H), 7.37 (d, 2H), 6.88 (s, 1H), 6.59 (d, 2H), 3.26 (m, 4H), 2.25 (s, 3H), 1.02 (t, 6H).

25

Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-diethylamino)phenyl)aminocarbonyl]pyrazole.

30

Standard transformation of the benzonitrile obtained in part B to the benzamidine via the ethyl imidate converted 0.24 g of the crude benzonitrile to 0.256 g of the benzamidine bis-TFA salt after HPLC purification. LRMS (ES+): 391 (M+H) HRMS (NH<sub>3</sub>-CI): Calc: 391.224635 Mass: 391.224109. 0.017 g of the benzamide was also isolated during HPLC purification.

35

LRMS (ESI+): 392 (M+H) HRMS (NH<sub>3</sub>-CI): calc: 392.208650  
mass:392.207700.

Examples 146 and 147

5           1-(3-amidinophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole (Example 146) and 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole (Example 147)

10   Part A. Preparation of 4-N-formylaminonitrobenzene.

          Treated 0.69 g of 4-aminonitrobenzene with acetic formic anhydride in THF at 0°C. Then warmed reaction mixture to 55 °C for 2H. Concentrated mixture under reduced pressure and  
15   placed residue on high vacuum to give the crude product. LRMS (NH<sub>3</sub>-CI): 184 (M+NH<sub>4</sub>).

          Part B. Preparation of 4-(1-tetrazolyl)nitrobenzene.

20           Made a solution of above compound, 2.63 g triphenylphosphine, 1.15 g TMS azide and 1.75 g DEAD reagent in THF. Let stir for 24H. Diluted reaction mixture with water and extracted with methylene chloride. Dried and concentrated organic extracts to give the crude product which  
25   was purified by standard chromatographic technique. LRMS (NH<sub>3</sub>-CI): 209 (M+NH<sub>4</sub>), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.35 (s, 1H), 8.48 (d, 2H), 8.20 (d, 2H).

          Part C. Preparation of 4-(1-tetrazolyl)aniline.

30           Treated 4-(1-tetrazolyl)nitrobenzene with 10% palladium on carbon in methanol and placed under 40psi of hydrogen for 2H. Passed reaction mixture through a 1 inch celite pad and concentrated filtrate to give the crude product. LRMS (NH<sub>3</sub>-  
35   CI): 162 (M+H), 179 (M+NH<sub>4</sub>), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 9.79 (s, 1H), 7.42 (d, 2H), 6.67 (d, 2H).

Part D. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole.

The pyrazole acid chloride was generated in the standard method and coupled to 0.26 g 4-(1-tetrazolyl)aniline using the standard DMAP coupling to give the 1-(3-cyanophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole. This crude material was used directly.

Part E. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole.

Standard transformation of the benzonitrile obtained in part D to the benzamidine via the ethyl imidate converted the crude benzonitrile to 0.014 g of the benzamidine TFA salt after HPLC purification. LRMS (ES<sup>+</sup>): 388 (M+H) HRMS (NH<sub>3</sub>-CI): Calc: 388.163431 Mass: 388.165343 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.79 (s, 1H), 10.01 (s, 1H), 9.40 (bs, 2H), 8.99 (bs, 2H), 7.93 (s, 1H), 7.85 (m, 4H), 7.77 (m, 2H), 7.67 (m, 1H), 7.04 (s, 1H), 2.31 (s, 3H). 0.007 g of the benzamide was also isolated during HPLC purification. LRMS (ESI<sup>+</sup>): 799 (2M+Na) 777 (2M+H) HRMS (NH<sub>3</sub>-CI): calc: 389.147447 mass:389.149952; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.77 (s, 1H), 10.00 (s, 1H), 7.94 (m, 1H), 7.87 (m, 6H), 7.51 (m, 1H), 6.96 (s, 1H), 2.30 (s, 3H).

#### Examples 148, 149, and 150

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-acetylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole, 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole, and 1-(3-amidinophenyl)-3-methyl-5-[(4'-piperizin-1-yl-phenyl)aminocarbonyl]pyrazole

Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole.

The pyrazole acid chloride was generated by the standard method and coupled to 0.23 g of 4-(N-boc-piperizine)aniline (which is readily available from commercially available 1-(4-nitrophenyl)piperazine) using the standard DMAP coupling to give the crude 1-(3-cyanophenyl)-3-methyl-5-((4'-N-tert-butyloxycarbonylpiperizine-1-phenyl)aminocarbonyl)pyrazole. The crude product was purified by standard chromatographic technique. LRMS (NH<sub>3</sub>-CI): 487 (M+H) <sup>1</sup>H NMR (DMSO-d<sub>6</sub>, 300 MHz)  $\delta$ : 10.60 (s, 1H), 7.90 (s, 1H), 7.81 (m, 1H), 7.73 (m, 1H), 7.61 (t, 1H), 7.47 (d, 2H), 6.90 (s, 1H), 6.88 (d, 2H), 3.41 (complex, 4H), 3.01 (complex, 4H), 2.28 (s, 3H), 1.37 (s, 9H).

Part B. Preparation of 1-(3-amidoximephenyl)-3-methyl-5-((4'-N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole.

Treated 0.29 g of 1-(3-cyanophenyl)-3-methyl-5-((4'-N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole with 0.15 g hydroxylamine hydrochloride and 0.11 g of sodium carbonate in ethanol/water. Warmed reaction mixture to reflux temperature for 5H. Worked up reaction mixture with aqueous washings, dried resulting organic, and concentrated in vacuo to give the crude amidoxime.

Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-((4'-N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole and 1-(3-amidinophenyl)-3-methyl-5-((4'-N-acetylpiperazin-1-yl)phenyl)aminocarbonyl]pyrazole.

Treated crude amidoxime with acetic acid and acetic anhydride for 0.5H. Added a catalytic amount of 10% palladium on carbon to reaction mixture and placed on Parr hydrogenator at 50 psi for 4H. Passed through a 1 inch celite pad and concentrated filtrate to give the crude benzamidine. Purified via standard HPLC technique. The N-acetyl compound LRMS

- (ES+): 446 (M+H, 100) HRMS (FAB+): calc.-446.230448 mass-446.231327 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.33 (s, 1H), 9.39 (bs, 2H), 9.04 (bs, 2H), 7.90 (s, 1H), 6.77 (d, 1H), 7.68 (m, 2H), 7.48 (d, 2H), 6.94 (s, 1H), 6.90 (d, 2H), 3.52 (m, 4H), 3.02 (M, 4H), 2.28 (s, 3H), 2.00-(s, 3H). 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-acetylpiperazin-1-yl)phenyl)aminocarbonyl]pyrazole was isolated as a by-product in addition to the N-boc compound LRMS (ES+): 504 (M+H) HRMS (NH<sub>3</sub>-CI): calc-504.272313 mass-504.272536 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.34 (s, 1H), 9.38 (bs, 2H), 9.05 (bs, 2H), 7.90 (m, 1H), 7.77 (m, 1H), 7.67 (m, 2H), 7.47 (d, 2H), 6.94 (s, 1H), 6.90 (d, 2H), 3.42 (m, 4H), 3.00 (m, 4H), 2.29 (s, 3H), 1.37 (s, 9H).
- Part D. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-piperizin-1-yl)phenyl)aminocarbonyl]pyrazole.

- 0.043 g of 1-(3-amidinophenyl)-3-methyl-5-[(4'-N-tert-butyloxycarbonylpiperizin-1-phenyl)aminocarbonyl]pyrazole was treated with TFA at ambient temperature for 3H. Concentrated reaction mixture under reduced pressure to give the crude product. Purified crude material by standard HPLC technique. LRMS (ES+): 404 (M+H) HRMS (NH<sub>3</sub>-CI): calc-404.219884 mass-404.221193 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.36 (s, 1H), 9.39 (bs, 2H), 9.18 (bs, 2H), 7.90 (s, 1H), 7.77 (d, 1H), 7.67 (m, 2H), 7.01 (d, 2H), 6.92 (m, 3H), 3.22 (m, 8H), 2.29 (s, 3H).

#### Example 151

- 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-cyclohexylphenyl)aminocarbonyl]pyrazole

Part A. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-[(4'-cyclohexylphenyl)aminocarbonyl]pyrazole.

- 0.25 g of N-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid was converted to its corresponding acid chloride by standard procedure and reacted with 0.15 g of 4-cyclohexylaniline in the presence of DMAP in methylene

chloride to afford the title compound after workup and purification by standard chromatographic technique. LRMS (ES+): 461 (M+Na+), 899 (Na+-dimer), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.57 (s, 1H), 8.13 (s, 1H), 7.95 (d, 1H), 7.86 (d, 1H), 7.69 (t, 1H), 7.65 (s, 1H), 7.50 (d, 2H), 7.15 (d, 2H), 2.41 (complex, 1H), 1.70 (complex, 5H), 1.25 (complex, 5H).

Part B. Preparation of 1-(3-amidinophenyl)-3-trifluoromethyl-5-((4'-cyclohexylphenyl)aminocarbonyl)pyrazole.

The cyano derivative was converted to the amidino derivative via the amidoxime as previously described. The amidoxime was reduced to the benzamidine by conversion to the corresponding acetate by acetic acid/acetic anhydride and catalytic reduction with 10% palladium on carbon under a hydrogen atmosphere, also previously described. The crude product was purified by standard HPLC technique to give the TFA salt. LRMS (ES+): 456 (M+H) HRMS (NH<sub>3</sub>-CI): calc-456.199783 mass-456.201120 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.62 (s, 1H), 9.40 (s, 2H), 9.16 (s, 2H), 7.99 (s, 1H), 7.88 (m, 2H), 7.72 (t, 1H), 7.69 (s, 1H), 7.50 (d, 2H), 7.14 (d, 2H), 2.41 (complex, 1H), 1.69 (complex, 5H), 1.25 (complex, 5H).

#### Example 152

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholino)-3'-chlorophenyl)aminocarbonyl]pyrazole

Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(N-morpholino)-3'-chlorophenyl)aminocarbonyl]pyrazole.

N-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid was converted to its corresponding acid chloride by standard procedure. 0.30 g of the acid chloride was reacted with 0.26 g of commercially available 2-chloro-4-morpholinoaniline in the presence of DMAP in methylene chloride to afford the product after workup and purification by standard chromatographic technique. LRMS (ES+): 422 (M+H), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.57 (s, 1H), 8.13 (s, 1H), 7.95 (d, 1H), 7.86



(d, 1H), 7.69 (t, 1H), 7.65 (s, 1H), 7.50 (d, 2H), 7.15 (d, 2H), 2.41 (complex, 1H), 1.70 (complex, 5H), 1.25 (complex, 5H).

5 Part B. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholino)-3'-chlorophenyl)]aminocarbonyl]pyrazole.

The cyano derivative was converted amidino derivative via the amidoxime as previously described. The amidoxime was  
10 reduced to the benzamidine by conversion to the corresponding acetate by acetic acid/acetic anhydride and catalytic reduction of the acetate with 10% palladium on carbon under a hydrogen atmosphere, also previously described. The crude product was purified by standard HPLC technique to give the  
15 bis TFA salt. LRMS (ES+): 439 (M+H) HRMS (NH<sub>3</sub>-CI): calc 439.164927 found 439.163814 <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz)δ: 10.54 (s, 1H), 9.38 (s, 2H), 9.06 (s, 2H), 7.89 (s, 1H), 7.78 (m, 2H), 7.67 (m, 2H), 7.51 (dd, 1H), 7.12 (d, 1H), 6.96 (s, 1H), 3.69 (t, 4H), 2.88 (t, 4H), 2.46 (m, 3H).

20

Example 153

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid  
salt

25

Part A. Preparation of Ethyl N-(3-cyanophenyl)glycine.

To a solution of 15.11 g (128 mmol) of 3-aminobenzonitrile in 200 mL of DMF under N<sub>2</sub> was added 23.50 g  
30 (141 mmol) of ethyl bromoacetate and 14.95 g (141 mmol) anhydrous sodium carbonate. The mixture was heated to 70°C for 5 hours and then cooled to room temperature. Water (500 mL) was added and the mixture stirred vigorously until a precipitate formed. The solid was collected, washed with 100  
35 mL water and then dried in vacuo to give 19.97 g (76%) of the desired compound as a yellow-orange solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.26 (t, 1H); 7.03 (d, 1H); 6.81 (d, 1H); 6.79 (s, 1H); 4.53 (br s, 1H); 4.03 (q, 2H); 3.92 (d, 2H); 1.21 (t, 3H).

Part B. Preparation of N-(3-cyanophenyl)glycine.

To a solution of 17.00 g (83.2 mmol) of ethyl N-(3-cyanophenyl)glycine in 100 mL of THF under N<sub>2</sub> was added 3.67 g (87.4 mmol) of lithium hydroxide monohydrate in 20 mL water. After 15 hours, the mixture was acidified with concentrated hydrochloric acid to pH 3 and a precipitate formed. The solid was collected, washed with 100 mL water and then dried in vacuo to give 14.15 g (97%) of the desired compound as a light yellow solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.28 (dt, 1H); 7.05 (dd, 1H); 6.83 (dd, 1H); 6.82 (d, 1H); 4.00 (s, 2H).

Part C. Preparation of N-(3-cyanophenyl)-N-nitrosoglycine.

Sodium nitrite (5.54 g, 80.3 mmol) in 15 mL of water was added to a suspension of N-(3-cyanophenyl)glycine (14.15 g, 80.3 mmol) in 65 mL of water under N<sub>2</sub>. This was allowed to stir at room temperature for 14 hours. The solution was acidified with concentrated hydrochloric acid to pH 3 and a precipitate formed. The solid was collected, washed with 50 mL water and then dried in vacuo to give 16.06 g (98%) of the desired compound as a grey solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 13.22 (br s, 1H); 8.10 (dd, 1H); 7.99 (ddd, 1H); 7.87 (dd, 1H), 7.72 (t, 1H), 4.78 (s, 2H).

Part D. Preparation of 1-(3-cyanophenyl)-4-oxy-1,2,3-oxadiazole.

N-(3-cyanophenyl)-N-nitrosoglycine (6.97 g, 34 mmol) was dissolved in 32 mL of acetic anhydride and heated to 70°C for 5 hours. The reaction mixture was cooled and then poured into 200 mL of ice-water. After stirring for 30 minutes to decompose the excess acetic anhydride, the reaction mixture was filter to provide 5.99 g (94%) of a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.08 (s, 1H), 8.02 (d, J=8.4, 1H), 7.99 (d, J=7.7, 1H), 7.82 (dd, J=8.4, 7.7, 1H), 6.81 (s, 1H).

Part E. Preparation of 1-(3-cyanophenyl)-4-oxy-5-methylthio-1,2,3-oxadiazole.

1-(3-cyanophenyl)-4-oxy-1,2,3-oxadiazole (1.48 g, 7.9 mmol) was dissolved in 30 mL of dry DMSO and cooled to 0°C. Acetyl chloride (1.25 g, 15.9 mmol) was added very slowly via syringe below the surface of the liquid under N<sub>2</sub>. The reaction mixture was allowed to stir at room temperature for 14 hours. The reaction mixture was diluted with 100 mL Et<sub>2</sub>O and washed twice with 25 mL saturated aqueous NaHCO<sub>3</sub>. Then washed three times with 25 mL water to remove the DMSO. The organic extract was dried with MgSO<sub>4</sub> and concentrated in vacuo to give 1.5 g of a red solid which was used without further purification. MS (NH<sub>3</sub>-CI) m/z 234.0 (M+H).

Part F. Preparation of methyl 1-(3-cyanophenyl)-3-methylthio-pyrazole-5-carboxylate.

The crude 1-(3-cyanophenyl)-4-oxy-5-methylthio-1,2,3-oxadiazole (0.95 g, 3.90 mmol) and methyl propiolate (3.28 g, 39.1 mmol) were dissolved in 40 mL of CH<sub>2</sub>Cl<sub>2</sub> and the quartz reaction vessel was purged with N<sub>2</sub>. The reaction mixture was irradiated in a Rayonet RPR-100 photochemical reactor for 14 hours. The crude product was concentrated in vacuo and then chromatographed with 20% EtOAc/hexanes on silica to provide 0.34 g (32%) of a yellow solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.77 (t, J=1.8, 1H); 7.70 (m, 2H); 7.57 (t, J=8.1, 1H); 6.94 (s, 1H); 3.83 (s, 3H); 2.57 (s, 3H).

Part G. Preparation of 1-(3-cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(thiomethyl)pyrazole.

4-Amino-2'-methylsulfonyl-[1,1']biphenyl (65.7 mg, 0.216 mmol) was suspended in 2 mL of CH<sub>2</sub>Cl<sub>2</sub> and 0.51 mL of a 2M solution of trimethylaluminum in heptane was added slowly via syringe. The reaction was stirred for 30 minutes at room temperature and methyl 1-(3-cyanophenyl)-3-methylthio-

pyrazole-5-carboxylate (56.2 mg, 0.206 mmol) was added. The reaction mixture was stirred at room temperature for an additional 14 hours. The aluminum reagent was quenched by careful addition of 1N HCl to pH 2. Then the reaction mixture  
5 extracted with 10 mL of CH<sub>2</sub>Cl<sub>2</sub> three times. The combined organic extracts were washed with water and brine, dried over MgSO<sub>4</sub> and the solvent evaporated. The desired product was obtained (83 mg, 74%) after silica gel chromatography with 30% EtOAc/hexane. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.16 (dd, J=7.7, 1.5, 1H); 7.84  
10 (br s, 1H); 7.84 (t, J=1.8, 1H); 7.76 (m, 1H); 7.70-7.46 (m, 8H); 7.50 (d, J=8.8, 2H); 7.25 (d, J=7.5, 1H); 6.81 (s, 1H); 2.62 (s, 3H).

Part H. Preparation of 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid salt.  
15

1-(3-Cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(thiomethyl)pyrazole (83 mg, 0.15  
20 mmol) was dissolved in 5 mL of methanol and 10 mL of chloroform. The reaction mixture was cooled in an ice bath and HCl gas was bubbled in for 30 minutes to saturate the solution. The mixture was sealed and allowed to stir at room temperature for 14 hours. The solvents were removed in vacuo  
25 and the resulting solid was used in the next step.

The imidate formed above was added to 0.15 g (1.6 mmol) of ammonium carbonate and 10 mL of methanol. The mixture was allowed to stir under N<sub>2</sub> for 14 hours. The solvent was removed at reduced pressure. The crude benzamidine was purified by  
30 HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 64 mg (84%) of the desired salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.66 (s, 1H); 9.41 (br s, 2H); 8.97 (br s, 2H); 7.96 (m, 2H); 7.79-7.66 (m, 7H); 7.63 (d, J=9.0, 2H); 7.56 (t, J=6.6, 1H); 7.33 (d, J=9.0, 2H); 7.27 (m, 1H); 7.19 (s, 1H); 2.55 (s, 3H).  
35 HRMS 507.1268 (M+H).

## Examples 154 and 155

1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfinyl)pyrazole, trifluoroacetic acid salt (Example 154) and 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole, trifluoroacetic acid salt (Example 155)

To a solution of 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid salt (54 mg, 0.11 mmol) in 10 mL methanol was added Oxone® (66 mg, 0.11 mol) and the reaction stirred for 14 hours. The solvent was removed at reduced pressure. The crude sulfoxide was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 22 mg (38%) of the desired salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.84 (s, 1H); 9.43 (br s, 2H); 9.00 (br s, 2H); 8.00 (s, 1H); 7.99 (m, 1H); 7.87 (m, 2H); 7.75 (m, 2H), 7.65 (d, J=9.6, 2H); 7.56 (m, 2H); 7.34 (d, J=8.4, 2H); 7.27 (m, 3H); 2.99 (s, 3H). HRMS 523.1220 (M+H). Another product, the sulfone, (28 mg, 47%), was isolated from the column. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.89 (s, 1H); 9.52 (br s, 2H); 9.09 (br s, 2H); 8.09 (s, 1H); 8.06 (d, J=7.3, 1H); 7.98 (m, 2H); 7.86 (s, 1H), 7.84 (t, J=9.0, 1H), 7.72 (d, J=8.8, 2H); 7.64 (m, 2H); 7.41 (d, J=8.4, 2H); 7.33 (m, 3H); 3.45 (s, 3H). HRMS 539.1175 (M+H).

## Example 156

1-(3-aminocarbonylphenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole

30

The title compound was prepared via the method described previously. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 5.85 (s, 2H); 7.10 to 8.25 (m, 12H). MS (ESI) 424.14 (M+H)<sup>+</sup>.

**Example 157**

**1-(3-aminocarbonylphenyl)-5-([(2'-aminosulfonyl-[1,1']-biphen-4-yl)methyl]tetrazole**

The title compound was prepared via the method described previously. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 5.85 (s, 2H); 7.15 to 8.25 (m, 12H). MS (ESI) 435.12 (M+H)<sup>+</sup>.

**Example 158**

**1-(3-amidinophenyl)-5-[(4'-cyclopentyloxyphenyl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt**

Part A: Standard coupling protocol of 4-cyclopentyloxy-aniline (obtained by the displacement of 4-fluoronitrobenzene with the anion of cyclopentanol, followed by catalytic (10% Pd/C) reduction in methanol) with the acid chloride derived for N1-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid afforded the amide precursor as a pale yellow oil; <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.79 (bs, 1H), 7.75-7.50 (m, 7H), 6.95 (d, 1H), 6.85 (m, 1H), 4.75 (m, 1H), 1.95-1.70 (m, 6H), 1.60 (bm, 2H), 2.30 (m, 3H) ppm; ESI mass spectrum m/z (rel intensity) 387 (M+H, 100).

Part B: The title compound was obtained as colorless crystals after purification (via standard techniques) following the standard Pinner/amidine reaction sequence. <sup>1</sup>HNMR(DMSO, d<sub>6</sub>) δ: 10.39 (s, 1H), 9.42 (bs, 2H), 9.05 (bs, 2H), 7.94 (s, 1H), 7.82-7.68 (cp, 3H), 7.71 (d, 2H), 6.97 (s, 1H), 6.88 (d, 2H), 4.77 (m, 1H), 2.33 (s, 3H), 1.84-1.59 (cp, 8H) ppm; ESI mass spectrum m/z (rel intensity) 404.2 (M+H, 100).

**Example 159**

**1-(3-amidinophenyl)-5-[(3-((pyrid-2-yl)methylamino)phenyl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt**

Part A: Standard coupling of 3-((pyrid-2-yl)methylamino)aniline [obtained in a two step sequence (condensation and reduction) from 3-nitroaniline and 2-pyridylcarboxaldehyde afforded the desired bis aniline

derivative;  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 8.58 (d,  $J = 5.13$ , 1H); 7.67 (t,  $J = 7.69$ , 1H); 7.35 (d,  $J = 7.69$ , 1H); 7.19 (m, 1H); 6.99 (t,  $J = 7.69$ , 8.06, 1H); 6.14 (m, 2H); 6.01 (m, 1H); 4.66 (brd, 1H); 4.44 (s, 2H); 3.56 (brd, 2H) ppm; Mass spectrum analysis (NH<sub>3</sub>-CI) 200 (M+H, 100)].

with the acid chloride derived from 1-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid afforded the coupled benzonitrile precursor which was then subjected to the standard Pinner amidine reaction sequence to afford the desired benzamidine compound as colorless crystals;  $^1\text{H NMR}(\text{DMSO})$   $\delta$ : 10.28 (s, 1H); 9.42 (s, 2H); 9.08 (s, 2H); 8.58 (d,  $J = 4.39$ , 1H); 7.83 (m, 3H); 7.72 (m, 2H); 7.46 (d,  $J = 8.06$ , 1H); 7.40 (t,  $J = 5.49, 6.59$ , 1H); 7.01 (m, 3H); 6.88 (d,  $J = 8.05$ , 1H); 6.34 (d,  $J = 8.06$ , 1H); 4.39 (s, 2H); 2.31 (s, 3H) ppm; ESI mass spectrum analysis  $m/z$  (rel intensity) 426.1 (M+H, 100); HRMS for C<sub>24</sub>H<sub>24</sub>N<sub>7</sub>O 426.204234 (calcd.), 426.201998 (obs).

#### Example 160

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]pyrazole

Part A. Preparation of N-(4-nitrophenyl)imidazole.

4-Imidazolo-nitrobenzene (5g) was hydrogenated (10% Pd/C) in 200mL methanol for 20h. the reaction mixture was filtered through a celite pad and evaporated the solvent to afford 3.99g of the crude product which was used directly in the next step. Mass spectrum analysis (H<sub>2</sub>O-GC/MS):160 (M+H, 100).

Part B. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]pyrazole.

The product from part A was then coupled to 1-(3-cyanophenyl)-3-methylpyrazole-5-carboxylic acid via the acid chloride methodology described previously to afford the desired amide which was then purified via standard reverse phase HPLC techniques to afford the desired material.

<sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.73 (s,1H) 9.35 (bs,1H) 8.13 (s,1H) 7.95 (s,1H) 7.90-7.60 (complex,8H) 7.0 (s,1H) 2.30 (s,3H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 369 (m+H, 100); HRMS calc. mass 369.146384; found 369.145884.

5

Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]pyrazole.

The product from part B was then subjected to the standard Pinner amidine reaction sequence to afford the desired benzamidine after HPLC purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.65 (s,1H) 9.40 (bs,2H) 9.00 (bs,2H) 8.19 (s,1H) 7.90 (s,1H) 7.80-7.55 (complex,8H) 7.06 (s,1H) 7.00 (s,1H) 2.30 (s,3H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 386 (M+H, 100). HRMS (FAB), calc. mass 386.172933; found 386.173388.

15

#### Example 161

20

1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-(N-morpholino)-3-chlorophenyl)aminocarbonyl]pyrazole

Part A. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-[(4'-(N-morpholino)-3-chlorophenylaminocarbonyl]pyrazole.

Standard coupling of commercially available 2-chloro-4-morpholinoaniline with N-(3-cyanophenyl)-3-trifluoromethylpyrazole-5-carboxylic acid via its acid chloride under usual conditions afforded the desired coupled product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.66 (s,1H), 8.12 (s,1H), 7.97 (d,1H), 7.87 (d,1H), 7.70 (complex,3H), 7.50 (dd,1H), 7.14 (d,2H), 3.70 (m,4H), 2.90 (m,4H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 476 (M+H, 100).

30

Part B: Preparation of 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-N-morpholino)-3-chlorophenyl)aminocarbonyl]pyrazole.

35

The cyano compound from part A was converted to the amidino derivative via the amidoxime as previously described.



The amidoxime was reduced to the title compound (acetic acid/acetic anhydride and catalytic reduction of the acetate with 10% palladium on carbon under a hydrogen atmosphere) as previously described. The crude product was purified by standard HPLC technique to afford the desired compound as its bis TFA salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.73 (s,1H) 9.41 (bs,2H) 9.09 (bs,2H) 7.98 (s,1H) 7.89 (m,2H) 7.73 (complex,3H) 7.50 (d,1H) 7.14 (d,1H) 3.69 (complex,4H) 2.89 (complex,4H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 493 (M+H, 100); HRMS(FAB+): calc-493.136662, obs. 493.136951.

#### Example 162

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-pyrrolidinocarbonyl)-3'-chlorophenyl)aminocarbonyl]pyrazole

Part A: Preparation of 4'-(pyrrolidinocarbonyl)-3-chloronitrobenzene.

To a dichloromethane solution of 4-nitro-3-chlorobenzoic acid (1.61g) was added N-methylmorpholine (1.93mL) and isobutylchloroformate (1.04mL) followed by the addition of pyrrolidine (0.67mL) and the reaction mixture was warmed to ambient temperature. Concentration of the reaction mixture followed by aqueous workup and extraction with ethylacetate afforded crude product which was used directly into the next reaction. LRMS(NH<sub>3</sub>-CI): 255 (m+H).

Part B. Preparation of 4'-(pyrrolidinocarbonyl)-3-chloroaniline.

The crude 4'-(pyrrolidinocarbonyl)-3-chloronitrobenzene was treated with a catalytic amount 10% palladium on carbon in 20mL methanol and placed under 10psi hydrogen for 15h. Passed through a 1" Celite pad and concentrated filtrate. The residue was washed with ethyl acetate and 3x20mL portions 1.0M HCl, dried (magnesium sulfate) and concentrated in vacuo. Recrystallized from methylene chloride/methanol to afford 1.80g of crystalline 4'-carboxamidopyrrolindino-3-

chloroaniline. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 6.94 (d, 1H, J=8.42), 6.55 (d, 1H, J=1.83), 6.47 (dd, 1H, J=8.43, J=7.69), 3.36 (t, 2H, J=6.23, J=6.95), 3.09 (t, 2H, J=6.22, J=6.23), 1.78 (m, 4H) ppm; Mass spectrum analysis (NH<sub>3</sub>-CI): 225 (m+H, 100).

5

Part C. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[(4'-pyrrolidinocarbonyl)-3-chlorophenyl]aminocarbonylpyrazole.

Standard coupling of the product from part B with the acid chloride derived from 1-(3-cyanophenyl)-3-methyl-pyrazole 5-carboxylic acid chloride afforded the desired coupled product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.71 (s, 1H), 7.97 (d, 1H), 7.84 (m, 2H), 7.76 (m, 1H), 7.63 (m, 2H), 7.32 (d, 1H), 7.00 (s, 1H), 3.42 (t, 2H), 3.06 (t, 2H), 2.29 (s, 3H), 1.80 (m, 4H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 434 (M+Na, 100).

15

Part D. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-pyrrolidinocarbonyl)-3-chlorophenyl]aminocarbonylpyrazole.

20

The benzonitrile product from part C was then converted to the desired benzamidine via standard conditions described previously. Purification via reverse phase HPLC afforded the title compound as its trifluoro-acetate salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.73 (s, 1H), 9.38 (s, 2H), 9.04 (s, 2H), 7.91 (s, 1H), 7.85 (s, 1H), 7.79 (d, 1H), 7.74 (d, 1H), 7.67 (d, 1H), 7.62 (m, 1H), 7.02 (s, 1H), 3.41 (t, 2H), 3.06 (t, 2H), 2.30 (s, 3H), 1.82 (m, 4H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 451 (M+H, 100). HRMS(CI): obs. 451.164788 calc. 451.164927.

25

30

#### Example 163

1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholinocarbonyl)-3-chlorophenyl)aminocarbonyl]pyrazole

35

Part A. Preparation of 4-(N-morpholinocarbonyl)-3-chloronitrobenzene.

To a dichloromethane solution of 4-nitrobenzoyl chloride (2.41g) was added morpholine (3.40mL) in 75mL methylene chloride at 0°C. The reaction mixture was warmed to ambient temperature over 20h, then diluted with water (100mL). The organic layer was separated, washed with water (50mL), 1.0M HCl (50mL), dried (magnesium sulfate) and concentrated in vacuo. The crude material was used directly into the next step without further purification. Mass spectrum analysis (NH<sub>3</sub>-CI): 237 (m+H, 100). The product obtained above was then subjected to catalytic reduction (10% palladium on carbon in 60mL methanol and placed under 60psi hydrogen for 3h), filtered through a celite pad and evaporated to afford the desired aniline derivative. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 7.09 (d,2H), 6.50 (d,2H), 3.54 (t,4H), 3.44 (t,4H), 3.29 (s,2H) ppm; Mass spectrum analysis (NH<sub>3</sub>-CI): 207 (m+H, 100).

Part B. Preparation of 1-(3-cyanophenyl)-3-methyl-5-[4'-(N-morpholinocarbonyl)-3-chlorophenyl]aminocarbonyl]pyrazole.

Standard coupling of the product from part A with the acid chloride derived from N-(3-cyanophenyl)-3-methylpyrazole-5-carboxylic acid followed by usual workup afforded the desired product after silica gel column chromatography (oil); <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.63 (s,1H), 7.94 (s,1H), 7.83 (d,1H,J=7.69), 7.75 (dd,1H,J=8.06,J=8.06), 7.70 (d,2H,J=8.42), 7.63 (t,1H,J=7.69,J=8.05), 7.37 (d,2H,J=8.06), 6.98 (s,1H), 3.28 (d,8H,J=6.96), 2.28 (s,3H); ESI mass spectrum analysis m/z (rel. intensity) 438 (M+Na), 416 (M+H, 100).

Part C. Preparation of 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholinocarbonyl)phenyl)aminocarbonyl]pyrazole.

Standard conversion of the product from part B to the benzamidine afforded after purification via reverse phase HPLC the desired product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ: 10.66 (s,1H), 9.38 (bs,2H), 9.04 (bs,2H), 7.90 (d,1H,J=9.52), 7.78 (d,1H,J=7.33), 7.73-7.62 (complex,4H), 7.37 (d,2H,J=8.42), 7.00 (s,1H), 3.55-3.46 (complex,8H), 2.30 (s,3H). ESI mass

spectrum analysis m/z (rel. intensity) 433 (M+H, 100); HRMS obs. 433.199045; calc.433.198814.

#### Example 164

5     **1-(3-Cyanophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid**

1-(3-Cyanophenyl)-3-trifluoromethyl-pyrazol-5-yl  
carboxylic acid (0.5g, 1.8mmol) was coupled with 4-imidazol  
10 aniline (0.3g,1.8mmol) by standard conditions and purified by HPLC to afford 0.67g(71%) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.99 (s,1H), 9.55 (s,1H), 8.22 (d,j=5.49Hz,2H), 8.04 (d,j=7.69Hz,1H), 7.96 (d,j=8.06Hz,1H), 7.89 (s+d,j=8.79Hz,3H), 7.80 (m,4H) ppm; HRMS 423.118119 (calc'd), 423.116015 (obs.);  
15 Analysis calc'd for C<sub>21</sub>H<sub>13</sub>F<sub>3</sub>N<sub>6</sub>O(TFA): C:51.50,H:2.63,N:15.67, found C:51.52,H:2.71,N:15.49.

#### Example 165

20     **1-(3-amidinophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid**

1-(3-Cyanophenyl)-5-[(4'-imidazol-1-ylphenyl)aminocarbonyl]-3-trifluoromethylpyrazole was subjected to  
25 standard Pinner amidine reaction sequence and purified under standard conditions to afford title amidine (79%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.02 (s,1H), 9.46 (s,1.5H),9.42 (s,1H), 9.22 (s,1.5H), 8.17 (s,1H), 8.06 (s,1H), 7.97 (t,j=7.69Hz,2H), 7.88 (d,j=8.79Hz,2H), 7.80 (m,3H), 7.79 (d,j=9.0Hz,2H) ppm; HRMS  
30 440.144668 (calc'd), 440.144557 (obs.); Analysis calc'd for C<sub>21</sub>H<sub>16</sub>F<sub>3</sub>N<sub>7</sub>O(TFA)<sub>2</sub>·(H<sub>2</sub>O)<sub>1</sub>: C:43.81,H:2.94,N:14.30,found C:43.76,H:2.70,N:13.95.

#### Example 166

35     **1-(3-amidinophenyl)-5-[(4'-(N-methyltetrazolon-1-yl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid**

Part A. 4-Nitrobenzoic acid was converted to the 4-nitrophenyltetrazolone according to the procedure of Toselli, M. and Zaneratio, P., *J.C.S. Perk. Trans.* **1992**, *1*, 1101.

<sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.46 (d, j=9.15Hz, 2H), 8.22 (d, j=9.16Hz, 2H).

5

Part B. To 4-nitrophenyltetrazolone (0.8g, 3.9mmol) in DMF (10mL) at 0°C was added iodomethane (0.38mL) and 60% sodium hydride (0.23g). The reaction was allowed to warm to ambient temperature and stirred 24h. The reaction was quenched with water and extracted with ethyl acetate and dried (MgSO<sub>4</sub>). The crude product was purified by chromatography on silica gel and recrystallized from methylene chloride/hexanes to afford 0.35g (41%) product, MS (DCI) m/z 192 (M+H-NO)<sup>+</sup>, 209 (M+NH<sub>4</sub>-NO)<sup>+</sup>.

10

Part C. The nitro compound (0.215g, 0.97mmol) from part B was hydrogenated under 1 atmosphere of hydrogen in the presence of a catalytic amount of 10% palladium on carbon to the aniline, Mass spectrum analysis (DCI) m/z 192 (M+H)<sup>+</sup>, 209 (M+NH<sub>4</sub>)<sup>+</sup>.

15

Part D. 1-(3-Cyanophenyl)-3-trifluoromethyl-pyrazol-5-yl carboxylic acid (0.38g, 1.4mmol) was coupled with the aniline from part C by standard procedure to afford the nitrile in 43% yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.04 (s, 1H), 7.95 (d, j=9.16Hz, 2H), 7.85 (s, 1H), 7.79 (m, 2H), 7.67 (m, 3H), 7.21 (s, 1H), 3.71 (s, 3H) ppm; MS (ESI) m/z= 454.9 (M+H)<sup>+</sup>, 477 (M+Na)<sup>+</sup>.

25

Part E. The nitrile from part D was subjected to the standard Pinner conditions to afford the title amidine in 53% yield. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.93 (s, 1H), 9.46 (s, 1.5H), 9.12 (s, 1.5H), 8.04 (s, 1H), 7.95 (d, j=7.69Hz, 2H), 7.84 (s, 4H), 7.81 (m, 2H), 3.61 (s, 3H) ppm; HRMS 472.145731 (calc'd), 472.145205 (obs.); Analysis calcd for C<sub>20</sub>H<sub>16</sub>F<sub>3</sub>N<sub>9</sub>O<sub>2</sub> (TFA) 1.2: C:44.23, H:2.85, N:20.73, found C:44.40, H:2.85, N:20.15.

30

35

#### Example 167

1-(3'-Aminocarbonylphenyl)-5-[(2'-aminosulfonylphenyl-[1,1']-biphen-4-yl)methylcarbonyl]-3-methyl-pyrazole

The title amide was isolated from the Pinner reaction via HPLC separation protocols. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.63 (s, 1H), 8.06 (s, 1H), 8.03 (dd, j=2.19, 7.32Hz, 1H), 7.87 (s, 1H), 7.61 (m, 2H), 7.53 (m+d, j=7.33Hz, 3H), 7.44-7.26 (m, 6H), 7.21 (s, 2H), 4.33 (s, 2H), 2.33 (s, 3H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 497 (M+Na, 100) 433 (M+H).

#### Example 168

1-((3-aminophenyl)-5-[(4'-(pyrrolidinomethyl)phenyl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt

Standard coupling of 4-(pyrrolidinomethyl)aniline with the acid chloride derived from 1-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid afforded the coupled benzonitrile precursor which was then subjected to the standard Pinner amidine reaction sequence to afford after purification the title compound as colorless crystals; <sup>1</sup>HNMR(DMSO) δ: 10.69 (s, 1H); 9.42 (s, 2H); 9.20 (s, 2H); 7.96 (s, 1H); 7.84 (m, 1H); 7.75-7.68 (m, 4H); 7.48 (d, 2H, J=8.79); 7.04 (s, 1H); 4.31 (m, 2H); 3.35 (brd, 2H); 3.05 (brd, 2H); 2.34 (s, 3H); 2.05 (brd, 2H); 1.85 (brd, 2H) ppm; ESI mass spectrum m/z (rel. intensity) 403 (M+H, 100); HRMS found for C<sub>23</sub>H<sub>27</sub>N<sub>6</sub>O 403.224635 (calcd), 403.222719 (obs).

25

#### Example 169

1-((3-aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

Part A: To commercially available 3-nitrophenylhydrazine hydrochloride (1.00 g, 5.27 mmol) in 15 mL of absolute ethanol was added 1,1,1-trichloro-4-methoxy-3-penten-2-one (1.15 g, 5.27 mmol) and the reaction brought to reflux for 12 h. The solvent was evaporated and the residue subjected to silica gel flash chromatography eluting with 20% ethyl acetate in hexanes. The first fraction to elute was the desired ethyl ((3-nitrophenyl)-3-methyl-5-pyrazole carboxylate. MS (ES+) 276.1 (M+H)<sup>+</sup> (100%). The ester (110 mg, 0.400 mmol) was coupled with (2'-tert-butylaminosulfonyl-[1,1']-biphen-4-

yl)amine (122 mg, 0.400 mmol) using Weinreb's trimethylaluminium procedure. After preparative TLC (eluent 50% ethyl acetate/hexanes) 178.2 mg (83% yield) of 1-(3-nitrophenyl)-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole was isolated as a colorless solid. MS (ES+) 551.24 (M+NH<sub>4</sub>)<sup>+</sup> (30%); 556.18 (M+Na)<sup>+</sup> (100%).

Part B: The product from part 170.5 mg (0.320 mmol) was refluxed in 5 mL of trifluoroacetic acid for 12 h. Preparative TLC (eluent 10% methanol/chloroform) afforded 1-(3-nitrophenyl)-3-methyl-5-[(2'-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole as a colorless solid. MS (ES+) 478.23 (M+H)<sup>+</sup> (30%); 500.21 (M+Na)<sup>+</sup> (100%). HRMS (FAB+) (M+H)<sup>+</sup>: calc. 478.118516; found 478.117673.

Part C: The product from part B 64.3 mg (0.135 mmol) was subjected to catalytic hydrogenation (5% Pd/C in ethanol under 1 atm of hydrogen) to afford the title compound as a colorless solid. <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 8.08 (d, J=7.7 Hz, 1H), 7.61-7.30 (m, 8H), 7.13 (t, J=7.7 Hz, 1H), 6.72 (m, 3H), 2.33 (s, 3H). MS (ESI+): 448.11 (M+H)<sup>+</sup> (35%); 470.16 (M+Na)<sup>+</sup> (100%). HRMS (FAB+) (M+H)<sup>+</sup>: calc. 448.144337; found 448.144965.

#### Example 170

1-(2'-Aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

The title compound was made in a similar manner to Example 169. <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 8.14-8.03 (m, 2H), 7.58-6.74 (m, 11H), 2.47 (s, 3H). MS (ES+) 448.12 (M+H)<sup>+</sup> (60%); 470.16 (M+Na)<sup>+</sup> (100%).

#### Example 171

1-(3-amino-4'-chlorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

The title compound was made in a similar manner to Example 169. <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 8.08 (d, J=6.9 Hz, 1H), 8.07-7.23

(m, 8H), 6.91 (d,  $J=2.2$  Hz, 1H), 6.75 (s, 1H), 6.66 (dd,  $J=8.43, 2.56$  Hz, 1H), 2.33 (s, 3H). MS (ES+) 482.0 (M+H)<sup>+</sup> (80%); 484.0 (30%); 504.0 (M+Na)<sup>+</sup> (100%); 506.0 (40%).

5

**Example 172**

**1-(3-amino-4'-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole**

The title compound was made in a similar manner to  
10 Example 169. <sup>1</sup>HNMR(CD<sub>3</sub>OD)  $\delta$ : 8.14-8.03 (m, 2H), 7.58-6.74 (m, 11H), 2.47 (s, 3H). MS (ES+) 466.0 (M+H)<sup>+</sup> (5%); 488.0 (M+Na)<sup>+</sup> (100%).

**Example 173**

15 **1-(3-amino-4'-methoxyphenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole**

The title compound was made in a similar manner to  
Example 169. <sup>1</sup>HNMR(CD<sub>3</sub>OD)  $\delta$ : 8.10 (d,  $J=6.6$  Hz, 1H), 7.63-7.31  
20 (m, 7H), 6.89-6.72 (m, 4H), 3.88 (s, 3H), 2.34 (s, 3H). MS (ES+) 478.1 (M+H)<sup>+</sup> (25%); 500.0 (M+Na)<sup>+</sup> (100%).

**Example 174**

25 **1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt**

Part A. Preparation of 1-(3-nitro-4-chlorophenyl)-5-carboethoxytetrazole.

30 4-Chloro-3-nitroaniline (10.36 g, 60 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (100 mL). Triethylamine (10 mL, 70 mmol) was added followed by ethyl oxalyl chloride (6.8 mL, 60 mmol). The mixture was stirred at room temperature under N<sub>2</sub> for 15 min. It was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water and brine.  
35 The CH<sub>2</sub>Cl<sub>2</sub> solution was dried over MgSO<sub>4</sub> and concentrated to a light yellow solid (15.53 g).

The above amide (5.5 g, 20.2 mmol) was refluxed 4 h with a solution of triphenylphosphine (7.87 g, 30 mmol) in 100 mL



of CCl<sub>4</sub> (The solution was stirred at 0°C for 15 min before the amide was added). The reaction mixture was cooled and the precipitate was filtered off. The filtrate was concentrated to a solid. It was then dissolved in 100 mL of CH<sub>3</sub>CN and NaN<sub>3</sub> (1.31 g, 1eq) was added. The mixture was stirred at room temperature under N<sub>2</sub> for 12 h. The solvent was removed. The solid was dissolved in EtOAc and washed with water and brine. It was dried over MgSO<sub>4</sub>, concentrated, and chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>) to afford 3.19 g of the desired product. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 1.35 (t, 3H); 4.42 (q, 2H); 7.50-7.70 (m, 2H); 8.10 (s, 1H). MS (DCI-NH<sub>3</sub>) 315 (M+NH<sub>4</sub>)<sup>+</sup>.

Part B. Preparation of 1-(3-nitro-4-chlorophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole.

2'-t-Butylaminosulfonyl-4-amino-[1,1']-biphenyl (1.33 g, 4.37 mmol) was dissolved in 40 mL of anhydrous CH<sub>2</sub>Cl<sub>2</sub>, and trimethylaluminum (11 mL of 2.M solution in heptane) was added slowly. The mixture was stirred at room temperature under N<sub>2</sub> for 15 min. Then, a solution of material from part A (1.30 g, 4.37 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added. The mixture was stirred at room temperature under N<sub>2</sub> for 18 h. The reaction mixture was quenched carefully with 1N HCl. It was diluted with CH<sub>2</sub>Cl<sub>2</sub> and washed with water and brine. The organic solution was then dried over MgSO<sub>4</sub>, concentrated, and chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>) to give 1.5 g of the desired product. MS(ESI) 554.1 (M-H)<sup>+</sup>.

Part C. Preparation of 1-(3-nitro-4-chlorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole.

The material from Part B (1.5 g, 2.7 mmol), and trifluoroacetic acid (20 mL) was stirred at room temperature under N<sub>2</sub> overnight. The trifluoroacetic acid was removed and chromatographed on silica gel (10% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>) to afford 0.72 g of desired product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 7.25 to 8.20 (m, 11H); 8.69 (s, 1H); 11.55 (s, 1H). MS (ESI) 497.9:499.9 (3:1) (M-H)<sup>+</sup>.

Part D. Preparation of 1-(3-amino-4-chlorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt.

5

The material from part C (0.72 g, 1.44 mmol) was dissolved in EtOAc (30 mL). SnCl<sub>2</sub> 2H<sub>2</sub>O (2.59 g, 11.52 mmol) was added. The reaction mixture was brought to reflux for 1 h and then cooled it to the room temperature. Saturated NaHCO<sub>3</sub> was added to the mixture until the pH 8.0. The mixture was partitioned between EtOAc and NaHCO<sub>3</sub> layer. The EtOAc layer was washed with water and brine. It was dried over MgSO<sub>4</sub> and concentrated. The solid was dissolved in CH<sub>3</sub>CN/TFA and purified by reversed phase HPLC to give 300 mg of the desired product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 6.80 to 8.00 (m, 11H); 11.40 (s, 1H). MS (DCI-NH<sub>3</sub>) 470.0 (M+H)<sup>+</sup>.

10

15

#### Example 175

1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole, trifluoroacetic acid salt

20

The title compound was prepared via the method of Example 171. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 6.80 to 8.40 (m, 10H); 11.70 (s, 1H). MS (ESI) 471.20 (M+H)<sup>+</sup>.

25

#### Example 176

1-(3-amino-4'-methoxyphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

30

The title compound was prepared via the method of Example 174. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 6.80 to 8.05 (m, 11H); 11.15 (s, 1H). MS (ESI) 466.0 (M+H)<sup>+</sup>.

35

#### Example 177

1-(3-aminomethylphenyl)-5-[(2'-aminosulfonylphenyl)pyrid-2-yl]aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid

- Part A: Ethyl-1-(3-cyanophenyl)-3-methyl-5-pyrazole-carboxylate (2.7g, 10.58mmol) was dissolved in methanol (50mL). To this solution was added glacial acetic acid (2mL) and 10% palladium on carbon (cat.). The reaction mixture was hydrogenated (50psi) for 12h, filtered over celite and evaporated to the crude benzylamine salt. Without further purification the crude amine was converted to the carbobenzyloxy derivative by treatment with CBzCl in saturated sodium bicarbonate solution. The organics were extracted with ethyl acetate (2x100mL) dried over magnesium sulfate and evaporated to the crude product (2.15g obtained). The oil was then hydrolysed with LiOH (0.22g, 5.5mmol) in aqueous THF for 16h. The reaction mixture was quenched with water (500mL) and unreacted products were extracted with ethyl acetate (2x100mL). The aqueous layer was carefully acidified (1NHCl) followed by extraction with ethyl acetate (2X100mL) dried (magnesium sulfate) and evaporated to pure acid (1.23g); ESI(-ve) 362 (M-H, 100).
- Part B: Standard coupling (TBTU, triethylamine in anhydrous THF) of the product from part A with 2-amino-5-(2'-tert-butylaminosulfonylphenyl)pyridine afforded the desired amide derivative which was dehydrogenated (10% Pd/C, methanol, balloon) overnight. The reaction mixture was filtered over celite and evaporated to a pale yellow oil. The desired product was obtained as colorless crystals after purification via standard reverse phase techniques; <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.35 (d, 1H), 8.19 (bs, 1H), 8.00 (t, 1H), 7.78 (dd, 1H), 7.63 (t, 2H), 7.77-7.37 (m, 6H), 7.06 (s, 1H), 4.13 (m, 2H), 2.30 (s, 3H) ppm; ESI mass spectrum analysis m/z (rel intensity) 463.3 (M+H, 100).

#### Example 178

- 1-(3-aminomethyl-4'-methylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4yl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt

- Part A: Ethyl 1-(3-cyano-4-methylphenyl)-3-methyl-5-pyrazole-carboxylate was prepared as colorless crystals following the standard condensation (3-cyano-4-methylphenyl-hydrazine and ethyl 2-(N-(methoxy)imino)-4-oxopentanoate in acetic acid) reaction protocol discussed previously. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.68 (s, 1H), 7.57 (dd, 1H), 7.58 (d, 1H), 6.82 (s, 1H), 4.24 (q, 2H), 2.40 (s, 3H), 2.37 (s, 3H), 1.27 (t, 3H) ppm; ESI mass spectrum analysis (270 (M+H, 100)).
- Part B: Standard Weinreb coupling protocol of the product from part A with 1-amino-2'-tert-butylaminosulfonyl-biphenyl afforded the desired coupled product. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.30 (bs, 1H), 8.13 (bd, 1H), 7.78-7.23 (m, 10H), 6.78 (s, 1H), 3.68 (s, 1H), 2.60 (s, 3H), 2.40 (s, 3H), 1.01 (s, 9H) ppm; ESI mass spectrum analysis ESI mass spectrum m/z (rel intensity) 550 (M+Na, 100).
- Part C: The product from part B was then hydrogenated at 50psi in acidic methanol as previously described, then treatment with TFA (neat) and purified via standard reverse phase chromatography to afford the title compound as colorless crystals. <sup>1</sup>HNMR(DMSO, d<sub>6</sub>) δ: 10.6 (s, 1H), 8.14 (bs, 2H), 8.01 (d, 1H), 7.68 (d, 2H), 7.54 (m, 2H), 7.26 (m, 5H), 6.91 (s, 1H), 4.07 (bd, 2H), 2.38 (s, 3H), 2.33 (s, 3H) ppm; ESI mass spectrum m/z (rel intensity) 476 (M+H, 100).

#### Example 179

1-(3-aminomethyl-4'-fluorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt

The title benzylamine was obtained from 3-cyano-4-fluorophenyl hydrazine via methods described previously. <sup>1</sup>HNMR(DMSO, d<sub>6</sub>) δ: 8.25 (bs, 3H), 8.00 (d, 1H), 7.78-7.23 (cp, 12H), 6.95 (s, 1H), 4.14 (m, 2H), 2.30 (s, 3H) ppm; ESI mass spectrum m/z (rel intensity) 480 (M+H, 100).

## Example 180

1-(3-aminomethylphenyl)-5-[(4'-(N-pyrrolidino-  
carbonyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole,  
trifluoroacetic acid

5

Part A. Preparation of 1-(3-cyanophenyl)-5-[(4'-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole.

10

1-(3-Cyanophenyl)-3-trifluoromethyl-pyrazol-5-yl  
carboxylic acid (0.5g, 1.8mmol) was coupled with 4-(N-pyrrolidinocarbonyl)aniline (0.3g, 1.8mmol) by standard conditions to afford 0.4g (56%) of a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 9.72 (s, 1H), 7.78-7.72 (m, 4H), 7.61 (t, j=7.69Hz, 1H), 7.23 (s, 4H), 3.67 (t, j=6.59Hz, 2H), 3.43 (t, j=6.59Hz, 2H), 1.98 (q, j=6.23Hz, 2H), 1.89 (q, j=6.23Hz, 2H) ppm; ESI mass spectrum analysis m/z (rel. intensity) 476 (M+Na, 100), 454.1 (M+H).

15

Part B. The nitrile from part A (0.4g, 0.88mmol), 10% palladium on carbon (50mg) and ethanol (20mL) was placed in a Parr apparatus and hydrogenated 18h at 40 psi. The reaction was filtered and concentrated. The crude product was purified by reverse phase HPLC and freeze-dried to afford 0.38g (76%) of the title amine. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.91 (s, 1H), 8.23 (brd s, 2H), 7.73 (m, 3H), 7.71 (d, j=8.79Hz, 2H), 7.59 (m, 2H), 7.54 (d, j=8.42Hz, 2H), 4.16 (d, j=5.50Hz, 2H), 3.45 (q, j=7.32Hz, 4H), 1.83 (brd m, 4H) ppm; Analysis calc'd for C<sub>23</sub>H<sub>22</sub>F<sub>3</sub>N<sub>5</sub>O<sub>2</sub> (TFA)1 (H<sub>2</sub>O)0.5: C:51.73, H:4.17, N:12.06, found C:51.45, H:3.95, N:11.73.

25

30

## Example 181

1-(3-Ethylcarboxyamidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methyl pyrazole

To 1-(3-cyanophenyl)-5-[(2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole (88mg, 0.15mmol) in DMF (5mL) was added ethyl chloroformate (0.017mL, 0.17mmol) and triethylamine (0.052mL, 0.037mmol) and the reaction was stirred 72h. The mixture was diluted with ethyl acetate and

35

washed successively with water and brine and dried (MgSO<sub>4</sub>). Purification by chromatography on silica gel using 3-10% methanol/methylene chloride as eluent afforded 27mg (33%) of the title compound. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.62 (s, 1H), 9.18 (s, 1H), 8.16 (s, 1H), 8.05 (m, 2H), 7.70 (d, 2H), 7.60 (5H, m), 7.37 (d, 2H), 7.30 (d, 1H), 7.24 (s, 2H), 6.95 (s, 1H), 4.10 (q, 2H), 2.35 (s, 3H), 1.20 (t, 3H) ppm; HRMS 547.176365 (calcd), 547.178880 (obs.).

10

#### Examples 182 and 183

1- (3- (1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt and 1- (3- (1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole, trifluoroacetic acid salt

Part A: The morpholino amidine compound was prepared from the precursor nitrile via the standard Pinner reaction protocol, with anhydrous morpholine as the nucleophile. Standard HPLC purification then afforded the desired morpholino amidine compound as colorless crystals; <sup>1</sup>HNMR(DMSO) δ: 11.39 (s, 1H); 9.67 (s, 1H); 9.27 (s, 1H); 8.62 (s, 2H); 8.09 (d, J=7.69, 1H); 7.79 (s, 1H); 7.73-7.61 (m, 5H); 7.42 (d, J = 7.32, 1H); 7.30 (s, 1H); 7.08 (s, 1H); 3.81 (brd, 2H); 3.74 (brd, 2H); 3.63 (brd, 2H); 3.37 (brd, 2H); 2.31 (s, 3H); 1.04 (s, 9H) ppm; ESI mass spectrum analysis m/z (rel intensity) 603.2 (M+H, 100).

Part B: Removal of the tert-butyl group was then effected by heating the product from part A in TFA, followed by standard HPLC purification techniques afforded the desired morpholino amidine compound as colorless crystals; <sup>1</sup>HNMR(DMSO) δ: 11.38 (s, 1H); 9.67 (s, 1H); 9.27 (s, 1H); 8.65 (s, 2H); 8.08 (m, 1H); 7.78 (s, 1H); 7.73-7.67 (m, 5H); 7.62 (m, 1H); 7.55 (s, 1H); 7.45 (m, 1H); 7.09 (s, 1H); 3.81 (brd, 2H); 3.74 (brd, 2H); 3.62 (brd, 2H); 3.37 (brd, 2H); 2.31 (s, 3H) ppm; ESI mass spectrum analysis m/z (rel intensity) 547.0 (M+H,

100). HRMS for  $C_{26}H_{27}N_8O_4S$  547.187599 (calcd), 547.186294 (obs).

#### Example 184

5

1-[3-[N-((5-methyl-2-oxo-1,3-dioxol-4-yl)methoxycarbonyl)amidino]phenyl]-5-((2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-methylpyrazole

Part A. To 4-hydroxymethyl-5-methyl-1,3-dioxol-2-one (0.227g, 1.75mmol) (Alpegiani, M. et al, Syn. Com. 1992, 22 (9), 1277) in chloroform (5mL) at 0°C was added pyridine (0.15mL) and 4-nitrophenyl chloroformate (0.387g, 1.9mmol). The reaction was allowed to warm to ambient temperature and was stirred 18h. The reaction mixture was washed with water, brine and dried (Na<sub>2</sub>SO<sub>4</sub>). The crude dioxolone was used in the next step.

Part B. To 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole (80mg, 0.14mmol) in DMF (1mL) was added the dioxolone from part A and triethylamine (0.038 mL). The reaction was stirred 18h. The reaction was diluted with ethyl acetate and washed with water and dried (MgSO<sub>4</sub>). Purification by chromatography on silica gel using 3-5% methanol in methylene chloride afforded 47mg (55%) of the title dioxolone. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.63 (s, 1H), 8.25 (s, 1H), 8.05 (t, 2H), 7.62 (d, 2H), 7.50 (m, 5H), 7.37 (m, 4H), 7.25 (s, 2H), 6.93 (s, 1H), 4.92 (s, 2H), 2.37 (s, 3H), 2.15 (s, 3H) ppm; HRMS 631.161109 (calcd), 631.160927 (obs.).

30

#### Example 185

1-(Pyrid-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

The title compound was prepared by previously described methodology using 2-pyridine hydrazine·HCl. LRMS (M+H)<sup>+</sup> m/z: 452.

## Example 186

1-(6-Bromopyridin-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

5 By using previously described methodology, ethyl 3-methyl-1-(pyridin-2-yl)-1H-pyrazolecarboxylate was obtained. This compound was then treated with N-bromosuccinamide according to the following procedure.

10 A mixture of 3-methyl-1-(pyridin-2-yl)-1H-pyrazolecarboxylic acid (7.0483 mmol, 1.63 g) and N-bromosuccinimide (2.51 g, 2.0 eq.) in carbon tetrachloride (40 mL) was stirred at ambient temperature for 18 h. The reaction mixture was filtered through celite to remove solid impurity and washed with carbon tetrachloride (30 mL). The filtrate  
15 was evaporated and purified by flash chromatography on a silica gel column (200 g) eluted with 3:1 hexane:ethyl acetate to give 0.258 g of pure 3-methyl-1-(6-bromopyridin-2-yl)-1H-pyrazolecarboxylic acid (12 %).

20 Thereafter, following previously described procedures the acid chloride of 3-methyl-1-(6-bromopyridin-2-yl)-1H-pyrazolecarboxylic acid was coupled with 3-fluoro-4-((2-N-t-butylsulfonamido)phenyl)aniline, and t-butyl protecting group removed with refluxing trifluoroacetic acid to obtain the title compound; LRMS (M+H)<sup>+</sup> m/z: 530.

25

## Example 187

1-(3-amino-4-chlorophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

30

The title compound was prepared by the same method described in Example 174. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.90 (s, 1H); 8.02 (d, 1H); 7.78 (d, 1H); 7.62 (m, 2H); 7.55 (s, 1H); 7.26-7.34 (m, 5H); 7.03 (s, 1H); 6.81 (d, 1H), 5.89 (bs, 2H). High  
35 resolution mass spectrum analysis: calcd 504.0412, found 504.0411.



## Example 188

1-(3-amino-4-chlorophenyl)-5-[(4'-(1-pyrrolidinocarbonyl)phenyl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

5

The title compound was prepared by the same method described in Example 174. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.26 (bs, 1H); 7.80 (t, 1H); 7.49 (d, J= 11.0 Hz, 1H); 7.42 (d, J= 8.4 Hz, 1H); 7.40 (d, J= 8.1 Hz, 1H); 7.04 (d, J= 2.6 Hz, 1H); 6.79 (dd, J= 8.4 and 2.6 Hz, 1H); 3.45 (t, J= 6.2 Hz, 2H), 3.40 (t, J= 5.8 Hz, 2H), 1.85 (m, 4H). ESI mass spectrum analysis m/z (relative intensity): 430.0 (M+H)<sup>+</sup>; 452.0, (M+Na)<sup>+</sup>.

10

## Example 189

15 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

1-(3-cyanophenyl)-5-[2'-(t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole prepared as shown in Part B of Example 24 (0.20 g, 0.40 mmol) was dissolved in 10 mL of EtOAc and 10 mL of EtOH. TFA (1 mL) and Palladium on carbon (10 %) were added. The mixture was hydrogenated at 30 psi for 18 h. The reaction mixture was filtered through celite and washed with EtOAc. The filtrate was concentrated to a brown oil. It was dissolved in 5 mL of TFA and refluxed under N<sub>2</sub> for 30 minutes. The solvent was removed in vacuo and the resulting material was purified by reversed phase HPLC to give 59.8 mg of the title compound with 98% purity. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.54 (s, 1H); 8.25 (bs, 3H); 8.02 (d, J= 6.3 Hz, 1H); 7.84 (bs, 1H); 7.77 (t, J= 5.8 Hz, 2H); 7.72 (t, J= 6.9 Hz, 2H); 7.60 (m, 2H); 7.39 (d, J = 8.8 Hz, 2H), 7.32 (m, 1H), 7.31 (s, 2H), 4.18, (bs, 2H). ESI mass spectrum analysis m/z (relative intensity): 450.2 (M+H, 100)<sup>+</sup>.

20

25

30

35

## Example 190

1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole, trifluoroacetic acid salt

The title compound was prepared by the same method described in Example 189. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.28 (s, 1H); 8.23 (bs, 3H); 7.99 (d, J= 6.6 Hz, 1H); 7.80 (bs, 1H); 7.70 (m, 2H); 7.60 (m, 2H); 7.41 (s, 2H); 7.31 (d, J = 9.5 Hz, 2H), 7.20 (d, J = 8.1 Hz, 1H), 4.14, (m, 2H). ESI mass spectrum analysis m/z (relative intensity): 467.9, (M+H, 100)+.

#### Example 191

1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole, trifluoroacetic acid salt

Part A: A solution of 3-amino-benzonitrile (6.3 g, 53.4 mmol) in ethyl alcohol (50 mL) was treated with n-butyl glyoxylate (7.0 g, 53.8 mmol). After stirring for 18h at rt, the reaction mixture was concentrated at reduced pressure. The residue was purified by flash-chromatography (hexane/ethyl acetate, 1:1) affording an imine (4.0 g, 33%) as a colorless oil. ESI mass spectrum analysis m/z (relative intensity): 232 (M+H, 100).

Part B: To the solution of the imine from part A (1.6 g, 6.9 mmol) in methyl alcohol (10 mL) was added potassium carbonate (1.9 g, 13.9 mmol) and tosylmethyl isocyanate (2.3 g, 11.8 mmol). The solution was stirred for 1h at rt, then solvent was removed under reduced pressure. The residue was treated with the saturated sodium chloride solution and the mixture was extracted with methylene chloride. The organic extract was concentrated and triturated with methyl alcohol. The precipitate was collected and dried to afford the desired methyl 1-(3-cyanophenyl)-imidazole-5-carboxylate (1.5 g, 94%). ESI mass spectrum analysis m/z (relative intensity): 227 (M+H, 100)

Part C: A solution of (2'-tert-butylaminosulfonyl-[1-1']-biphen-4-yl)amine (3.5 mmol) in methylene chloride (3 mL) was treated dropwise with AlMe<sub>3</sub> (2M in hexanes, 1.8 mL, 3.5 mmol). The resultant reaction mixture was stirred for 0.5h at rt, then treated with the product from part B (0.16 g, 0.7 mmol) and allowed to stir for 18h. The mixture was carefully quenched

with 10% HCl, extracted with methylene chloride, dried over magnesium sulfate and concentrated. Purification by flash chromatography (methanol/methylene chloride, 1:9) afforded the coupled amide compound (0.22 g, 28%). ESI mass spectrum analysis m/z (relative intensity): 500 (M<sup>+</sup>, 100). Reduction of the benzonitrile to the benzylamine followed by standard HPLC purification protocols via methods previously described afforded pure titled compound as colorless crystals. <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 8.61 (bs, 1H), 8.14 (bs, 1H), 8.09 (dd, J = 7.7Hz, 1H), 7.65-7.50 (m, 12H), 7.40 (dd, J = 8.8Hz, 2H), 7.32 (dd, J = 7.3Hz, 1H), 4.91 (s, 3H)ppm. ESI mass spectrum analysis m/z (relative intensity): 448.2 (M+H, 100).

#### Example 192

15     1-(3-aminomethylphenyl)-5-[(2'-methylsulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole, trifluoroacetic acid salt

The title compound was prepared in a similar manner to Example 197. <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 8.57 (s, 1H), 8.15 (m, 2H), 7.72-7.58 (m, 12H), 7.40 (m, 3H), 4.22 (s, 2H), 2.72 (s, 3H)ppm. ESI mass spectrum analysis m/z (relative intensity): 447 (M+H, 100).

#### Example 193

25     1-(3-amidinophenyl)-5-[(2'-aminosulfonyl[1,1']-biphen-4-yl)aminocarbonyl]imidazole, trifluoroacetic acid salt

The benzonitrile obtained in part C in Example 197 was subjected to the Pinner-amidine reaction protocol and further purified via methods described previously to obtain the title compound as colorless crystals. ESI mass spectrum analysis m/z (relative intensity): <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 8.76 (s, 1H), 8.21 (s, 1H), 8.07 (d, J = 7.7Hz, 1H), 7.98 (d, J = 8.4Hz, 1H), 7.89 (d, J = 8.4Hz, 1H), 7.79 (t, J = 7.7Hz, 1H), 7.59 (m, 3H), 7.50 (t, J = 7.7Hz, 1H), 7.38 (d, J = 8.5Hz, 2H), 7.30 (d, J = 8.7Hz, 1H)ppm. ESI mass spectrum analysis m/z (relative intensity): 461.2 (M+H, 100).

## Example 194

1-[3-(methylaminomethyl)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-  
[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole,  
trifluoroacetic acid salt

5

Part A. Preparation of ethyl 1-[3-(N-t-butoxycarbonyl-aminomethyl)phenyl]-3-methylpyrazolecarboxylate.

To a solution of 1.52 g (5.14 mmol) of ethyl 1-[3-(aminomethyl)phenyl]-3-methylpyrazolecarboxylate hydrochloride in 10 mL of THF under N<sub>2</sub> was added 1.49 g (14.7 mmol) of triethylamine and 1.35 g (6.17 mmol) di-t-butyl dicarbonate. The mixture was allowed to stir at room temperature for 16 hours. Water (25 mL) was added and the mixture was extracted with 25 mL ether three times. The combined organic extracts were dried over MgSO<sub>4</sub> and the solvent evaporated to give the desired product (1.85 g, 74%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.34 (m, 4H); 6.81 (s, 1H); 4.87 (b s, 1H); 4.37 (d, J = 7, 2H); 4.22 (q, J = 7, 2H); 2.35 (s, 3H); 1.45 (t, 9H); 1.24 (t, J = 7, 3H).

Part B. Preparation of ethyl 1-[3-(N-t-butoxycarbonyl-N-methylaminomethyl)phenyl]-3-methylpyrazolecarboxylate.

To a solution of 1.85 g (5.15 mmol) of ethyl 1-[3-(N-t-butoxycarbonylaminomethyl)phenyl]-3-methylpyrazolecarboxylate in 10 mL of THF under N<sub>2</sub> was added 0.15 g (5.88 mmol) of 95% sodium hydride. After 1 hour, the gas evolution ceased and 0.83 g (5.88 mmol) of methyl iodide was added. The mixture was allowed to stir at room temperature for 16 hours. Water (25 mL) was added and the mixture was extracted with 25 mL ether three times. The combined organic extracts were dried over MgSO<sub>4</sub> and the solvent evaporated and then chromatographed with 20% EtOAc/hexanes on silica to give the desired product (0.52 g, 27%) as a white solid. An additional 0.83 g of non-methylated starting material was also isolated. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.40 (m, 1H); 7.30 (m, 3H); 6.81 (s, 1H); 4.47 (b s, 2H); 4.22

(q, J = 7, 2H); 2.83 (b m, 3H); 2.34 (s, 3H); 1.47 (b s, 9H); 1.23 (t, J = 7, 3H).

Part C. Preparation of 1-[3-(N-t-butoxycarbonyl-N-methylaminomethyl)phenyl]-3-methylpyrazolecarboxylic acid.

To a solution of 0.52 g (1.39 mmol) of ethyl 1-[3-(N-t-butoxycarbonyl-N-methylaminomethyl)phenyl]-3-methylpyrazolecarboxylate in 5 mL of THF was added 1.4 mL (1.4 mmol) of 1M aqueous lithium hydroxide. The mixture was allowed to stir at room temperature for 6 hours. Water (10 mL) was added and the mixture was extracted with 25 mL ether twice. The aqueous layer was acidified with 1N HCl to pH 4 and extracted with 25 mL ether three times. The combined organic layers from the second set of extractions were dried over MgSO<sub>4</sub> and the solvent evaporated to give the desired product (0.35 g, 74%) as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.38 (m, 4H); 6.87 (s, 1H); 4.46 (b s, 2H); 2.83 (b m, 3H), 2.37 (s, 3H), 1.46 (b s, 9H).

Part D. Preparation of 1-[3-(methylaminomethyl)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methyl)pyrazolecarboxamide, trifluoroacetic acid salt.

To a solution of 1-[3-(N-t-butoxycarbonyl-N-methylaminomethyl)phenyl]-3-methylpyrazolecarboxylic acid (0.176 g, 0.509 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 10μL of DMF and oxalyl chloride (97 mg, 0.763 mmol). The solution was allowed to stir for 1.5 hours under Ar and then solvent was evaporated under high vacuum. The resulting solid was redissolved in 10 mL and triethylamine (0.15 g, 1.53 mmol) and 2'-(t-butylaminosulfonyl)-3-fluoro-[1,1']-biphenyl (0.172 g, 0.534 mmol) were added. After stirring for 16 hours under Ar, the reaction mixture was added to water and extracted with ethyl acetate. The solvent was evaporated and the mixture was dissolved in 5 mL of TFA. This solution was heated to 50°C for 4 hours, cooled to room temperature and the solvent evaporated. The crude benzylamine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 60 mg (19%) of the

desired salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.75 (br s, 2H); 8.00 (m, 1H); 7.63-7.15 (m, 10H); 6.94 (s, 1H); 4.15 (b t, J = 6, 2H); 2.54 (t, J = 5, 2H); 2.45 (s, 3H). ESI mass spectrum analysis m/z (relative intensity): 494.1 (M+H, 100).

5

**Example 195**

**1-[3-(methylaminomethyl)phenyl]-5-[(2'-methylsulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt**

10

To a solution of 1-[3-(N-t-butoxycarbonyl-N-methylaminomethyl)phenyl]-3-methylpyrazolecarboxylic acid (0.176 g, 0.509 mmol) in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 10μL of DMF and oxalyl chloride (97 mg, 0.763 mmol). The solution was allowed to stir for 1.5 hours under Ar and then solvent was evaporated under high vacuum. The resulting solid was redissolved in 10 mL and triethylamine (0.15 g, 1.53 mmol) and 2'-(methylsulfonyl)-3-fluoro-[1,1']-biphenyl (0.172 g, 0.534 mmol) were added. After stirring for 16 hours under Ar, the reaction mixture was added to water and extracted with ethyl acetate. The solvent was evaporated and the mixture was dissolved in 5 mL of TFA. This solution was heated to 50°C for 4 hours, cooled to room temperature and the solvent evaporated. The crude benzylamine was purified by HPLC (C18 reversed phase) eluting with 0.5% TFA in H<sub>2</sub>O/CH<sub>3</sub>CN to give 140 mg (45%) of the desired salt. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.76 (br s, 2H); 8.06 (dd, J = 8, 1, 1H); 7.77-7.61 (m, 4H); 7.52-7.31 (m, 5H); 7.19 (dd, J = 8, 1.5, 1H); 6.95 (s, 1H); 4.17 (b t, J = 6, 2H); 2.90 (s, 3H); 2.54 (t, J = 5, 2H); 2.29 (s, 3H). ESI mass spectrum analysis m/z (relative intensity): 492.2 (M+H).

20  
25  
30**Example 196**

**1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-methoxy-3-trifluoromethyl pyrazole, trifluoroacetic acid salt**

35

Part A. To 1-(3-cyanophenyl)-4-methoxy-3-trifluoromethylpyrazole carboxylic acid (0.69 g, 2.2 mmol) was

added  $\text{CH}_2\text{Cl}_2$  (15 mL), oxalyl chloride (0.27 mL, 3.1 mmol), and three drops of DMF. The reaction was stirred for 2h. The solvents were removed and fresh  $\text{CH}_2\text{Cl}_2$  (15 mL), 4-bromo-aniline (0.38 g, 2.2 mmol) and DMAP (0.68 g, 5.5 mmol) were added and the reaction was stirred 18h. Dilution with  $\text{CH}_2\text{Cl}_2$ , followed by washing successively with 1N HCl, saturated  $\text{NaHCO}_3$ , brine, drying ( $\text{MgSO}_4$ ) and recrystallization with  $\text{CH}_2\text{Cl}_2$ /hexanes afforded 0.5 g (48%) pure product and 0.43 g from filtrate.  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 8.90 (s, 1H), 7.79 (m, 2H), 7.72 (dd, J = 1.83, 6.96 Hz, 1H), 7.63 (t, J = 8.06 Hz, 1H), 7.46 (s, 4H), 4.15 (s, 3H) ppm; ESI mass spectrum analysis m/z (relative intensity): 482-484 (M+H, 100).

Part B To the bromo compound (0.4 g, 0.86 mmol) from Part A was added 2-thiomethyl phenylboronic acid (0.18 g, 1.1 mmol), 2M  $\text{Na}_2\text{CO}_3$  (1 mL), toluene (15 mL), and ethanol (15 mL). The mixture was degassed and tetrakis(triphenylphosphine) palladium (0) (40 mg) was added and the reaction was heated to reflux 18h. The reaction was cooled, filtered, concentrated, and extracted with ethyl acetate and dried ( $\text{MgSO}_4$ ). The compound was purified by chromatography on silica gel eluting with (4:1) hexanes/ethyl acetate to afford 0.195 g (46%) yellow solid.  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 8.95 (s, 1H), 7.80 (m, 3H), 7.63 (d, J = 8.42 Hz, 2H), 7.61 (m, 1H), 7.44 (d, J = 8.43 Hz, 2H), 7.34 (m, 2H), 7.20 (m, 2H), 4.15 (s, 3H), 2.37 (s, 3H) ppm.

Part C To the product (0.19 g, 0.37 mmol) of Part B in  $\text{CH}_2\text{Cl}_2$  (15 mL), cooled to  $0^\circ\text{C}$ , m-chloroperbenzoic acid (0.33 g, 1.1 mmol) was added. The reaction warmed to ambient temperature overnight. The reaction was washed with water, sodium bisulfite solution,  $\text{NaHCO}_3$  and dried ( $\text{MgSO}_4$ ). The compound was purified by chromatography on silica gel eluting with (1:1) hexanes/ethyl acetate to afford 0.192 g (95%) yellow solid.  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 9.02 (s, 1H), 8.24 (dd, J = 1.46, 7.69 Hz, 1H), 7.80 (m, 3H), 7.66 (d, J = 8.06 Hz, 2H), 7.65 (m, 3H), 7.49 (d, J = 8.79 Hz, 2H), 7.37 (dd, J = 1.46, 7.69 Hz, 1H), 4.18 (s, 3H), 2.68 (s, 3H); ESI mass spectrum analysis m/z (relative intensity): 563 (M+Na, 100).

Part D The product of Part C was hydrogenated in EtOH/TFA with 10% palladium on carbon catalyst at 50 psi for 24h.

Purification by reverse phase HPLC and freeze-drying afforded the title compound 0.16 g (62.6%), <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.11 (s, 1H), 8.25 (brd s, 2H), 8.10 (d, J = 8.06Hz, 1H), 7.77 (s+d, J = 8.79Hz, 2H), 7.69 (s+d, J = 7.32Hz, 3H), 7.60 (s+m, 3H), 7.41 (m, 3H), 4.15 (brd s, 2H), 3.95 (s, 3H) 2.88 (s, 3H) ppm; HRMS 545.147037 (calc'd), 545.146284 (obs.); Elemental analysis calc'd for C<sub>26</sub>H<sub>23</sub>F<sub>3</sub>N<sub>4</sub>O<sub>4</sub>S(TFA)(H<sub>2</sub>O)1.3: C:49.31, H:3.93, N:8.22, found C:49.46, H:3.62, N:8.09.

#### Example 197

1- (3-aminomethylphenyl)-5-[(2-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

Part A. To 1-(3-cyanophenyl)-3-trifluoromethylpyrazole carboxylic acid (0.29 g, 1.0 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added oxalyl chloride (0.135 mL, 1.6 mmol) and several drops DMF. The reaction was stirred for 2h, then concentrated. To the acid chloride was added fresh CH<sub>2</sub>Cl<sub>2</sub> (40 mL), 2-fluoro-4-(N-pyrrolidinocarbonyl)aniline (0.22 g, 1 mmol), and DMAP (0.32 g, 2.6 mmol) and the reaction was stirred 18h. The reaction was washed successively with 1N HCl, NaHCO<sub>3</sub>, and dried (MgSO<sub>4</sub>). The compound was purified by chromatography on silica gel eluting with (1:1.5) hexanes/ethyl acetate to afford 0.345 g (71%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 9.03 (s, 1H), 7.86 (m, 4H), 7.63 (t, J = 8.05Hz, 1H), 7.55 (s, 1H), 7.21 (m, 2H), 3.67 (t, J = 8.05Hz, 2H), 3.43 (t, J = 6.59Hz, 2H), 2.02 (q, J = 6.22Hz, 2H), 1.92 (q, J = 6.22Hz, 2H) ppm; ESI mass spectrum analysis m/z (relative intensity): 472.1 (M+H)<sup>+</sup>, 494 (M+Na)<sup>+</sup>.

Part B. The product of Part A was hydrogenated in EtOH/TFA with 10% palladium on carbon catalyst at 50 psi for 24h. Purification by reverse phase HPLC and freeze-drying afforded 0.34 g (80%) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.8 (s, 1H), 8.23 (s, 2H), 7.72 (m+d, J = 8.06Hz, 3H), 7.59 (m, 3H), 7.49 (dd, J =



- 1.84, 11.36Hz, 1H), 7.39 (dd, J = 8.06, 1.83Hz, 1H), 4.15 (q, J = 5.86Hz, 2H), 3.47 (t, J = 6.6Hz, 2H), 3.42 (t, J = 6.2Hz, 2H), 1.89 (m, 4H) ppm; ESI mass spectrum analysis m/z: 476.2 (M+H)<sup>+</sup>; Elemental analysis calc'd for C<sub>23</sub>H<sub>21</sub>F<sub>4</sub>N<sub>5</sub>O<sub>2</sub> (TFA) (H<sub>2</sub>O) 0.5: C:50.17, H:3.87, N:11.70, found C:50.05, H:3.87, N:11.43

#### Example 198

1-(3-aminomethylphenyl)-5-[(3-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

- Part A. 1-(3-Cyanophenyl)-3-trifluoromethylpyrazole carboxylic acid and 3-fluoro-4-(N-carboxylpyrrolidino)aniline were coupled via the acid chloride as in the previous Example in 81% yield.
- <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ 10.01 (s, 1H), 7.79 (m, 4H), 7.61 (t, J = 7.69Hz, 1H), 7.16 (dd, J = 1.84, 10.99Hz, 1H), 7.06 (t, J = 8.06Hz, 1H), 6.93 (dd, J = 1.83, 8.05Hz, 1H), 3.68 (t, J = 6.59Hz, 2H), 3.34 (t, J = 6.59Hz, 2H), 2.00 (q, J = 6.59Hz, 2H), 1.94 (q, J = 6.59Hz, 2H) ppm; ESI mass spectrum analysis m/z (relative intensity): 472.1 (M+H)<sup>+</sup>, 494 (M+Na)<sup>+</sup>.

- Part B. The product of Part A was hydrogenated in EtOH/TFA with 10% palladium on carbon catalyst at 50 psi for 24h. Purification by reverse phase HPLC and freeze-drying afforded 0.38 g (84%) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.07 (s, 1H), 8.24 (s, 2H), 7.73 (m, 3H), 7.63 (m, 3H), 7.50 (m, 2H), 4.16 (d, J = 5.49Hz, 2H), 3.47 (t, J = 6.23Hz, 2H), 3.23 (t, J = 6.23Hz, 2H), 1.89 (m, 4H) ppm; HRMS 476.170963 (calc'd), 476.171044 (obs.); Elemental Analysis calc'd for C<sub>23</sub>H<sub>21</sub>F<sub>4</sub>N<sub>5</sub>O<sub>2</sub> (TFA) (H<sub>2</sub>O) 0.5: C:50.17, H:3.87, N:11.70, found C:50.17, H:3.85, N:11.48.

#### Example 199

1-(3-aminomethylphenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole, trifluoroacetic acid salt

1-(3-Cyanophenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole (synthesis

previously described) was hydrogenated in EtOH/TFA with 10% palladium on carbon catalyst at 50 psi for 24h. Purification by reverse phase HPLC and freeze-drying afforded the title compound. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.92 (s,1H), 8.24 (bd s,2H), 8.10 (d, J = 7.69Hz,1H), 7.79 (m,6H), 7.60 (m,3H), 7.41 (s+d, J = 8.79Hz,3H), 4.17 (q, J = 5.12Hz,2H), 2.85 (s,3H)ppm, HRMS 515.136472 (calc'd), 515.137193 (obs).

#### Example 200

10 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']  
biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole,  
trifluoroacetic acid salt

Part A. 1-(3-Cyanophenyl)-3-trifluoromethylpyrazole carboxylic  
15 acid and 1-(2'-tertbutylaminosulfonyl-[1,1']-3-  
fluorobiphenylaniline were coupled via the acid chloride as  
in previous Examples in 76% yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.31 (t, J =  
8.43Hz,1H), 8.18 (dd, J = 1.47,7.69Hz,1H), 8.04 (s,1H), 7.88 (d,  
J = 1.46Hz,1H), 7.83 (m,2H), 7.68 (d, J = 8.06Hz,1H), 7.62  
20 (m,2H), 7.42 (dd, J = 1.83,11.72Hz,1H), 7.29 (d, J =  
1.47Hz,1H), 7.22 (m,2H), 3.69 (s,1H), 1.07 (s,9H)ppm; ESI mass  
spectrum analysis m/z (relative intensity): 607.9 (M+Na, 100).

Part B. The product of Part A was refluxed in TFA for 30  
25 minutes then hydrogenated in EtOH/TFA with 10% palladium on  
carbon catalyst at 50 psi for 24h and then with platinum (II)  
oxide catalyst at 50 psi for 24h. Purification by reverse  
phase HPLC and freeze-drying afforded 0.16 g (44%) product.  
<sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.71 (s,1H), 8.24 (bd s,2H), 8.05 (dd, J =  
30 1.47,6.96Hz,1H), 7.74 (s,1H), 7.69 (s,1H), 7.66 (m,6H), 7.43  
(s,2H), 7.35 (m,2H), 7.23 (d, J = 8.42Hz,1H), 4.16 (q, J =  
5.49Hz,2H)ppm; ESMS 534.1 (M+H); Elemental Analysis calc'd for  
C<sub>24</sub>H<sub>19</sub>F<sub>4</sub>N<sub>5</sub>O<sub>3</sub>S(TFA)1.1 (H<sub>2</sub>O)0.6: C:46.99, H:3.21, N:10.46, found  
C:47.06, H:2.86, N:10.37.

35

#### Examples 201 and 202

1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)-[1,6-  
dihydro]pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole,

trifluoroacetic acid salt and 1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole, trifluoroacetic acid salt.

- 5 1-(3-Cyanophenyl)-5-[(5-(2'-tertbutylaminosulfonylphenyl)-4-yl)pyrimid-2-yl)aminocarbonyl]-3-trifluoro-methyl pyrazole (0.3 g, 0.5 mmol) (synthesis previously described) was hydrogenated in ethanol/acetic acid for 24h at 40 psi, first with 10% palladium on carbon and then with added platinum (II) oxide. The reaction was filtered, concentrated, and refluxed in TFA for 30 minutes. Purification by reverse phase HPLC and freeze-drying afforded small amounts of two products. The dihydro-compound was the first product obtained (64.5 mg). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 9.76 (s, 1H), 9.10 (s, 1H), 8.22 (brd, 2H), 7.95 (dd, J = 1.10, 7.69Hz, 1H), 7.65 (s, 1H), 7.61 (m, 5H), 7.49 (s, 2H), 7.41 (dd, J = 1.46, 7.32Hz, 1H), 7.19 (s, 1H), 6.10 (d, J = 4.40Hz, 1H), 4.22 (s, 2H), 4.15 (q, J = 5.86Hz, 2H)ppm; HRMS 520.137869 (calc'd); 520.138256 (obs); Elemental Analysis calc'd for C<sub>22</sub>H<sub>20</sub>F<sub>3</sub>N<sub>7</sub>O<sub>3</sub>S(TFA)<sub>2</sub>: C:41.77, H:2.97, N:13.12, found C:41.98, H:3.02, N:12.97. The second product was the pyrimidyl analog. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.61 (s, 1H), 8.66 (s, 2H), 8.24 (brd, 2H), 8.08 (dd, J = 2.20, 6.95Hz, 1H), 7.73 (m, 4H), 7.60 (m, 5H), 7.48 (m, 1H), 4.16 (m, 2H); HRMS 518.122219 (calc'd); 518.122803 (obs); Elemental Analysis calc'd for C<sub>22</sub>H<sub>18</sub>F<sub>3</sub>N<sub>7</sub>O<sub>3</sub>S(TFA)<sub>1.3</sub> (H<sub>2</sub>O) C:43.79, H:3.03, N:14.53, found C:43.92, H:2.99, N:14.37.

#### Example 203

- 1-[3-(2'-ethylaminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole, trifluoroacetic acid salt.

- Part A. To 1-(3-cyanophenyl)-5-hydroxymethyl-3-trifluoromethyl pyrazole (1.8 g, 6.7 mmol) in DMF (12 mL) was added tert-butyldimethylsilylchloride (1 g, 7.1 mmol) and imidazole (0.94 g, 13.8 mmol). The reaction was stirred for 3h, then partitioned between ethyl acetate and water. Extraction with ethyl acetate, drying (MgSO<sub>4</sub>) and purification

by chromatography on silica gel eluting with (4:1) hexanes/ethyl acetate to afford 1.88 g (73%).

Part B. To the product of Part A (0.4 g, 1.0 mmol) in THF (15 mL) at 0°C was added methyl-magnesium chloride (0.9 mL, 2.6 mmol) and the reaction was stirred at ambient temperature for 2h. After cooling to 0°C, methanol (25 mL) and then sodium borohydride (0.2 g, 5 mmol) were added and the reaction was stirred for 1h. The reaction was quenched with water, filtered and concentrated. The residue was extracted into ethyl acetate and dried (MgSO<sub>4</sub>). The crude oil was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, cooled to 0°C and di-tert-butylcarbamate (0.23 g, 1.1 mmol) and triethylamine (0.15 mL) were added. The reaction was stirred 18h, then washed with saturated ammonium chloride, brine and dried (MgSO<sub>4</sub>). The crude material was dissolved in THF and tetrabutylammonium fluoride in THF (1.46 mL) was added. The reaction stirred for 3h, then concentrated. The residue was dissolved in CH<sub>2</sub>Cl<sub>2</sub>, and washed with water, brine and dried (MgSO<sub>4</sub>). Purification by chromatography on silica gel eluting with (2:1) hexanes/ethyl acetate afforded 0.187 g (47%).  
<sup>1</sup>H NMR (CDCl<sub>3</sub>) δ 7.58 (s, 1H), 7.47 (m, 2H), 7.38 (m, 1H), 6.70 (s, 1H), 4.92 (bd, 1H), 4.78 (m, 1H), 4.65 (m, 2H), 2.91 (bd, 1H), 1.49 (d, J = 6.96 Hz, 3H), 1.40 (s, 9H) ppm; MS ESI mass spectrum analysis m/z (relative intensity): 407.8 (M+Na, 100).

Part C. To the product of Part B (0.17 g, 0.44 mmol) in acetonitrile (5 mL) at 0°C was added a few crystals of ruthenium (III) chloride and aqueous solution of sodium periodate (0.2 g, 0.9 mmol). The reaction was stirred 18h, then filtered and concentrated. The aqueous residue was extracted with ethyl acetate and dried (MgSO<sub>4</sub>). ESI (-ve) mass spectrum analysis m/z (relative intensity): 398 (M-H, 100).

Part D. To the product of Part C (0.17 g, 0.4 mmol) and 4-bromoaniline (0.073 g, 0.4 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.11 g, 0.57 mmol). The reaction was stirred 18h, then washed with water, brine and dried (MgSO<sub>4</sub>). Filtration through a plug of

silica gel eluting with (1:1) hexanes/ethyl acetate afforded 0.148 g of a white foam. ESI mass spectrum analysis m/z (relative intensity): 575-577 (M+Na)<sup>+</sup>.

- 5 Part E. The product of Part D (0.14 g, 0.26 mmol) was coupled to 2-tert-butylsulfonamide phenyl boronic acid by standard Suzuki procedure. The crude product of this reaction was heated to reflux in TFA for 20 minutes. Purification by reverse phase HPLC and freeze-drying afforded 77 mg product (46%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.86 (s, 1H), 8.32 (brd, 2H), 8.04 (dd, j=7.69, 1.42Hz, 1H), 7.76 (s, 1H), 7.68 (d, j=8.42Hz, 2H), 7.67 (m, 6H), 7.39 (d, J = 8.79Hz, 2H), 7.32 (dd, J = 9, 1.32Hz, 1H), 7.29 (s, 2H), 4.56 (m, 1H), 1.52 (d, J = 6.96Hz, 3H)ppm; HRMS 530.147371 (cal'd), 530.148939 (obs); Elemental Analysis calc'd for C<sub>25</sub>H<sub>22</sub>F<sub>3</sub>N<sub>5</sub>O<sub>3</sub>S(TFA)1.1: C:49.88, H:3.55, N:10.69, found C:49.49, H:3.49, N:10.60.

#### Example 204

- 1-[[3-(1-(N-morpholino)imino)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt.

- To 1-(3-cyanophenyl)-5-[(2'-tert-butylaminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole (0.23 g, 0.39 mmol) in (2:1) CHCl<sub>3</sub>/MeOH (30 mL) at 0°C was bubbled HCl gas for 15 minutes. The flask was sealed and placed in the refrigerator for 18h. The solvent was removed and morpholine (0.2 mL) and fresh methanol were added. The reaction was stoppered and stirred for 48h. The solvent was removed and the residue was heated to reflux in TFA for 15 minutes. Purification by reverse phase HPLC and freeze-drying afforded 0.146 g product (51%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 10.70 (s, 1H), 9.69 (s, 1H), 9.32 (s, 1H), 8.05 (dd, j=6.96, 2.20Hz, 1H), 7.94 (s, 1H), 7.89 (d, J = 8.05Hz, 1H), 7.80 (m, 2H), 7.65 (m, 3H), 7.42 (s, 2H), 7.35 (d, J = 8.50Hz, 2H), 7.23 (d, J = 9.52Hz, 1H), 3.81 (bs, 2H), 3.74 (bd s, 2H), 3.56 (bd s, 2H), 3.32 (bd s, 2H)ppm; ESMS 616.9 (M+H). Elemental Analysis calc'd for

$C_{28}H_{24}F_4N_6O_4S(TFA)1.1$  ( $H_2O$ )1.2: C:47.50, H:3.63, N:11.01, found C:47.39, H:3.28, N:10.69.

#### Example 205

5    1-(3-aminomethylphenyl)-5-[2-(2'-aminosulfonyl-[1,1']-biphen-4-yl)-1-hydroxyethyl]-3-trifluoromethyl-pyrazole, trifluoroacetic acid salt

Part A. To 1-(3-cyanophenyl)-3-trifluoromethylpyrazole-5-carboxylic acid (1 g, 3.6 mmol) in  $CH_2Cl_2$  (40 mL) was added oxalyl chloride (0.4 mL, 4.9 mmol) and several drops of DMF. The reaction was stirred for 3h, then the solvent was removed in vacuo. In a separate flask, dibromoethane (0.1 mL), was added to activated Zn (0.35 g, 5.3 mmol) in THF (5mL). The mixture was heated to reflux for 5 minutes, then cooled to 0°C and 4-bromo-benzylbromide (1.1 g, 4.3 mmol) in THF (5 mL) was added slowly over 0.5h. The reaction was kept at 0°C for 3h, then cannulated into a mixture of CuCN (0.38 g, 4.3 mmol), LiCl (0.36 g, 8.5 mmol) and THF (10 mL) at -78°C. The reaction was warmed to -20°C for 5 minutes, then recooled to -78°C. The solid acid chloride was suspended in THF (20 mL) and added to the above cold mixture. The reaction was allowed to slowly warm to room temperature, then, filtered and concentrated. Purification by chromatography on silica gel eluting with (2:1) hexanes/ethyl acetate afforded 0.55 g (37%) white foam. MS (ESI) m/z= 433.9-432 (M-H)<sup>+</sup>.

Part B. The product of Part A (0.53 g, 1.2 mmol) was coupled by standard Suzuki procedures to 2-tert-butylaminosulfonyl-phenyl boronic acid (0.39 g, 1.7 mmol). Purification by chromatography on silica gel eluting with (4:1) hexanes/ethyl acetate afforded 0.32g (46%) the keto-nitrile coupled product. MS (ESI) m/z= 565 (M-H)<sup>+</sup>.

Part C. To the product from Part B (0.05 g, 0.08 mmol) was added  $CH_2Cl_2$  (10mL) and tetra-N-butylammonium borohydride (0.08 g, 0.31 mmol) and the mixture was heated to reflux 18h. The solvent was removed and replaced with 10% HCl and heated to

reflux for 1h. The reaction was cooled, extracted with diethyl ether, basefied with 50% NaOH, extracted with ethyl acetate and dried (MgSO<sub>4</sub>). The diethyl ether layer contained tert-butyl protected intermediate. The ether was concentrated and the residue heated in TFA for 15 minutes. All product was combined and purification by reverse phase HPLC and freeze-drying afforded 0.01 g of product (18%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.23 (brd, 2H), 8.03 (d, j=6.96Hz, 1H), 7.63 (m, 6H), 7.28 (s+d, j=7.69Hz, 3H), 7.18 (s, 2H), 7.11 (s+d, j=6.59Hz, 3H), 5.83 (m, 1H), 4.81 (m, 1H), 4.15 (m, 2H), 3.09 (d, j=6.60Hz, 2H)ppm; HRMS 517.152122(calc'd), 517.152222(obs.).

#### Example 206

1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole, trifluoroacetic acid salt

Part A. To 1-(3-cyanophenyl)-3-trifluoromethylpyrazole-5-carboxylic acid (1 g, 3.6 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added oxalyl chloride (0.43 mL, 4.9 mmol) and several drops of DMF. The reaction was stirred 18h, then the solvent was removed in vacuo. Fresh CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added followed by 4-bromo-2-fluoroaniline (0.68 g, 3.6 mmol) and 4-dimethylaminopyridine (1.09 g, 8.9 mmol). After stirring 18h, the reaction was washed with 1N HCl, sat'd NaHCO<sub>3</sub>, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated to afford 1.55 g crude bromide. ESI (-ve) mass spectrum analysis m/z (relative intensity) 450.8-452.8 (M-H, 100).

Part B. The bromide from Part A (0.5 g, 1.1 mmol), 2-thiomethyl phenylboronic acid (0.26 g, 1.5 mmol), and 2M Na<sub>2</sub>CO<sub>3</sub> (2 mL), were combined in (1:1) ethanol/toluene (20 mL) and degassed by bubbling nitrogen through for 30minutes. Tetrakis-triphenylphosphine palladium(0) (50 mg) was added and the reaction heated to reflux 18h. The reaction was cooled, concentrated, extracted with ethyl acetate and dried (MgSO<sub>4</sub>). The coupled product was purified through a plug of silica gel using (1:1) hexane/ethyl acetate as eluent and carried onto the

- next step. The thiomethyl compound was dissolved in  $\text{CH}_2\text{Cl}_2$  (50 mL), cooled to  $0^\circ\text{C}$ , and MCPBA (0.67 g, 2.2 mmol) was added. The reaction was stirred 48h, then washed successively with aqueous sodium bisulfite, brine, and dried ( $\text{MgSO}_4$ ). The sulfone was purified through a plug of silica gel using (1:1) hexane/ethyl acetate as eluent to afford 0.34 g.
- $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.25 (t, 1H), 7.90-7.15 (m, 12H), 2.39 (s, 3H) ppm. ESI mass spectrum analysis  $m/z$  550.7 ( $\text{M}+\text{Na}$ )<sup>+</sup>, 526.7 ( $\text{M}-\text{H}$ )<sup>+</sup>.
- 10 Part C. The product of Part B (0.34 g, 0.6 mmol) was hydrogenated in (1:2) methanol/ethanol (70 mL) and TFA (1 mL) with 10% palladium on carbon catalyst at 50 psi for 24h. Purification by reverse phase HPLC and freeze-drying afforded 0.21 g (50%) product.  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$  10.75 (s, 1H), 8.23 (m, 3H), 8.11 (dd,  $j=7.69$ , 1.46Hz, 1H), 7.96 (dd,  $j=6.96$ , 1.47Hz, 1H), 7.81 (m, 8H), 7.26 (dd,  $j=1.47$ , 8.06Hz, 1H), 4.16 (q,  $j=5.49\text{Hz}$ , 2H), 2.94 (s, 3H) ppm; ESI mass spectrum analysis  $m/z$  532.9 ( $\text{M}+\text{H}$ , 100); Elemental Analysis calc'd for  $\text{C}_{25}\text{H}_{20}\text{F}_4\text{N}_4\text{O}_3\text{S}(\text{TFA})1.1$ : C:49.65, H:3.23, N:8.52, found
- 15 C:49.73, H:2.98, N:8.40.
- 20

#### Example 207

- 1-((3-aminomethylphenyl)-5-[(5-(2'-methylsulfonyl-phenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl pyrazole, trifluoroacetic acid salt.
- 25

- Part A. 1-(3-Cyanophenyl)-3-trifluoromethylpyrazole carboxylic-5-acid (2.2, 7.8 mmol) was heated to reflux in methanol containing con. sulfuric acid (1 mL) for 48h. The solvent was removed and the residue was dissolved in ethyl acetate, washed with  $\text{NaHCO}_3$  (sat.), brine and dried ( $\text{MgSO}_4$ ). The ester was hydrogenated in MeOH/TFA with 10% palladium on carbon catalyst at 40 psi for 24h. The reaction was filtered and concentrated. The residue was suspended in  $\text{CH}_2\text{Cl}_2$ , cooled to  $0^\circ\text{C}$  and 1N NaOH (35 mL) and benzyl chloroformate (1.2 mL, 8.6 mmol) were added. The reaction was stirred 2h then separated and the organics dried ( $\text{MgSO}_4$ ) and concentrated. The residue was dissolved in MeOH, cooled to  $0^\circ\text{C}$  and a solution of
- 30
- 35



LiOH (0.5 g, 11.8 mmol) in water was added. The reaction was stirred 18h. The reaction was concentrated and the residue was acidified and extracted with ethyl acetate and dried (MgSO<sub>4</sub>) to afford 1.83 g (57%) white solid. ESI mass spectrum analysis  
5 m/z (relative intensity): 417.9 (M-H, 100).

Part B. The acid from Part A (0.46 g, 1.1 mmol) was coupled with 2-amino-5-(2'-methylsulfonylphenyl)pyrimidine (0.31 g, 1.1 mmol) by the standard acid chloride procedure to afford 0.3 g  
10 (42%) of the carbobenzyloxy protected intermediate. The intermediate was heated to reflux in TFA for 45 minutes and purification by reverse phase HPLC and freeze-drying afforded 0.16 g (23% overall) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.65 (s, 1H), 8.72 (s, 2H), 8.24 (bd, 2H), 8.15 (d, J = 7.69Hz, 1H), 7.87  
15 (m, 4H), 7.58 (s+m, 3H), 7.54 (d, J = 7.32Hz, 1H), 4.16 (q, J = 5.49Hz, 2H), 3.07 (s, 3H) ppm; HRMS 517.126970 (calc'd), 517.125600 (obs); Elemental Analysis calc'd for C<sub>23</sub>H<sub>19</sub>F<sub>3</sub>N<sub>6</sub>O<sub>3</sub>S(TFA) 1.2: C:46.70, H:3.12, N:12.86, found C:46.78, H:3.04, N:12.56.

20

#### Example 208

1-[3-amidinophenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole,  
25 trifluoroacetic acid salt

25

The nitrile prepared as in Example 206 was subjected to standard Pinner reaction conditions and purification by reverse phase HPLC and freeze-drying afforded 0.067 g (27%) of the desired titled product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.74 (s, 1H), 9.45  
30 (s, 1.5H), 9.13 (s, 1.5H), 8.11 (d, J = 7.69Hz, 1H), 8.04 (s, 1H), 7.95 (d, J = 8.42Hz, 2H), 7.81 (m, 5H), 7.44 (m, 2H), 7.26 (d, J = 8.42Hz, 1H), 2.94 (s, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity): 546 (M+H, 100).

35

#### Example 209

1-[3-amidinophenyl]-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole,  
trifluoroacetic acid salt

The nitrile prepared in Example 207 was subjected to standard Pinner reaction conditions and purification by reverse phase HPLC and freeze-drying afforded 0.042 g (25%) product.  
5 HRMS 547.117549 (calc'd), 547.117400 (obs).

#### Example 210

1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylmethyl]-3-trifluoromethyl pyrazole, trifluoroacetic  
10 acid salt

Part A. To the N-carbobenzyloxy protected carboxylic acid (5 g, 11.9 mmol) (described in Example 207) in  $\text{CH}_2\text{Cl}_2$  (100 mL) was added oxalyl chloride (1.5 mL, 16.7 mmol) and DMF (0.5 mL). The  
15 reaction was stirred 18h, then the solvents were removed and the resultant yellow solid set aside. In a separate flask, dibromoethane (0.3 mL), was added to activated Zn (1.87 g, 28 mmol) in THF (30 mL). The mixture was heated to reflux for 5 minutes, then cooled to  $0^\circ\text{C}$  and 4-bromo-benzylbromide (5.96  
20 g, 24.9 mmol) in THF (45 mL) was added slowly over 0.5h. The reaction was kept at  $0^\circ\text{C}$  for 3h, then cannulated into a mixture of CuCN (2.24 g, 25 mmol), LiCl (1.52 g, 36 mmol) and THF (15 mL) at  $-78^\circ\text{C}$ . The reaction was warmed to  $-20^\circ\text{C}$  for 5 minutes, then recooled to  $-78^\circ\text{C}$ . The solid acid chloride was suspended in  
25 THF (50 mL) and added to the above cold mixture. The reaction was kept at  $-78^\circ\text{C}$  for 1h,  $0^\circ\text{C}$  for 1h, then at  $20^\circ\text{C}$  for 1h. The reaction was quenched with saturated  $\text{NH}_4\text{Cl}$ , filtered, and extracted with ethyl acetate. The aqueous layer was carefully acidified, extracted with ethyl acetate and the combined  
30 organic layers dried ( $\text{Na}_2\text{SO}_4$ ). Purification by chromatography on silica gel eluting with (1:1) hexanes/ethyl acetate and recrystallization ( $\text{CH}_2\text{Cl}_2$ /hexanes) afforded 2.8 g pure product and 2.5 g slightly impure product from the filtrate.  
35  $^1\text{H}$ NMR ( $\text{CDCl}_3$ )  $\delta$ : 7.47 (d,  $j = 8.4\text{Hz}$ , 2H), 7.42 (m, 8H), 7.08 (d,  $J = 8.4\text{Hz}$ , 2H), 7.00 (d,  $J = 8.4\text{Hz}$ , 1H), 5.13 (s, 2H), 4.43 (d,  $J = 5.9\text{Hz}$ , 2H), 4.09 (s, 2H), 3.11 (AB,  $J = 13.5, 46.9\text{Hz}$ , 2H) ppm; ESI (-ve) mass spectrum analysis  $m/z$  (relative intensity): 569.7-571.6 (M-H) $^+$ .

Part B. The product of Part A (0.5 g, 0.88 mmol) was coupled by standard Suzuki procedures with 2-tert-butylaminosulfonylphenyl boronic acid (0.3 g, 1.1 mmol). Purification by chromatography on silica gel eluting with (2:1) hexanes/ethyl acetate afforded 0.36 g coupled product. Deprotection in boiling TFA (20 minutes), and purification by reverse phase HPLC and freeze drying afforded 0.2 g (64%) product. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.16 (m, 3H), 8.13 (dd, J = 6.9, 2.2 Hz, 1H), 7.61 (m, 5H), 7.45 (m, 1H), 7.33 (m, 7H), 4.45 (s, 2H), 4.14 (d, J = 5.9 Hz, 2H) ppm; ESI mass spectrum analysis m/z (relative intensity): 514.8 (M+H, 100); Elemental Analysis calc'd for C<sub>25</sub>H<sub>21</sub>F<sub>3</sub>N<sub>4</sub>O<sub>3</sub>S(TFA)1.3: C:50.02, H:3.39, N:8.45, found C:50.10, H:3.35, N:8.39.

15

**Example 211**

1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonylmethyl)pyrazole, trifluoroacetic acid salt

20 Part A. The pyrazole (1 g, 3.92 mmol) obtained in part B of Example 10 was dissolved in CCl<sub>4</sub>, then NBS (1.1 g, 6.27 mmol) and benzoylperoxide (0.038 g, 0.5 mmol) were added. The mixture was heated to reflux for 18 hr. After removal of the solvent, 50 mL water was added, then extracted with EtOAc, washed the organic layer with brine and dried over MgSO<sub>4</sub>. Filtration and concentration of the filtrate in vacuo was followed by purification using flash chromatography (2:3 / Hexane:Methylene chloride) to afford 0.55 g of the desired bromomethyl product as a light yellow solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.77-7.69 (m, 3H); 7.61 (t, J = 7.69, 1H); 7.13 (s, 1H); 4.51 (s, 2H); 4.32 (q, J = 6.95, 2H); 1.33 (t, J = 6.96, 3H) ppm; Ammonia CI mass spectrum analysis m/z (relative intensity): 334.0 (97) and 336.0 (100).

35 Part B. To the product of part A (0.55 g, 1.65 mmol) in DMF was added KSMc (0.16 g, 1.81 mmol). The mixture was heated to reflux over night. The solution was quenched with water (100 mL) and extracted with EtOAc. The organic layer was washed with

brine and dried over  $\text{MgSO}_4$ . Filtration, bubbling air through the filtrate for 2h. and concentration of the filtrate in vacuo was followed by purification using flash chromatography (3:2/Hex:EtOAc) to afford 0.14 g methylsulfonylmethyl compound as a colorless oil. Ammonia-Cl mass spectrum analysis  $m/z$  (relative intensity): 334.1 ( $M+H$ , 100).  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 7.77-7.69 (m, 4H); 7.61 (t,  $J = 8.05$ , 1H); 4.38 (s, 2H); 4.30 (q,  $J=6.96$ , 2H); 2.94 (s, 3H); 1.32 (t,  $J = 6.96$ , 3H) ppm.

Part C. Standard Weinreb coupling procedures of the product from part B with 2'-tert-butylaminosulfonyl-[1,1']-biphenyl aniline followed by the usual acid quench and silica gel flash chromatography afforded 0.13 g of the desired coupled product. ESI mass spectrum analysis  $m/z$  (relative intensity): 613.8 (75).  $^1\text{H NMR}(\text{CDCl}_3)$   $\delta$ : 8.35 (s, 1H); 8.16 (m, 1H); 7.82 (s, 1H); 7.75-7.55 (m, 8H); 7.50-7.45 (m, 2H); 7.30 (m, 1H); 7.16 (s, 1H); 4.42 (s, 2H); 3.00 (s, 3H); 1.02 (s, 9H) ppm.

Part D. To the product from part C (0.13 g, 0.22 mmol) dissolved in ethanol (50 mL) was added 10% Pd/C (20 mg) and 2 mL AcOH. Hydrogenation of this solution on the Parr at 50psi for 18h followed by filtration through a pad of Celite and concentration afforded a crude reduced product which was treated TFA (6 mL) and heated to reflux for 50 min. After removal of the solvent and purification via standard HPLC reverse phase techniques and lyophilization afforded the title compound as a colorless solid. ESI mass spectrum analysis  $m/z$  (relative intensity): 540.1 ( $M+H$ , 100).

#### Example 212

1-(3-amidino)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylaminosulfonylmethyl)pyrazole, trifluoroacetic acid salt.

Part A. To the product (1.1 g, 3.29 mmol) from part A (Example 211) in DMF was added  $\text{NaN}_3$  (0.24 g, 3.62 mmol). The mixture was stirred at R.T. for 18h. The reaction mixture was quenched with water (200 mL) and extracted with EtOAc. Washed the

organic layer with water and brine and dried over  $\text{MgSO}_4$ . The mixture was filtered and concentrated to afford 0.93 g of the crude azidomethyl compound. ESI mass spectrum analysis  $m/z$  (relative intensity): 297.1 ( $M+H$ , 100).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 7.77 (m, 3H); 7.59 (m, 1H); 7.08 (s, 1H); 4.44 (s, 2H); 4.30 (q,  $J = 7$ , 2H); 1.31 (t,  $J = 7$ , 3H) ppm.

Part B. To the product (0.54 g, 1.82 mmol) from part A in THF, was added  $\text{PPh}_3$  (0.53 g, 2.01 mmol). The reaction mixture was stirred at rt for 4h and the solvent was evaporated. HCl (1N, 50 mL) was added and the organics were extracted with EtOAc. The organic layer was washed with brine and dried over  $\text{MgSO}_4$ . Evaporation in vacuo afforded the desired aminomethyl compound (0.32 g) as a white solid. ESI mass spectrum analysis  $m/z$  (relative intensity): 271.1 ( $M+H$ , 100).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 7.77 (s, 1H); 7.72 (m, 2H); 7.59 (m, 1H); 7.01 (s, 1H); 4.30 (q,  $J = 7$ , 2H); 3.96 (s, 2H); 1.31 (t,  $J = 7$ , 3H) ppm.

Part C. To the product (0.43 g, 1.59 mmol) from part B in  $\text{CH}_2\text{Cl}_2$  was added triethylamine (1.5eq.). The reaction mixture was cooled to  $0^\circ\text{C}$  and  $\text{CH}_3\text{SO}_2\text{Cl}$  (1eq.) was added. The reaction mixture was stirred at R.T. for 18hr. diluted with  $\text{CH}_2\text{Cl}_2$  and washed with 1N HCl,  $\text{NaHCO}_3$  (sat.), brine, then dried over  $\text{MgSO}_4$ . Evaporation in vacuo was followed by purification via flash chromatography (4:1/Hex:EtOAc) to afford 0.42 g of the desired methylsulfonamide pyrazole precursor. Ammonia CI mass spectrum analysis  $m/z$  (relative intensity): 349.0 ( $M+H$ , 100).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 7.76 (m, 2H); 7.73 (m, 1H); 7.61 (m, 1H); 7.08 (s, 1H); 4.44 (d,  $J = 6.3$ , 2H); 4.29 (q,  $J = 7.3$ , 2H); 3.325 (s, 1H); 3.01 (s, 3H); 1.31 (t,  $J = 7.3$ , 3H) ppm.

Part D. Standard Weinreb coupling procedures of the product from part B with 2'-tert-butylaminosulfonyl-[1,1']-biphenylamine followed by the usual acid quench and silica gel flash chromatography afforded the desired coupled product. ESI (-ve) mass spectrum analysis  $m/z$  (relative intensity): 605.1 ( $M-H$ , 100).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.55 (s, 1H); 8.16 (m, 1H); 7.74

(m, 5H); 7.56 (m, 6H); 7.30 (m, 1H); 7.02 (s, 1H); 4.46 (d, 2H); 3.81 (s, 1H); 3.06 (s, 3H); 1.04 (s, 9H) ppm.

Part C. Standard Pinner-amidine reaction protocol followed by  
5 purification via reverse phase HPLC techniques and  
lyophilization afforded the desired compound as colorless  
crystals. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 10.69 (s, 1H); 9.43 (s, 2H); 9.15  
(s, 2H); 8.05 (m, 1H); 7.95 (s, 1H); 7.85 (m, 1H); 7.80 (m,  
1H); 7.70 (m, 4H); 7.60 (m, 2H); 7.35 (m, 2H); 7.30 (m, 1H);  
10 7.20 (m, 3H); 4.28 (d, J = 6.1, 2H); 2.97 (s, 3H) ppm. ESI mass  
spectrum analysis m/z (relative intensity): 568.0 (100) HRMS for  
C<sub>25</sub>H<sub>26</sub>N<sub>7</sub>O<sub>5</sub>S<sub>2</sub> 568.143686 (calcd); 568.145471 (obs).

#### Example 213

15 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-  
(methylaminosulfonylmethyl)pyrazole, trifluoroacetic acid salt

Part A: Standard Weinreb coupling of the product from part C  
20 (Example 203) with 4-bromo-2-fluoroaniline afforded the desired  
amide. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.13 (t, J=8.4, 1H); 7.90 (brd, 1H);  
7.79 (m, 1H); 7.78 (m, 2H); 7.61 (m, 1H); 7.35 (m, 2H); 6.96  
(s, 1H); 4.86 (m, 1H); 4.44 (d, J = 6.2, 2H); 3.04 (s, 3H) ppm.  
ESI (-ve) mass spectrum analysis m/z (relative intensity):  
25 489.8 (85) and 491.8 (100).

Part B: Standard Suzuki coupling of the product from part A  
with 2-thiomethylboronic acid afforded the desired 2'-  
thiomethyl-biphenyl intermediate. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.25 (brd,  
30 1H); 8.00 (brd, 1H); 7.83 (s, 1H); 7.75 (m, 2H); 7.62 (m, 1H);  
7.35 (m, 6H); 6.96 (s, 1H); 4.85 (m, 1H); 4.48 (d, J = 5.9,  
2H); 3.05 (s, 3H); 2.39 (s, 1H) ppm. ESI mass spectrum  
analysis m/z (relative intensity): 557.9 (M+Na, 100). ESI (-ve)  
mass spectrum analysis m/z (relative intensity): 533.8 (M-H,  
35 100).

Part C: To the product from part B (0.54 g, 1.01 mmol) in  
CH<sub>2</sub>Cl<sub>2</sub> was added MCPBA (0.52 g, 3.03 mmol) and the reaction

mixture was stirred at R.T. for overnight. The mixture was then CH<sub>2</sub>Cl<sub>2</sub> and washed with NaHCO<sub>3</sub> (sat.), sodium bisulfite, brine and dried over MgSO<sub>4</sub>. Filtration and concentration of the filtrate in vacuo was followed by purification using flash chromatography (1:1/Hex:EtOAc) to afford 0.53 g of the sulfonylmethyl derivative as white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 10.53 (s, 1H); 8.07 (m, 1H); 7.97 (s, 1H); 7.85 (m, 1H); 7.8 (m, 6H); 7.41 (m, 1H); 7.35 (m, 1H); 7.23 (m, 2H); 4.23 (s, 2H); 2.94 (s, 3H); 2.89 (s, 3H)ppm. ESI (-ve) mass spectrum analysis m/z (relative intensity): 565.8 (70).

Part D: The product from part C was hydrogenated as described previously to afford the desired benzylamine analog as colorless crystals following reverse phase HPLC and lyophilization techniques. <sup>1</sup>HNMR(DMSO) δ: 10.53 (s, 1H); 8.16 (brd, 2H); 8.07 (m, 1H); 7.75 (m, 1H); 7.72 (m, 4H); 7.49 (m, 5H); 7.21 (m, 2H); 4.23 (d, J=6.2, 2H); 4.09 (m, 2H); 2.93 (s, 3H); 2.90 (s, 3H)ppm. ESI mass spectrum analysis m/z (relative intensity): 571.9 (M+H, 100). HRMS calc'd for C<sub>26</sub>H<sub>27</sub>N<sub>5</sub>O<sub>5</sub>FS<sub>2</sub> 572.143766 (calcd); 572.145154 (obs).

#### Example 214

1-(3-(N-carboxymethyl)amidinophenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

To the Example 92 compound (100 mg, 0.19 mmol) in DMF was added methylchloroformate (36 mg, 0.38 mmol) and Et<sub>3</sub>N. The mixture was stirred at R.T. for 2.5hr. Diluted with 100 mL water and extracted with EtOAc, the organic layer was washed with water, brine and dried over MgSO<sub>4</sub>, filtered and concentrated in vacuo and purified using reverse phase HPLC techniques to afford the the desired carbamate [ESI mass spectrum analysis m/z (relative intensity): 590.9 (100)], which was then treated with TFA and heated to gentle reflux for 0.5h. Evaporation of the TFA followed by purification via HPLC reverse phase and lyophilization afforded the title compound. <sup>1</sup>HNMR(DMSO) δ: 11.34 (s, 1H); 8.61 (s, 2H); 8.01 (m, 1H); 7.95

(m, 1H): 7.80 (m, 1H); 7.69 (m, 1H); 7.64 (m, 3H); 7.49 (s, 2H); 7.40 (m, 1H); 7.03 (s, 1H); 3.79 (s, 3H); 2.28 (s, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity): 535.0 (M+H, 100). HRMS calc'd for C<sub>24</sub>H<sub>22</sub>N<sub>8</sub>O<sub>3</sub>S 535.151213 (calcd);  
5 535.151600 (obs).

#### Example 215

1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt  
10

Part A: Standard Weinreb coupling of 2'sulfonylmethyl-biphenylamine to the pyrazole ester obtained in part B of Example 10 followed by standard workup afforded after silica gel purification the desired coupled amide precursor.  
15 <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.24 (d, J=7.7, 1H); 7.87 (s, 1H); 7.81 (s, 1H); 7.76 (m, 1H); 7.69 (m, 6H); 7.45 (m, 2H); 7.35 (m, 1H); 6.71 (s, 1H); 2.68 (s, 3H); 2.42 (s, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity): 478.9 (M+Na, 100). ESI (-ve) mass spectrum analysis m/z (relative intensity): 454.9 (M-H, 20 100).

Part B: To the product from from part A( 0.48 g, 1.05 mmol) in EtOH was added 10% Pd/C (80 mg) and 1 mL TFA. The mixture was hydrogenated on a Parr apparatus at 50psi for 18hr. After  
25 filtration through a pad of Celite and concentration the filtrate in vacuo, purified using reversed phase prep HPLC to afford the title compound. <sup>1</sup>HNMR(DMSO) δ: 10.65 (s, 1H); 8.17 (brd, 2H); 8.06 (d, J=7.7, 1H); 7.75 (m, 5H); 7.49 (m, 6H); 6.92 (s, 1H); 4.10 (m, 2H); 2.81 (s, 3H); 2.29 (s, 3H) ppm. ESI  
30 mass spectrum analysis m/z (relative intensity): 460.9 (M+H, 100). HRMS calc'd for C<sub>25</sub>H<sub>25</sub>N<sub>4</sub>O<sub>3</sub>S 461.164738 (calcd); 461.164405 (obs).

#### Example 216

35 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-methyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl pyrazole.  
trifluoroacetic acid salt



Part A: Standard Weinreb coupling of 2'-tert-butylaminosulfonyl-2-methyl-biphenylamine with the previously obtained pyrazole ester afforded the desired coupled amide precursor. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.17 (d, J = 7.7, 1H); 7.83 (m, 4H); 7.64 (m, 2H); 7.56 (m, 2H); 7.4 (m, 3H); 7.15 (s, 1H); 3.61 (s, 1H); 2.36 (s, 3H); 1.04 (s, 9H) ppm. ESI mass spectrum analysis m/z (relative intensity): 604.1 (M+Na, 100). ESI (-ve) mass spectrum analysis m/z (relative intensity): 580.3 (M-H, 100).

Part B: Reduction of the benzonitrile to the benzylamine followed by removal of the tert-butyl group and standard reverse phase HPLC purification afforded the title compound. <sup>1</sup>HNMR(DMSO) δ: 10.33 (s, 1H); 8.23 (bd, 2H); 8.02 (m, 1H); 7.76 (s, 1H); 7.66 (m, 6H); 7.40 (d, J = 8.1, 1H); 7.31 (m, 5H); 4.15 (m, 2H); 2.25 (s, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity): 530.2 (M+H, 100).

#### Example 217

1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole. trifluoroacetic acid salt

Standard Weinreb coupling of 4-bromo-2-fluoro-aniline with the previously obtained 1,2,3-triazole-5-carboxylic acid as used in the preparation of Example 46 afforded after purification via flash silica-gel chromatography the coupled amide triazole derivative. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.23 (s, 1H); 8.11 (m, 1H); 7.86 (m, 4H); 7.68 (m, 1H); 7.34 (m, 2H) ppm. ESI (-ve) mass spectrum analysis m/z (relative intensity): 383.8 (100) and 385.7 (80). Standard Suzuki coupling of this intermediate with 2-thiomethyl boronic acid followed by oxidation with MCPBA in dichloromethane afforded the desired biphenylsulfonyl derivative. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.34 (m, 3H); 8.05 (bd, 1H); 7.93 (m, 3H); 7.74 (m, 3H); 7.37 (m, 2H); 7.24 (m, 1H); 2.74 (s, 3H) ppm. ESI (-ve) mass spectrum analysis m/z (relative intensity): 459.9 (M-H, 100). This intermediate was then reduced to the benzylamine and purified via standard

conditions described previously.  $^1\text{H NMR}$  (DMSO)  $\delta$ : 10.76 (s, 1H); 8.53 (s, 1H); 8.21 (bd, 2H); 8.05 (m, 1H); 7.77 (m, 7H); 7.39 (m, 2H); 7.22 (m, 1H); 4.14 (m, 2H); 2.89 (s, 3H) ppm. ESI (-ve) mass spectrum analysis  $m/z$  (relative intensity): 465.9 (M+H, 100). HRMS calc'd for  $\text{C}_{23}\text{H}_{21}\text{N}_5\text{O}_3\text{FS}$ , 466.134915, found 466.136900.

#### Example 218

1-(3-aminomethyl-4-methyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole. trifluoroacetic acid salt

Part A: To a cold ( $0^\circ\text{C}$ ) acidic (Con HCl, 100 mL) solution of 2-methyl-4-amino-benzonitrile (10 g, 78.12 mmol) was added sodium nitrite (8.08 g, 117.19 mmol) previously dissolved in water (20 mL). The reaction temperature was kept cold throughout the addition of sodium nitrite. After stirring for an additional 0.5h a solution of  $\text{SnCl}_2$  in con HCl (50 mL) was added dropwise. A precipitate immediately ensued. The reaction mixture was allowed to stir at  $0^\circ\text{C}$  for an additional 18h then filtered, washed with cold water (1000 mL) followed by a solution of Petroleum ether/ether (2:1, 500 mL). The residue was dried in vacuo overnight to afford a total weight of 8.15 g crude hydrazine tin salt.

Part B: The tin salt obtained in part A was stirred in glacial acetic acid (100 mL). To this solution was then added methoxyoxime derived from ethyl 2,4-dioxovalerate (4.59 g, 24.55 mmol). The reaction mixture was gently refluxed overnight. Acetic acid was evaporated off and the residue was then quenched with water (200 mL). The organics were extracted with ethyl acetate (2X100 mL) washed with saturated sodium bicarbonate (2X50 mL), brine (50 mL) and dried (magnesium sulfate). Column chromatography (silica gel, ethyl acetate:hexane 2:8) then afforded the desired pyrazole carboxylate (4 g) as a pale yellow oil which crystallized on standing.

Part C: The product from part B was then subjected to the standard Weinreb trimethylaluminum coupling protocol with 2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl-amine as described previously. The crude was then subjected to silica gel column chromatography (methylene chloride:methanol, 9:1) to afford pure material in 90% yield. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.30 (bs, 1H), 8.13 (bd, 1H), 7.78-7.23 (m, 10H), 6.78 (s, 1H), 3.68 (s, 1H), 2.60 (s, 3H), 2.40 (s, 3H), 1.01 (s, 9H)ppm; ESI mass spectrum analysis m/z (relative intensity) 528 (M+H, 100).

Part D: The product from part D was then subjected reduction (Parr apparatus) at 50psi hydrogen pressure in an acidic media (methanol, acetic acid) using 10% palladium on carbon overnight. The solvents were evaporated and the crude was then stirred in TFA (reflux) for 0.5h. Evaporation of the solvents then afforded crude benzylamine product which was then subjected to a preparative HPLC purification technique (acetonitrile:water, gradient containing 5% TFA) to afford the desired benzylamine as flaky colorless crystals. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.6 (s, 3H), 8.14 (bs, 2H), 8.01 (d, 1H), 7.68 (d, 2H), 7.64-7.54 (m, 2H), 7.38-7.26 (m, 5H), 6.91 (s, 1H), 4.07 (bd, 2H), 2.38 (s, 3H), 2.33 (s, 3H)ppm; ESI mass spectrum analysis m/z (relative intensity) 476.2 (M+H, 100).

#### Example 219

1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

The pyrazole compound was prepared from readily accessible 4-fluoro-3-cyano-phenylhydrazine.tin salt (obtained from the corresponding aniline) and the oxime derived from ethyl-2,4-dioxovalerate via procedures described previously. Standard Weinreb coupling of the pyrazole with 2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl-amine afforded the desired coupled amide precursor which was then subjected to the standard reductive protocol (50psi hydrogen pressure, methanol:acetic acid) using 10% palladium on carbon.

Evaporation of the solvents followed by treatment with TFA for 0.5h and then preparative HPLC as described before afforded the title compound as colorless crystals. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.25 (bs, 3H), 8.00 (d, 1H), 7.78-7.23 (cp, 12H), 6.95 (s, 1H), 4.14 (m, 2H), 2.30 (s, 3H) ppm. -ESI mass spectrum analysis m/z(relative intensity) 480.2 (M+H, 100).

#### Example 220

1-(3-aminomethyl-4-chloro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

The pyrazole compound was prepared from readily accessible 4-chloro-3-cyano-phenylhydrazine tin salt (obtained from the corresponding aniline) and the oxime derived from ethyl-2,4-dioxovalerate via procedures described previously. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.78 (d, 1H), 7.86-7.55 (m, 2H), 6.86 (s, 1H), 4.24 (q, 2H), 2.35 (s, 3H), 1.28 (t, 3H) ppm; ESI mass spectrum analysis m/z(relative intensity) 290 (M+H, 100). Standard Weinreb coupling of the pyrazole obtained above with 2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl-amine afforded the desired coupled amide precursor which was then subjected to the treatment with tetrabutylammonium borohydride (1.5equiv.) in dichloromethane for 24h. The solvent was evaporated and the residue was then gently refluxed in TFA for 0.5h. Evaporation of the solvent followed by preparative HPLC as described before afforded the title compound as colorless crystals: <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.25 (bs, 3H), 8.00 (d, 1H), 7.78-7.23 (cp, 12H), 6.95 (s, 1H), 4.14 (m, 2H), 2.30 (s, 3H) ppm. ESI mass spectrum analysis m/z(relative intensity) 497.1 (M+H, 100).

#### Example 221

1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethylpyrazole, trifluoroacetic acid salt

The reduction of the benzonitrile precursor prepared as described before (10% palladium on carbon, methanol/acetic acid

at 50psi hydrogen) afforded the title compound as colorless crystals after preparative HPLC purification and lyophilization techniques. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.68 (s, 1H), 8.27 (bs, 2H), 8.02 (dd, 1H), 7.81 (m, 1H), 7.73 (s, 1H), 7.68-7.60 (m, 4H), 7.61-7.43 (m, 3H), 7.38-7.30 (m, 2H), 7.20 (dd, 2H), 4.18 (bd, 2H)ppm; ESI mass spectrum analysis m/z(relative intensity) 551.9 (M+H, 100).

#### Example 222

10 1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole, trifluoroacetic acid salt

Part A: The coupling of ethyl 1-(3-cyanophenyl)-3-methyl-5-carboxylate with 2'-tert-butylaminosulfonyl-3-fluoro-[1,1']-biphen-4-yl-amine via the Weinreb protocol as described previously afforded the desired coupled amide compound. In this case 1.5 equivalents of the biphenyl analog was used to facilitate the coupling. Purification via silicagel(methylene chloride/methanol, 9/1) afforded pure amide (60%) as a pale yellow oil. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.35 (t, 1H), 8.15 (dd, 1H), 7.96 (m, 1H), 7.82 (s, 1H), 7.78-7.68 (m, 4H), 7.60-7.48 (m, 4H), 7.20 (m, 1H), 6.74 (s, 1H), 3.67 (s, 1H), 2.04 (s, 3H), 1.04 (s, 9H) ppm; ESI mass spectrum analysis m/z(relative intensity) 553.9 (M+Na, 100). ESI (-ve) mass spectrum analysis m/z(relative intensity) 529.9 (M-H, 100).

Part B: The product obtained from part A was then converted to the corresponding benzylamine via the reductive methodology (10% Pd/C, MeOH/AcOH at 50psi hydrogen pressure) described previously. Evaporation of the solvent followed by standard removal of the tert-butyl group with TFA and purification via preparative HPLC techniques afforded pure title compound as colorless crystals (60%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.42 (s, 1H), 8.20 (bs, 2H), 8.02 (dd, 1H), 7.70-7.59 (m, 4H), 7.55-7.29 (m, 6H), 7.19 (dd, 1H), 6.97 (s, 1H), 4.11 (bd, 2H), 2.50 (s, 2H)ppm; ESI mass spectrum analysis m/z(relative intensity) 480 (M+H, 100).

## Example 223

1-(3-aminomethyl)phenyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-  
biphen-4-yl)aminocarbonyl]-3-trifluoromethylpyrazole,  
trifluoroacetic acid salt

Part A: The coupling of ethyl 1-(3-cyanophenyl)-3-methyl-5-carboxylate with 2'-methylsulfonyl-3-fluoro-[1,1']-biphen-4-yl-amine (previously prepared via the Suzuki coupling methodology of 2-thiomethylphenylboronic acid with 4-bromo-2-fluoro aniline) via the Weinreb protocol as described previously afforded the desired coupled amide compound. Purification via silica gel (methylene chloride/methanol, 9/1) afforded pure amide (80%) as a colorless solid. The amide was also obtained by first coupling (Weinreb) of 2-fluoro-4-bromo-aniline with the above pyrazole carboxylate followed by Suzuki coupling with 2-thiomethyl-phenylboronic acid and oxidation to the sulfonyl derivative. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.39 (t, 1H), 8.20 (dd, 1H), 7.96 (bd, 1H), 7.83 (s, 1H), 7.78-7.59 (m, 5H), 7.41-7.35 (t, 2H), 7.17 (d, 1H), 6.74 (s, 1H), 2.73 (s, 3H), 2.40 (s, 3H) ppm; ESI mass spectrum analysis m/z(relative intensity) 593 (M+Na, 100). ESI (-ve) mass spectrum analysis m/z(relative intensity) 572 (M-H, 100).

Part B: Reduction of the product from part A via procedures described previously and HPLC purification afforded the desired compound as colorless crystals (70%). <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.45 (s, 1H), 8.20 (bs, 3H), 8.08 (dd, 1H), 7.80-7.66 (m, 4H), 7.55-7.37 (m, 5H), 7.21 (dd, 1H), 6.98 (s, 1H), 4.12 (s, 2H), 2.94 (s, 3H), 2.50 (s, 3H) ppm; ESI mass spectrum analysis m/z(relative intensity) 479 (M+H, 100).

## Example 224

1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole bis-trifluoroacetate

Part A. Preparation of N-(3-fluoro-4-nitrophenyl)morpholine.

3,4-Difluoronitrobenzene (10.0 g, 62.86 mmol) was dripped into a cooled solution (0°C) of morpholine (6.03 mL, 69.14 mmol), diisopropylamine (11.83 mL, 67.89 mmol) and 35 mL ethyl acetate over 1.5H. The reaction mixture was allowed to warm to ambient temperature over 48H. Diluted reaction mixture with 25 mL methylene chloride, 100 mL ethyl acetate and 50 mL water. Separated and extracted aqueous 2x25 mL EtOAc. Combined organics, dried over magnesium sulfate and concentrated under reduced pressure to give a yellow solid. The crude material was recrystallized from acetone and water to give 12.55 g of a yellow crystalline solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 7.99 (m, 2H) 7.14 (t, 1H, J = 8.79 Hz) 3.71 (bt, 4H, J = 4.56 Hz) 3.23 (bt, 4H, J = 4.76 Hz). ESI mass spectrum analysis m/z (relative intensity) 227 (M+H).

Part B. Preparation of N-(3-fluoro-4-aminophenyl)morpholine.

N-(3-fluoro-4-nitrophenyl)morpholine (6.01 g, 26.59 mmol) and a catalytic amount of palladium on carbon (10%) were suspended in 100 mL methanol in a Parr flask. The reaction mixture was placed on the Parr Hydrogenator at 60 psi for 2H. The reaction mixture was passed through a Celite pad and the filtrate was concentrated under reduced pressure to give 4.50 g of N-(3-fluoro-4-aminophenyl)morpholine an off-colored solid. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 6.73 (t, 1H, J = 9.34), 6.28 (m, 2H), 3.64 (bt, 4H, J = 4.58 Hz), 2.76 (bt, 4H, J = 4.58 Hz). ESI mass spectrum analysis m/z (relative intensity) 197 (M+H, 100). <sup>19</sup>F NMR (DMSO-d<sub>6</sub>) δ: -124.455.

Part C. Preparation of 1-(3-cyanophenyl)-3-methyl-5-((3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

Dimethylaminopyridine (0.28 g, 2.25 mmol) was added to a solution of 1-(3-cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid chloride (0.46 g, 1.88 mmol) and N-(3-fluoro-4-aminophenyl)morpholine (0.37 g, 1.88 mmol) in 20 mL methylene chloride. The reaction mixture was stirred at ambient temperature for 72H and then concentrated under reduced

pressure. The resulting residue was purified via flash chromatography to give 0.070 g of pure 1-(3-cyanophenyl)-3-methyl-5-((3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ:

5 10.50 (s, 1H), 7.93 (s, 1H), 7.83 (d, 1H, J = 7.33Hz), 7.73 (d, 1H, J = 8.79Hz), 7.62 (t, 1H, J = 7.87Hz), 7.53 (m, 1H), 7.34 (d, 1H, J = 9.15Hz), 6.99 (t, 1H, J = 9.34Hz), 6.93 (s, 1H), 3.69 (bt, 4H, J = 4.58Hz), 2.92 (bt, 4H, J = 4.58Hz), 2.28 (s, 3H). ESI mass spectrum analysis m/z(relative intensity) 406 (M+H, 100), 833 (2M+Na). <sup>19</sup>F NMR (dmsO-d<sub>6</sub>, 300MHz) δ: -122.081.

10

Part D. Preparation of 1-(3-amidinophenyl)-3-methyl-5-((3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

15 1-(3-Cyanophenyl)-3-methyl-5-((3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole (0.070 g, 0.173 mmol) was dissolved in 2 mL chloroform and 2 mL ethanol at 0°C. Hydrogen chloride gas was bubbled into the reaction mixture for 1H. The reaction mixture was allowed to warm to ambient

20 temperature over 15H and was concentrated under reduced pressure. The resulting solid was placed under high vacuum for 2H. Then the crude imidate was dissolved in 2 mL ethanol and ammonium carbonate (0.25 g, 2.60 mmol) was added to the solution at ambient temperature. The reaction mixture was stirred for

25 72H and concentrated under reduced pressure. The crude product was purified by standard HPLC methods to give 0.016 g of pure 1-(3-amidinophenyl)-3-methyl-5-((3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ:

30 10.53 (s, 1H), 9.40 (s, 2H), 9.12 (s, 2H), 7.93 (d, 1H, J = 1.71Hz, 2H), 7.81 (m, 1H), 7.70 (m, 2H), 7.53 (dd, 1H, J = 15Hz), 7.35 (d, 1H, J = 8.79Hz), 7.01 (m, 2H), 3.72 (bt, 4H, J = 4.52Hz), 2.95 (bt, 4H, J = 4.6Hz), 2.29 (s, 3H). ESI mass spectrum analysis m/z(relative intensity) 423 (M+H, 100). <sup>19</sup>F NMR(DMSO-d<sub>6</sub>) δ: -73.790 and -121.040. HRMS Calculated for

35 C<sub>22</sub>H<sub>24</sub>N<sub>6</sub>O<sub>2</sub>F<sub>1</sub>: 423.194478, found 423.192755.



## Example 225

**1-(3-aminomethylphenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole bis-trifluoroacetate**

- 5 Part A. Preparation of 1-(3-cyanophenyl)-3-methyl-5-((3'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

Dimethylaminopyridine(0.18 g,1.47 mmol) was added to a solution of N-(cyanophenyl)-3-methyl-pyrazole-5-carboxylic acid  
10 chloride(0.30 g,1.22 mmol) and previously described N-(3-fluoro-4-aminophenyl)morpholine(0.24 g,1.22 mmol) in 20 mL methylene chloride. The reaction mixture was stirred at ambient temperature for 72H and then concentrated under reduced pressure. The resulting residue was purified via flash  
15 chromatography to give 0.070 g of pure 1-(3-cyanophenyl)-3-methyl-5-((3'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.50 (s,1H), 7.93 (s,1H), 7.83 (d, 1H, J = 7.33Hz), 7.73 (d, 1H, J = 8.79Hz), 7.62 (t, 1H, J = 7.87Hz), 7.53 (m,1H), 7.34  
20 (d, 1H, J = 9.15Hz), 6.99 (t, 1H, J = 9.34Hz), 6.93 (s,1H), 3.69 (bt, 4H, J = 4.58Hz), 2.92 (bt, 4H, J = 4.58Hz), 2.28 (s,3H). ESI mass spectrum analysis m/z(relative intensity) 406 (M+H, 100) 833 (2M+Na). <sup>19</sup>F NMR (DMSO-d<sub>6</sub>) δ: -122.078.

- 25 Part B. Preparation of 1-(3-aminomethylphenyl)-3-methyl-5-((3'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)-pyrazole.

1-(3-Cyanophenyl)-3-methyl-5-((3'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole(0.21 g,0.519 mmol) was  
30 suspended with a catalytic amount of palladium on carbon(10%) in 15 mL methanol and 1 mL trifluoroacetic acid. The reaction mixture was placed on the Parr Hydrogenator at 60psi for 20H. The reaction mixture was passed through a Celite pad and the filtrate was concentrated under reduced pressure. The crude  
35 material was purified by standard HPLC methods to give pure 1-(3-aminomethylphenyl)-3-methyl-5-((3'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.53 (s,1H), 8.18 (bs,2H), 7.60 (s,1H), 7.53 (dd, 1H, J<sub>1</sub> =

15.0Hz,  $J_2 = 2.2\text{Hz}$ ), 7.44 (m, 2H), 7.33 (d, 2H,  $J = 7.33\text{Hz}$ ), 6.98 (dd, 1H,  $J_1 = 9.3\text{Hz}$ ,  $J_2 = 9.2\text{Hz}$ ), 6.86 (s, 1H) 4.07 (bt, 2H,  $J_1 = 2.9\text{Hz}$ ,  $J_2 = 2.6\text{Hz}$ ), 3.69 (bt, 4H,  $J_1 = 4.4\text{Hz}$ ,  $J_2 = 4.8\text{Hz}$ ), 2.91 (bt, 4H,  $J_1 = 4.9\text{Hz}$ ,  $J_2 = 4.8\text{Hz}$ ), 2.47 (s, 3H). ESI mass spectrum analysis m/z (relative intensity) 410 (M+H, 100).  $^{19}\text{F}$  NMR (DMSO-d<sub>6</sub>)  $\delta$ : -74.991 and -122.105. HRMS calculated for C<sub>22</sub>H<sub>25</sub>N<sub>5</sub>O<sub>2</sub>F: 410.199224, found 410.197598.

#### Example 226

10 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-((3-fluoro-4-(2-methylimidazol-1-yl)phenyl)aminocarbonyl)pyrazole bis-trifluoroacetate

Part A. Preparation of 3-fluoro-4-(2-methylimidazol-1-yl)nitro-benzene.

2-Methylimidazole (1.0 g, 12.18 mmol) was added to a solution of 3,4-difluoronitrobenzene in 100 mL DMF. Added potassium carbonate (2.02 g, 14.61 mmol) to the reaction mixture and stirred vigorously for 24H. Concentrated reaction mixture under reduced pressure and took up residue in 100 mL ethyl acetate. Washed organics 6x50 mL water and 3x50 mL brine solution. Dried resulting organics over magnesium sulfate and concentrated resulting organics under reduced pressure to give 1.66 g of crude 3-fluoro-4-N-(2-methylimidazol-1-yl)nitro-benzene.  $^1\text{H}$ NMR (dmsO-d<sub>6</sub>, 300MHz)  $\delta$ : 8.42 (dd, 1H,  $J_1 = 2.4\text{Hz}$ ,  $J_2 = 10\text{Hz}$ ), 8.21 (m, 1H), 7.86 (t, 1H,  $J = 8.4$ ), 7.34 (s, 1H), 6.98 (s, 1H), 2.21 (s, 1H). ESI mass spectrum analysis m/z (relative intensity) 221.9 (M+H, 100).  $^{19}\text{F}$  NMR (DMSO-d<sub>6</sub>)  $\delta$ : -118.512.

Part B. Preparation of 3-fluoro-4-(2-methylimidazol-1-yl)aniline.

3-Fluoro-4-N-(2-methylimidazol-1-yl)nitrobenzene (1.66 g, 7.51 mmol) was added to a suspension of palladium on carbon (10%) in 30 mL menthanol. The reaction mixture was placed on the Parr Hydrogenator at 60psi for 20H. Filtered reaction mixture through a Celite pad and concentrated filtrate

under reduced pressure to give 1.40 g of the crude 3-fluoro-4-N-(2-methylimidazol-1-yl)aniline. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>, 300MHz) δ:

7.02 (m, 2H), 6.83 (s, 1H), 6.43 (m, 2H), 5.70 (bs, 1H), 2.07 (s, 3H). ESI mass spectrum analysis m/z (relative intensity).

5 <sup>19</sup>F NMR (DMSO-d<sub>6</sub>) δ: -124.344.

Part C. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-((3'-fluoro-4'-(2-methylimidazol-1-yl)phenyl)-aminocarbonyl)pyrazole.

10

Dimethylaminopyridine (0.19 g, 1.56 mmol) was added to a solution of N-(3-cyanophenyl)-3-trifluoromethyl-pyrazole-5-carboxylic acid chloride (0.39 g, 1.30 mmol) and 3-fluoro-4-(2-methylimidazol-1-yl)aniline (0.25 g, 1.30 mmol) in 10 mL methylene chloride. The reaction mixture was stirred at ambient temperature for 5H and then concentrated under reduced pressure. The resulting residue was purified via flash chromatography to give 0.16 g of pure 1-(3-cyanophenyl)-3-trifluoromethyl-5-((3'-fluoro-4'-(2-methylimidazol-1-yl)phenyl)-aminocarbonyl)pyrazole. ESI mass spectrum analysis m/z (relative intensity) 455.2 (M+H, 100).

15

20

Part D. Preparation of 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-((3'-fluoro-4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl)pyrazole bis-trifluoroacetate.

25

Standard transformation of the benzonitrile (0.16 g, 0.344 mmol) obtained in part C to the benzylamine via catalytic hydrogenation yielded 0.050 g 1-(3-methylaminophenyl)-3-trifluoromethyl-5-((3'-fluoro-4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl)pyrazole bis-trifluoroacetate after HPLC purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 11.25 (s, 1H), 7.91-7.52 (m, 10H), 4.12 (m, 2H), 2.43 (s, 3H). ESI mass spectrum analysis m/z (relative intensity) 459.1 (M+H, 100). HRMS(NH<sub>3</sub>-CI):

30

35

Calculated for C<sub>22</sub>H<sub>19</sub>N<sub>6</sub>O<sub>4</sub>F<sub>4</sub>: 459.155647, found 459.154688.

## Example 227

## 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole

- 5 Part A. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole.

Diethylazodicarboxylate (0.41 mL, 2.59 mmol) was dripped into a solution of previously described 1-(3-cyanophenyl)-3-trifluoromethyl-5-hydroxymethylpyrazole (0.46 g, 1.73 mmol), 4-hydroxy-[1,1']-biphenyl (0.44 g, 2.59 mmol), and triphenylphosphine (0.68 g, 2.59 mmol) in 15 mL THF over 1h. Let reaction mixture stir for 48h. Diluted reaction mixture with 30 mL water and extracted with ethyl acetate. Combined organics, dried over magnesium sulfate and purified crude material by flash chromatography to give 0.040 g of pure 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl) pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.01 (m, 1H), 7.91 (m, 1H), 7.75 (m, 1H), 7.58 (m, 4H), 7.44 (m, 2H), 7.34 (m, 1H), 6.99 (m, 2H), 6.88 (s, 1H), 5.05 (s, 2H). ESI mass spectrum analysis m/z (relative intensity) 420 (M+H, 100), 437 (M+NH<sub>4</sub>, 63).

## Examples 228 and 229

- 1-(3-amidinophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole (Example 228) and 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole (Example 229)

- Part A. Preparation of 1-(3-amidinophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole and 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole.

- Standard Pinner-amidine transformation of the 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole as previously described afforded 0.022 g of 1-(3-amidinophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl)pyrazole trifluoroacetate. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.42

(bs, 2H), 9.16 (bs, 2H), 8.06 (s, 1H), 8.01 (d, 1H, J = 8.1Hz), 7.91 (d, 1H, J = 8.1Hz), 7.79 (t, 1H, J = 8.1Hz), 7.56 (m, 4H), 7.39 (m, 2H), 7.28 (m, 1H), 7.23 (m, 1H), 7.01 (d, 2H, J = 8.79Hz), 5.26 (s, 2H). ESI mass spectrum analysis m/z (relative intensity) 437 (M+H, 100). HRMS(NH<sub>3</sub>-CI): Calculated for C<sub>24</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub>: 437.158921, found 437.157809 and 0.002 g of 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethylpyrazole after HPLC purification. <sup>1</sup>HNMR(dmsod<sub>6</sub>, 300MHz) δ: 8.18 (s, 1H) 7.99 (d, 1H, J = 7.7Hz), 7.78 (d, 1H, J = 9.2Hz), 7.68-7.53 (m, 5H), 7.39 (t, 2H, J = 7.7Hz), 7.27 (dd, 2H, J<sub>1</sub> = 7.3Hz, J<sub>2</sub> = 7.0Hz), 7.18 (s, 1H), 7.01 (d, 2H, J = 8.8Hz), 5.21 (s, 2H). ESI mass spectrum analysis m/z (relative intensity) 479 (M+H+MeCN). HRMS(NH<sub>3</sub>-CI): Calculated for C<sub>24</sub>H<sub>20</sub>N<sub>4</sub>O<sub>3</sub>: 437.15892, found 437.157809.

#### Examples 230 and 231

1-(3-amidinophenyl)-3-trifluoromethyl-5-((2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole bis-trifluoro acetate and 1-(3-Carboxamidophenyl)-3-trifluoromethyl-5-((2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole

#### Part A. Preparation of 2-fluoro-4-(-N-morpholino)aniline.

Morpholine (10.0 mL, 115 mmol) was added to a mixture of 4-bromo-2-fluoroaniline (1.03 g, 5.42 mmol), copper(I) bromide (0.039 g, 0.27 mmol) and potassium carbonate (1.50 g, 10.84 mmol). The reaction mixture was heated to 130°C for 48h, concentrated under reduced pressure and purified by flash chromatography to afford 0.11 g of pure 2-fluoro-4-(-N-morpholino)aniline. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 6.73 (dd, 1H, J<sub>1</sub> = 8.8Hz, J<sub>2</sub> = 9.9Hz), 6.64 (dd, 1H, J<sub>1</sub> = 2.6Hz, J<sub>2</sub> = 13.2Hz), 6.57 (m, 1H), 3.85 (m, 4H), 3.02 (m, 4H). ESI mass spectrum analysis m/z (relative intensity) 197 (M+H, 100). <sup>19</sup>FNMR (dmsod<sub>6</sub>, 300MHz) δ: -133.

Part B. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-((2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

2-Fluoro-4-(N-morpholino)aniline (0.11 g, 0.56 mmol) in 5 mL methylene chloride was dripped into a stirring solution of N-(3-cyanophenyl)-3-trifluoromethyl-pyrazole-5-carboxylic acid chloride (0.17 g, 0.56 mmol) and dimethylaminopyridine (0.082 g, 0.67 mmol) in 10 mL methylene chloride. The reaction mixture was stirred at ambient temperature for 20h. The reaction mixture was concentrated under reduced pressure and purified by flash chromatography to give 0.19 g of pure 1-(3-cyanophenyl)-3-trifluoromethyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl]pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 7.94 (m, 1H), 7.86 (s, 1H), 7.77 (m, 3H), 7.61 (dd, 2H, J<sub>1</sub> = 7.7Hz, J<sub>2</sub> = 8.1Hz), 7.12 (s, 1H), 3.85 (m, 4H), 3.14 (m, 4H).

Part C. Preparation of 1-(3-amidinophenyl)-3-trifluoro-methyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)amino-carbonyl]pyrazole bis-trifluoroacetate and 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl]pyrazole.

Standard Pinner-amidine transformation of the 1-(3-cyanophenyl)-3-trifluoromethyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl]pyrazole afforded 0.10 g of 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl]pyrazole bis-trifluoroacetate. <sup>1</sup>HNMR(dmso-d<sub>6</sub>, 300MHz) δ: 10.35 (s, 1H), 9.40 (bs, 2H), 9.11 (bs, 2H), 7.96 (s, 1H), 7.87 (t, 2H), 7.72 (t, 1H), 7.67 (s, 1H), 7.27 (t, 1H), 6.84-6.71 (m, 2H), 3.70-3.66 (m, 4H), 3.09-3.06 (m, 4H). ESI mass spectrum analysis m/z (relative intensity) 476.5 (M+H, 100). HRMS(CI): Calculated for C<sub>22</sub>H<sub>21</sub>N<sub>6</sub>O<sub>2</sub>F<sub>4</sub> 477.166212, found 477.166415. <sup>19</sup>F NMR (dmso-d<sub>6</sub>, 300MHz) δ: -61.354, -74.772 and 0.002 g of 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-[(2'-fluoro-4'-(N-morpholino)phenyl)aminocarbonyl]pyrazole after HPLC purification. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.31 (s, 1H), 8.10 (s, 1H), 7.98-7.94 (m, 2H), 7.64-7.50 (m, 2H), 7.33-7.27 (m, 1H), 6.83-6.70 (m, 2H), 3.70-3.66 (m, 4H), 3.09-3.06 (m, 4H). ESI mass spectrum analysis m/z (relative intensity) 477.5 (M+H, 100). <sup>19</sup>F NMR

(DMSO-d<sub>6</sub>) δ: -61.274, -74.363. HRMS(CI): Calculated for C<sub>22</sub>H<sub>20</sub>N<sub>5</sub>O<sub>3</sub>F<sub>4</sub> 478.150228, found 478.147507.

### Example 232

5 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-((3-trifluoromethyl-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole bis-trifluoroacetate

Part A. Preparation of 3-trifluoromethyl-4-N-  
10 morpholinoaniline.

3-Trifluoromethyl-4-N-morpholinonitrobenzene (1.0 g, 3.62 mmol) was added to a suspension of palladium on carbon (10%) in 25 mL methanol. The reaction mixture was placed under 1 atmosphere H<sub>2</sub> at ambient temperature for 24h. Passed reaction mixture through a Celite pad and concentrated filtrate under reduced pressure to give the desired aniline in quantitative yeild. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 7.20 (d, 1H, J=7.2Hz), 6.92 (d, 1H, J = 2.6), 6.81 (dd, 1H, J<sub>1</sub> = 2.9Hz, J<sub>2</sub> = 8.4Hz), 3.80 (m, 4H), 3.74 (bs, 2H), 2.83 (bt, 4H, J = 4.4Hz), 3.70-3.66 (m, 4H), 3.09-3.06 (m, 4H). Ammonia CI mass spectrum analysis m/z (relative intensity) 247 (M+H, 100).

Part B. Preparation of 1-(3-cyanophenyl)-3-trifluoromethyl-5-  
25 ((3'-trifluoromethyl-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

Dimethylaminopyridine (0.25 g, 2.01 mmol) was added to a slight suspension of 3-trifluoromethyl-4-N-morpholinoaniline (0.41 g, 1.67 mmol) and N-(3-cyanophenyl)-3-trifluoromethyl-pyrazole-5-carboxylic acid chloride (0.50 g, 1.67 mmol) in 20 mL methylene chloride. The reaction mixture was stirred at ambient temperature for 24h, concentrated under reduced pressure and purified by flash chromatography to give 0.38 g of pure 1-(3-cyanophenyl)-3-trifluoromethyl-5-((3'-trifluoromethyl-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR (dmsO-d<sub>6</sub>, 300MHz) δ: 7.79 (m, 5H), 7.62 (dd, 1H, J<sub>1</sub> = 7.7Hz, J<sub>2</sub> = 8.1Hz), 7.38 (d, 1H, J = 8.4Hz), 7.15 (s, 1H), 3.83 (bt,

4H, J = 4.4), 2.91 (bt, 4H, J = 4.4Hz). Ammonia CI mass spectrum analysis m/z (relative intensity) 510 (M+H, 100). <sup>19</sup>F NMR (DMSO-d6) δ: -61.033 and -62.854.

- 5 Part C. Preparation of 1-(3-amidinophenyl)-3-trifluoromethyl-5-((3'-trifluoromethyl-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole bis-trifluoroacetate.

10 The 1-(3-cyanophenyl)-3-trifluoromethyl-5-((3'-trifluoromethyl-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole (0.38 g, 0.75 mmol) was transformed to the corresponding benzylamine by standard catalytic reduction as described previously to afford 0.19 g of 1-(3-amidinophenyl)-3-trifluoromethyl-5-((3'-trifluoromethyl-4'-(N-morpholino)phenyl)aminocarbonyl)-pyrazole  
15 bis-trifluoroacetate after HPLC purification. <sup>1</sup>H NMR (DMSO-d6) δ: 10.92 (s, 1H), 7.99 (d, 1H, J = 2.6Hz), 7.88-7.85 (m, 1H), 7.68 (s, 1H), 7.63 (s, 1H), 7.55-7.49 (m, 4H), 4.11 (bs, 2H), 3.67-3.64 (m, 4H), 2.78 (bt, 4H, J = 4.4Hz). ESI mass spectrum analysis m/z (relative intensity) 514 (M+H, 100). <sup>19</sup>F NMR (dms-  
20 d6, 300MHz) δ: -59.557, -61.305, and -74.290. HRMS (CI): Calculated for C23H22N5O2F6: 514.167770 found 514.166332.

### Example 233

25 1-(3-aminomethylphenyl)-3-ethyl-5-[(3-fluoro-2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

Part A. Preparation of ethyl-2,4-dioxohexanoate.

30 Sodium metal (16.50 g, 717.39 mmol) was dissolved in 200 mL ethanol. When the solution had cooled 2-butanone (64.26 mL, 717.39 mmol) was added to the solution. After 0.10h diethyl oxalate (97.43 mL, 717.39 mmol) was added to the reaction mixture. Warmed reaction mixture to 65°C for 4h, concentrated  
35 under reduced pressure and treated with 200 mL 1.0M hydrochloric acid solution. Extracted with 200 mL EtOAc and washed organics 2x150 mL water and 2x150 mL brine solution. Dried resulting organics over magnesium sulfate, concentrated



under reduced pressure and purified by flash chromatography to give 21.13 g of pure ethyl-2,4-dioxohexanoate. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 14.40 (bs,1H), 6.38 (s,1H), 4.40-4.32 (m,2H), 2.54 (q, 2H, J = 7.7Hz), 1.41-1.36 (m,3H), 1.18 (t, 3H, J = 7.2Hz).

5

Part B. Preparation of ethyl(2-methylimino)-4-oxohexanoate.

Ethyl 2,4-dioxohexanoate(21.13 g,0.12 mmol) and methoxylamine hydrochloride(10.26 g,0.12 mmol) were added to a suspension of 3Å molecular sieves(30 g) in 500 mL anhydrous ethanol. The reaction mixture was stirred mechanically for 24h. Then the suspension was filtered through a Celite pad and the resulting filtrate was concentrated to give the crude product. Flash chromatography of the crude material gave 6.07 g of pure ethyl(2-methylimino)-4-oxohexanoate. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 4.33 (q, 2H, J = 7.2Hz), 4.06 (s,3H), 3.71 (s,3H), 2.51 (q, 2H, J = 7.2Hz), 1.35 (t, 3H, J = 7.2Hz), 1.08 (t, 3H, J = 7.2Hz). Ammonia CI mass spectrum analysis m/z(relative intensity) 201 (M+H,60), 219 (M+NH<sub>4</sub>,100).

20

Part C. Preparation of ethyl(N-(3-cyanophenyl)-3-ethyl)pyrazole-5-carboxylate

To a solution of ethyl(2-methoxyimino)-4-oxohexanoate(1.0 g, 4.98 mmol) in 50 mL glacial acetic acid was added 3-cyanophenylhydrazine hydrochloride(0.84 g,4.98 mmol). The reaction mixture was warmed to reflux temperature for 4h, concentrated under reduced pressure and purified by flash chromatography to give 0.98 g of ethyl(N-(3-cyanophenyl)-3-ethyl)pyrazole-5-carboxylate. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 7.77-7.76 (m,1H), 7.72-7.68 (m, 2H), 7.56 (t,1H,J=8.0Hz), 6.89 (s,1H), 4.30-4.23 (m,2H), 2.73 (q, 2H, J = 8.0Hz), 1.33-1.27 (m,6H). Ammonia CI mass spectrum analysis m/z(relative intensity) 270 (M+H, 100).

30

Part D. Preparation of 1-(3-cyanophenyl)-3-ethyl-5-((4-bromo-2-fluorophenyl))aminocarbonyl)pyrazole.

35

To a solution of 4-bromo-2-fluoroaniline (2.06 g, 10.82 mmol) and ethyl (3-cyanophenyl)-3-ethylpyrazole-5-carboxylate (0.97 g, 3.61 mmol) in 20 mL methylene chloride was added trimethylaluminum (2.0M in hexanes, 5.41 mL, 10.82 mmol) in a dropwise fashion over 0.3h. The reaction mixture was stirred at ambient temperature for 72h, quenched carefully with 1.0M hydrochloric acid solution, washed 4x50 mL 1.0M hydrochloric acid solution, dried over magnesium sulfate and concentrated under reduced pressure. The resulting residue was purified by flash chromatography to afford 0.23 g of 1-(3-cyanophenyl)-3-ethyl-5-[(4-bromo-2-fluorophenyl)aminocarbonyl]pyrazole. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.17 (t, 1H, J = 8.0Hz), 7.82 (m, 2H), 7.71 (m, 2H), 7.56 (dd, 1H, J<sub>1</sub> = 8.0Hz, J<sub>2</sub> = 7.7Hz), 7.33 (m, 1H), 6.72 (s, 1H), 2.77 (m, 2H), 1.34 (t, 3H, J = 7.7Hz). Ammonia CI mass spectrum analysis m/z (relative intensity) 415 (M+H, 100).

Part E. Preparation of 1-(3-cyanophenyl)-3-ethyl-5-[(3-fluoro-2-tertbutylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

To a nitrogen purged solution of 1-(3-cyanophenyl)-3-ethyl-5-[(4-bromo-2-fluorophenyl)aminocarbonyl]pyrazole (0.23 g, 0.56 mmol), 2-tert-butylaminosulfonylphenylboronic acid (0.17 g, 0.67 mmol) and sodium carbonate (0.12 g, 1.12 mmol) in 10 mL ethanol and 20 mL toluene was added catalytic tetrakis-triphenylphosphine palladium. The reaction mixture was heated to 80°C for 15h, concentrated under reduced pressure and purified by flash chromatography to afford 0.13 g of 1-(3-aminomethylphenyl)-3-ethyl-5-[(2'-fluoro-2-tertbutylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole. <sup>1</sup>H NMR (DMSO-d<sub>6</sub>) δ: 8.36 (t, 1H, J = 8.0Hz), 8.16 (m, 1H), 7.97 (bd, 1H, J = 3.0Hz), 7.85 (s, 1H), 7.77 (d, 1H, J = 8.1Hz), 7.70 (d, 1H, J = 7.8Hz), 7.54 (m, 3H), 7.41 (dd, 1H, J<sub>1</sub> = 1.8Hz, J<sub>2</sub> = 11.7Hz), 7.25 (m, 2H), 6.76 (s, 1H), 3.67 (s, 1H), 2.79 (q, 2H, J = 8.0Hz), 1.36 (t, 3H, J = 8.0Hz), 1.06 (s, 9H). Ammonia CI mass spectrum analysis m/z (relative intensity) 546 (M+H, 100). <sup>19</sup>F NMR (dmsO-d<sub>6</sub>, 300MHz) δ: -130.963.

Part F. Preparation of 1-(3-aminomethylphenyl)-3-ethyl-5-[(3-fluoro-(2-tertbutylaminosulfonyl-[1,1']-biphen-4-yl))aminocarbonyl]pyrazole.

5

Standard transformation of the benzonitrile obtained in part C to the benzylamine via the catalytic reduction followed by treatment with refluxing trifluoroacetic acid converted the 1-(3-cyanophenyl)-3-ethyl-5-[(3-fluoro-2-tertbutylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole to 1-(3-aminomethylphenyl)-3-ethyl-5-[(3-fluoro-2-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole trifluoroacetate. The crude product was purified by standard HPLC purification technique.

<sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.01-7.98 (m, 1H), 7.63-7.56 (m, 4H), 7.45-7.25 (m, 5H), 7.16 (d, 1H, J = 8.4Hz), 6.96 (s, 1H), 3.95 (s, 2H), 2.66 (q, 2H, J = 7.7Hz), 1.24 (t, 3H, J = 7.7Hz). ESI mass spectrum analysis m/z (relative intensity) 493.9 (M+H, 100). HRMS(CI): Calculated for C<sub>25</sub>H<sub>24</sub>N<sub>5</sub>O<sub>3</sub>FS 493.158390, found 493.156279.

20

#### Example 234

1-(3-Aminomethylphenyl)-3-ethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl))aminocarbonyl]pyrazole trifluoroacetate

Part A. Preparation of 1-(3-cyanophenyl)-3-ethylpyrazole-5-carboxylic acid chloride.

To a chilled solution (0°C) of ethyl 1-(3-cyanophenyl)-3-ethylpyrazole-5-carboxylate (7.13 g, 26.51 mmol) in 100 mL water and 150 mL tetrahydrofuran was added lithium hydroxide (1.33 g, 31.81 mmol). The reaction mixture was allowed to warm to ambient temperature overnight and was concentrated under reduced pressure. The resulting aqueous solution was washed 3x100 mL diethylether and acidified with concentrated hydrochloric acid solution to give a white precipitate that was isolated by vacuum filtration. The white solid was placed under high vacuum for 24h and a portion (0.31 g, 1.27 mmol) was treated with oxalyl chloride (0.17 mL, 1.90 mmol) and

dimethylformamide (0.1 mL) in 10 mL methylene chloride. After 24h at ambient temperature the reaction mixture was concentrated and the resulting solid was placed under high vacuum to give the crude 1-(3-cyanophenyl)-3-ethylpyrazole-5-carboxylic acid chloride. The crude acid chloride was used without further purification.

Part B. Preparation of 1-(3-cyanophenyl)-3-ethyl-5-((3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole.

10

To a solution of 2-fluoro-2'-methylsulfonylphenyl)aniline hydrochloride(0.38 g,1.27 mmol) and crude 1-(3-cyanophenyl)-3-ethylpyrazole-5-carboxylic acid chloride(1.27 mmol) in 10 mL dichloromethane was added dimethylaminopyridine (0.34 g, 2.79 mmol). The reaction mixture was stirred at ambient temperature for 24h, concentrated under reduced pressure and purified by flash chromatography to afford 0.23 g of 1-(3-cyanophenyl)-3-ethyl-5-((2'-fluoro-2-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.42 (s,1H), 8.06 (dd,1H,J<sub>1</sub>=2.0Hz,J<sub>2</sub>=8.0Hz), 7.95-7.94 (m,1H), 7.85-7.60 (m,6H), 7.42-7.32 (m,2H), 7.20 (dd, 1H, J<sub>1</sub> = 2.0Hz, J<sub>2</sub> = 8.0Hz), 7.08 (s,1H), 2.89 (s,3H), 2.67 (q, 2H, J = 7.7Hz), 1.24 (t, 3H, J = 7.7Hz). Ammonia CI mass spectrum analysis m/z(relative intensity) 489 (M+H,100).

25

Part C. Preparation of 1-(3-aminomethylphenyl)-3-ethyl-5-(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole.

30

To a suspension of 1-(3-cyanophenyl)-3-ethyl-5-((2'-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole (0.103 g,0.211 mmol) and cobalt chloride (0.003 g,0.021 mmol) in 10 mL methanol was added sodium borohydride (0.016 g,0.422 mmol). After 1h additional sodium borohydride (0.016 g, 0.422 mmol) was added. Let reaction mixture stir for 2h. Then concentrated reaction mixture under reduced pressure and took up resulting residue in 1.0M hydrochloric acid solution to give a white precipitate.

35

Isolated precipitate by vacuum filtration and purified solid by HPLC to give 0.030 g of pure 1-(3-aminomethylphenyl)-3-ethyl-5-(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole trifluoroacetate. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ:

5 10.45 (s, 1H) 8.06 (dd, 1H, J<sub>1</sub>=2.0Hz, J<sub>2</sub> = 8.0Hz), 7.77-7.61 (m, 5H), 7.47-7.31 (m, 4H), 7.21-7.17 (m, 1H), 7.01 (s, 1H), 4.07-4.06 (m, 2H), 2.90 (s, 3H), 2.66 (q, 2H, J=7.7Hz), 1.24 (t, 3H, J=7.7Hz). ESI mass spectrum analysis m/z (relative intensity) 493 (M+H, 100). HRMS Calculated for C<sub>26</sub>H<sub>26</sub>N<sub>4</sub>O<sub>3</sub>FS:

10 493.170966, found 493.172100.

#### Example 235

1-(3-Aminomethylphenyl)-3-ethyl-5-[(2-fluoro-4-(2-methylsulfonylimidazol-1-yl)phenyl)]aminocarbonyl)pyrazole trifluoroacetate

15

Part A. Preparation of 4-(2'-methylthioimidazol-1-yl)nitrobenzene.

20 To a stirred suspension of potassium carbonate (40.07 g, 22.60 mmol) in 175 mL acetone was added 1-(4-nitrophenyl)imidazoline-2-thione (5.0 g, 22.60 mmol). Dripped iodomethane (1.44 mL, 23.05 mmol) into reaction mixture and heated to reflux temperature for 20h. Concentrated reaction mixture under

25 reduced pressure and took up resulting solid in 200 mL water. Extracted aqueous three times with ethyl acetate. Combined extracts, dried over magnesium sulfate and concentrated in vacuo to give 5.29 g of crude 4-(2'-methylthioimidazol-1-yl)nitrobenzene. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.45 (s, 1H) 8.06 (dd, 1H, J<sub>1</sub> = 2.0Hz; J<sub>2</sub> = 8.0Hz), 8.38-8.33 (m, 2H), 7.77-7.72 (m, 2H), 7.61 (d, 1H, J = 1.5Hz), 7.14 (d, 1H, J = 1.5Hz), 2.52 (s, 3H). ESI mass spectrum analysis m/z (relative intensity) 236 (M+H, 100).

30

35 Part B. Preparation of 4-(2'-methylsulfonylimidazol-1-yl)nitrobenzene.

To a cooled solution (0°C) of 4-(2'-methylthioimidazol-1-yl)nitrobenzene (1.05 g, 4.47 mmol) in 40 mL dichloromethane was added meta-chloroperoxybenzoic acid (1.54 g, 8.94 mmol). The reaction mixture was allowed to warm to ambient temperature over 20H. Washed reaction mixture 3x75 mL 1.0M sodium hydroxide solution. Dried resulting organics over magnesium sulfate and concentrated under reduced pressure to give 0.98 g of crude 4-(2'-methylsulfonylimidazol-1-yl)nitrobenzene. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 8.39 (d, 2H, J = 8.7Hz), 7.73 (d, 2H, J = 8.7Hz), 7.28-7.23 (m, 2H), 3.43 (s, 3H). Ammonia CI mass spectrum analysis m/z(relative intensity) 268 (M+H, 100).

Part C. Preparation of 4-(2'-methylsulfonylimidazol-1-yl)aniline.

Standard catalytic reduction of 4-(2'-methylsulfonylimidazol-1-yl)nitrobenzene (0.98 g, 3.67 mmol) with palladium on carbon(10%) in methanol gave 0.80 g of 4-(2'-methylsulfonylimidazol-1-yl)aniline.

<sup>1</sup>HNMR (CDCl<sub>3</sub>) δ: 7.24 (d, 2H, J=8.7Hz), 7.15 (dd, 2H, J<sub>1</sub> = 18.3Hz, J<sub>2</sub> = 18.6Hz), 6.72 (d, 2H, J = 8.7Hz), 3.30 (s, 3H). Ammonia CI mass spectrum analysis m/z(relative intensity) 238 (M+H, 100).

Part C. Preparation of 1-(3-cyanophenyl)-3-ethyl-5-((2'-fluoro-4'-(2-methylsulfonylimidazol-1-yl)phenyl))aminocarbonylpyrazole.

Dimethylaminopyridine (0.42 g, 3.48 mmol) was added to a solution of 4-(2'-methylsulfonylimidazol-1-yl)aniline (0.37 g, 1.58 mmol) and 1-(3-cyanophenyl)-3-ethylpyrazole-5-carboxylic acid chloride (1.58 mmol) in 15 mL dichloromethane. The reaction mixture was stirred at ambient temperature for 15H, concentrated under reduced pressure and purified by flash chromatography to give 0.37 g of 1-(3-cyanophenyl)-3-ethyl-5-[(2'-fluoro-4'-(2-methylsulfonylimidazol-1-yl)phenyl)]aminocarbonylpyrazole. ESI mass spectrum analysis m/z(relative intensity) 460.9 (M+H, 100), 482.9 (M+Na).

Part D. Preparation of 1-(3-aminomethylphenyl)-3-ethyl-5-[(2'-fluoro-4'-(2-methylsulfonylimidazol-1-yl)phenyl)]aminocarbonylpyrazole.

5

Standard catalytic reduction of 1-(3-cyanophenyl)-3-ethyl-5-[(2'-fluoro-4'-(2-methylsulfonylimidazol-1-yl)phenyl)]aminocarbonylpyrazole with palladium on carbon (10%) in methanol gave 0.10 g of 1-(3-aminomethylphenyl)-3-ethyl-5-[(2'-fluoro-4'-(2-methylsulfonylimidazol-1-yl)phenyl)]aminocarbonylpyrazole trifluoroacetate after HPLC purification. <sup>1</sup>HNMR(CDCl<sub>3</sub>, 300MHz) δ: 10.78 (s, 1H), 7.76 (d, 2H, J = 8.8Hz), 7.63 (d, 2H, J = 1.1Hz), 7.49-7.35 (m, 5H), 7.26 (d, 1H, J = 1.1Hz), 6.98 (s, 1H), 4.08 (s, 2H), 3.35 (s, 3H), 2.67 (q, 2H, J = 7.7Hz), 1.24 (t, 3H, J = 7.7Hz). ESI mass spectrum analysis m/z (relative intensity) 464.9 (M+H, 100). HRMS calculated for C<sub>23</sub>H<sub>25</sub>N<sub>6</sub>O<sub>3</sub>S: 465.170886, found 465.172332.

10

15

#### Example 236

20 1-[(6-(aminomethyl)pyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

Part A. Preparation of ethyl 1-[pyrid-2-yl]-3-methylpyrazole-5-carboxylate.

25

To a solution of 2-hydrazinopyridine (0.68 g, 6.24 mmol) in 15 mL of glacial acetic acid was added ethyl 2-methoxyimino-4-oxopentanoate (0.90 g, 4.80 mmol). The resulting mixture was allowed to stir at 100° C for 2 h. The reaction mixture was cooled to room temperature and concentrated *in vacuo*. The residue was diluted with ethyl acetate, washed with saturated aq sodium carbonate and brine, dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 3:1 hexanes/ethyl acetate) to give 0.4 g (36%) of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.45 (dd, 1H), 7.82 (td, 1H), 7.61 (d, 1H), 7.29 (dd, 1H), 6.70 (s, 1H), 4.25 (q, 2H),

30

35

2.38 (s, 3H), 1.23 (t, 3H). Ammonia CI mass spectrum analysis  
m/z(relative intensity) 232 (M+H, 100).

Part B. Preparation of ethyl 1-[6-cyanopyrid-2-yl]-3-  
5 methylpyrazole-5-carboxylate.

To a solution of of ethyl 1-[pyrid-2-yl]-3-  
methylpyrazole-5-carboxylate (1.4 g, 6.05 mmol) in 10 mL of  
glacial acetic acid was added 6 mL (large excess) of 30% H<sub>2</sub>O<sub>2</sub>.  
10 The reaction was stirred at 100° C for 3 h and then was allowed  
to cool to room temperature and was poured into saturated aq  
sodium carbonate. The resulting mixture was extracted with  
ethyl acetate and the combined organic extracts were washed  
with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The  
15 resulting crude N-oxide was dissolved in 20 mL of  
tetrahydrofuran and then there was added trimethylsilyl cyanide  
(2.4 mL, 18.2 mmol) followed by dimethylcarbamoyl chloride (1.7  
mL, 18.2 mmol). The reaction was allowed to stir at 65° C for  
18 h. The reaction was allowed to cool and was diluted with  
20 ethyl acetate, washed with saturated aq sodium bicarbonate and  
brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue  
was purified by flash chromatography (elution with 3:1  
hexanes/ethyl acetate) to give 0.66 g (43%) of the title  
compound as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.98 (m, 2H), 7.61  
25 (td, 1H), 6.67 (s, 1H), 4.38 (q, 2H), 2.38 (s, 3H), 1.32 (t,  
3H). Ammonia CI mass spectrum analysis m/z(relative intensity)  
257 (M+H, 100).

Part C. Preparation of 1-[(6-cyanopyrid-2-yl)]-3-methyl-5-  
30 [(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-  
yl)aminocarbonyl]pyrazole.

To a solution of (2'-tert-butylaminosulfonyl-[1,1']-  
biphen-4-yl)amine (0.24 g, 0.78 mmol) in 20 mL of methylene  
35 chloride at 25° C was added trimethylaluminum (1.2 mL of a 2.0  
M solution in toluene, 2.34 mmol) dropwise. The resulting  
solution was allowed to stir until no more gas evolution was  
observed (~ 15 min). To this solution was added ethyl 1-[6-



cyanopyrid-2-yl]-3-methylpyrazole-5-carboxylate (0.20 g, 0.78 mmol) as a solution in methylene chloride. The resulting solution was stirred at 40° C for 3 h and then was cooled to 25° C and quenched by the addition of saturated aq NH<sub>4</sub>Cl.

- 5 After diluting with ethyl acetate, the layers were separated and the organic layer was washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 1:1 hexanes/ethyl acetate) to afford 0.15 g (38%) of the title compound as a solid.
- 10 <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 10.63 (s, 1H), 8.20 (t, 1H), 8.08 (d, 1H), 7.98 (m, 2H), 7.64 (d, 2H), 7.59 (td, 1H), 7.51 (td, 1H), 7.34 (d, 2H), 7.28 (d, 1H), 6.80 (s, 1H), 6.46 (s, 1H), 2.31 (s, 3H), 0.97 (s, 9H). ESI mass spectrum analysis m/z(relative intensity) 515.1 (M+H, 100).

15

Part D. Preparation of 1-[(6-(aminomethyl)pyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

- 20 To a solution of 1-[(6-cyanopyrid-2-yl)]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (0.14 g, 0.27 mmol) in 15 mL of absolute ethanol was added 12 N HCl (0.023 mL, 0.27 mmol) and 10% Pd/C catalyst (30 mg). The resulting mixture was stirred
- 25 under 1 atm of H<sub>2</sub> for 18 h. The mixture was then filtered through a pad of celite and was concentrated in vacuo. The residue was taken up in 3 mL of trifluoroacetic acid and stirred at 80° C for 20 min. This solution was cooled and concentrated in vacuo. The residue was purified by prep HPLC
- 30 (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 70 mg (45%) of the title compound as a white powder. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ 10.56 (s, 1H), 8.18 (broad s, 3H), 8.02 (m, 2H), 7.64 (m, 4H), 7.58 (m, 2H), 7.45 (d, 1H), 7.33 (d, 2H), 7.27 (m, 2H), 6.84 (s, 1H),
- 35 4.02 (broad q, 2H), 2.30 (s, 3H). ESI mass spectrum analysis m/z(relative intensity) 462.9 (M+H, 100).

## Example 237

1-[(6-(*N*-hydroxyamidino)pyrid-2-yl)]-3-methyl-5-[(2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole

- 5 Preparation of 1-[(6-(*N*-hydroxyamidino)pyrid-2-yl)]-3-methyl-5-[(2'-*tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

To a solution of 1-[(6-cyanopyrid-2-yl)]-3-methyl-5-[(2'-  
10 *tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (0.11 g, 0.21 mmol) in 5 mL of absolute ethanol was added hydroxylamine hydrochloride (0.054 g, 0.77 mmol) and sodium carbonate (0.039 g, 0.36 mmol). This mixture was stirred at 80° C for 1 h and then was allowed to  
15 cool. The mixture was diluted with ethyl acetate, washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The solid residue was triturated with ether to afford 80 mg (68%) of the title compound as a white solid. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 10.79 (s, 1H), 9.95 (s, 1H), 8.0 (dd, 1H), 7.95 (t, 1H), 7.80 (d, 1H), 7.68  
20 (m, 3H), 7.59 (td, 1H), 7.51 (td, 1H), 7.35 (m, 3H), 6.68 (s, 1H), 6.65 (s, 1H), 5.43 (broad s, 2H), 2.31 (s, 3H), 0.96 (s, 9H). ESI mass spectrum analysis m/z(relative intensity) 548.1 (M+H, 100).

25

## Example 238

1-[(6-amidinopyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

- Preparation of 1-[(6-amidinopyrid-2-yl)]-3-methyl-5-[(2'-  
30 aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

To a solution of 1-[(6-cyanopyrid-2-yl)]-3-methyl-5-[(2'-  
35 *tert*-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (0.28 g, 0.54 mmol) in 20 mL of absolute ethanol was added triethylamine (0.38 mL, 2.7 mmol). Hydrogen sulfide gas was bubbled slowly through this solution (excess H<sub>2</sub>S was scrubbed through Chlorox bleach) for 20 min.

The flask was stoppered tightly and allowed to stand at room temperature overnight. The solution was concentrated *in vacuo*. The crude thioamide residue was dissolved in 10 mL of acetone and then there was added 2 mL (large excess) of methyl iodide. 5 The resulting solution was stirred at 60° C for 2 h and then was cooled and concentrated *in vacuo*. The residue was dissolved in methanol and then there was added ammonium acetate (1.8 mL of a 1.5 M solution in methanol, 2.7 mmol). The resulting mixture was stirred at 60° C for 2 h and then was 10 cooled and concentrated *in vacuo*. The residue was dissolved in trifluoroacetic acid and stirred at 80° C for 20 min and then was allowed to cool and was concentrated *in vacuo*. The residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to 15 afford 78 mg (24%) of the title compound as a white powder. <sup>1</sup>HNMR(d<sub>6</sub>-DMSO) δ: 10.70 (s, 1H), 9.36 (broad s, 2H), 9.04 (broad s, 2H), 8.31 (t, 1H), 8.13 (m, 2H), 8.00 (d, 1H), 7.63 (d, 2H), 7.58 (m, 2H), 7.34 (d, 2H), 7.28 (d, 1H), 7.23 (broad s, 2H), 6.87 (s, 1H), 2.33 (s, 3H). ESI mass spectrum analysis 20 m/z(relative intensity) 476.2 (M+H, 100). HRMS: calculated for C<sub>23</sub>H<sub>22</sub>N<sub>7</sub>O<sub>3</sub>S: 476.150485; Observed: 476.152830.

#### Example 239

1-[6-amidinopyrid-2-yl]-3-methyl-5-[3-fluoro-(2'- 25 methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt

Part A. Preparation of ethyl 1-[(6-thiocarbonylamino)pyrid-2-yl]-3-methylpyrazole-5-carboxylate.

30 To a solution of ethyl 1-[6-cyanopyrid-2-yl]-3-methylpyrazole-5-carboxylate in 100 mL of absolute ethanol was added triethylamine (2.7 mL, 19.4 mmol). Hydrogen sulfide gas was slowly bubbled through this solution (excess H<sub>2</sub>S was 35 scrubbed through Chlorox bleach) for 20 min. The flask was stoppered tightly and allowed to stand at room temperature overnight. The solution was concentrated *in vacuo*. The residue was dissolved in ethyl acetate, washed with 10% aq HCl

and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo to afford 1.1 g (97%) of the title compound which was sufficiently pure to be used without purification. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 9.01 (broad s, 1H), 8.55 (dd, 1H), 7.92 (t, 1H), 7.82 (dd, 1H), 7.58 (broad s, 1H), 6.66 (s, 1H), 4.22 (q, 2H), 2.33 (s, 3H), 1.18 (t, 3H).  
ESI (-ve) mass spectrum analysis m/z(relative intensity) 288.9 (M-H, 100).

Part B. Preparation of 1-[(6-thiocarbonylamino)pyrid-2-yl]-3-methyl-5-[(1-bromo-3-fluorophenyl-4-yl)aminocarbonyl]pyrazole.

To a solution of 4-bromo-2-fluoroaniline (2.17 g, 11.4 mmol) in 150 mL of methylene chloride was added trimethylaluminum (11.4 mL of a 2M solution in toluene, 22.8 mmol) dropwise. This solution was stirred until gas evolution ceased (15-20 min) and then there was added ethyl 1-[(6-thiocarbonylamino)pyrid-2-yl]-3-methylpyrazole-5-carboxylate (1.1 g, 3.8 mmol) in methylene chloride. The resulting solution was allowed to stir at reflux for 18 h and then it was cooled and quenched by dropwise addition of saturated aq ammonium chloride. The mixture was diluted with ethyl acetate, the layers were separated, the organic layer was washed with water and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The solid residue was purified by trituration with ether and the remaining solid was dried in vacuo to afford 1.26 g (76%) of the title compound. <sup>1</sup>HNMR(d<sub>6</sub>-DMSO) δ: 10.62 (broad s, 1H), 10.20 (broad s, 1H), 8.84 (broad s, 1H), 8.33 (dd, 1H), 8.12 (t, 1H), 7.98 (d, 1H), 7.72 (t, 1H), 7.58 (dd, 1H), 7.39 (d, 1H), 6.75 (s, 1H), 2.30 (s, 3H)ppm.

Part C. Preparation of 1-[(6-(N-tert-butylloxycarbonyl)aminoiminomethyl)pyrid-2-yl]-3-methyl-5-[(1-bromo-3-fluorophenyl-4-yl)aminocarbonyl]pyrazole.

To a solution of 1-[(6-thiocarbonylamino)pyrid-2-yl]-3-methyl-5-[(1-bromo-3-fluorophenyl-4-yl)aminocarbonyl]pyrazole (1.09 g, 2.51 mmol) in 100 mL of acetone was added 12 mL (large excess) of methyl iodide. The resulting solution was stirred

at 60° C for 2 h and then was cooled and concentrated in vacuo. The residue was dissolved in methanol and then there was added ammonium acetate (8.3 mL of a 1.5 M solution in methanol, 12.5 mmol). The resulting mixture was stirred at 60° C for 2 h and  
5 then was cooled and concentrated in vacuo to afford 1.0 g of the crude amidine. To 0.5 g (1.2 mmol) of this residue in 10 mL of pyridine was added di-tert-butyl dicarbonate (0.52 g, 2.4 mmol) and 4-dimethylaminopyridine (0.29 g, 2.4 mmol). This mixture was allowed to stir at room temperature for 18 h and  
10 then was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with water, 10% aq HCl and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 1:1 hexanes/ethyl acetate) to afford 0.15 g (24%) of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ:  
15 9.08 (broad s, 1H), 8.22 (m, 3H), 7.95 (d, 1H), 7.85 (t, 1H), 7.25 (m, 2H), 6.53 (s, 1H), 2.33 (s, 3H), 1.49 (s, 9H)ppm. ESI mass spectrum analysis m/z 516.9/518.9 (M+H)+.

Part D. Preparation of 1-[(6-(N-tert-  
20 butyloxycarbonyl)aminoiminomethyl)pyrid-2-yl]-3-methyl-5-[3-fluoro-(2'-thiomethoxy-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole.

To a solution of 1-[(6-(N-tert-  
25 butyloxycarbonyl)aminoiminomethyl)pyrid-2-yl]-3-methyl-5-[(1-bromo-3-fluorophenyl-4-yl)aminocarbonyl]pyrazole (0.15 g, 0.29 mmol) in 15 mL of benzene was added 2-thiomethoxyphenyl boronic acid (0.07 g, 0.42 mmol), tetrabutylammonium bromide (0.01 g, 0.03 mmol), sodium carbonate (0.09 g, 0.85 mmol) and 0.80 mL of  
30 water. This mixture was degassed with a stream of nitrogen and then tetrakis triphenylphosphine palladium (0.06 g, 0.05 mmol) was added. The mixture was stirred at 80° C for 24 h. The reaction was allowed to cool and then was diluted with ethyl acetate, washed with saturated aq sodium bicarbonate and brine,  
35 dried (MgSO<sub>4</sub>), filtered through celite and concentrated in vacuo to afford 0.157 g (95%) of the title compound. This material was sufficiently pure to be used without purification.

<sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 8.40 (t, 1H), 8.02 (broad s, 2H), 7.60-7.20 (m, 10H), 6.56 (s, 1H), 2.34 (s, 3H), 2.33 (s, 3H), 1.46 (s, 9H) ppm. ESI mass spectrum analysis m/z (relative intensity) 560.9 (M+H, 100).

5

Part E. Preparation of 1-[6-amidinopyrid-2-yl]-3-methyl-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

10

To a solution of 1-[(6-(N-tert-butyloxycarbonyl)aminoiminomethyl)pyrid-2-yl]-3-methyl-5-[3-fluoro-(2'-thiomethoxy-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole (0.157 g, 0.28 mmol) in 20 mL of methylene chloride was added 3-chloroperoxybenzoic acid (0.17 g, 0.99 mmol). The resulting mixture was stirred at room temperature for 24 h and then was diluted with ethyl acetate, washed with saturated aq sodium metabisulfite, saturated aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was dissolved in 5 mL of trifluoroacetic acid and stirred at 80° C for 20 min. The reaction was cooled and concentrated in vacuo. The residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 80 mg (47%) of the title compound as a white powder. <sup>1</sup>HNMR(d<sub>6</sub>-DMSO) δ: 10.52 (s, 1H), 9.42 (broad s, 2H), 9.08 (broad s, 2H), 8.31 (t, 1H), 8.12 (m, 3H), 7.78-7.73 (m, 3H), 7.42 (d, 1H), 7.32 (d, 1H), 7.20 (d, 1H), 6.89 (s, 1H), 2.89 (s, 3H), 2.33 (s, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity) 493.9 (M+H, 100).

30

#### Example 240

1-(3-aminomethylphenyl)-3-methyl-5-((2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole trifluoroacetate

35

Part A: Preparation of 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-5-((2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole.

To a solution of 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-pyrazole-5-carboxylic acid (183 mg, 0.5 mmol) in DMF (10 mL) was added PyBrop (bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, 280 mg, 0.6 mmol) and the resulting solution was stirred at room temperature for 10 min. N,N-diisopropylethylamine (1 mL) was added and stirred for additional 10 min. To this solution was then added 2-methoxy-4-N-morpholine-aniline (125 mg, 0.6 mmol) and the resulting mixture was stirred at 60°C for 3 hours. After the mixture was cooled to room temperature, to it was added DOWEX (50WX8-100 ion-exchange resin, 0.5 g) and stirred for additional 0.5h. The mixture was filtered and the residue was washed with EtOAc (50 mL). The filtrate was washed with brine (5 x 10 mL), dried over MgSO<sub>4</sub>, and purified by column chromatography with EtOAc to give the product (261 mg, 95%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.42-7.31 (m, 10H), 7.03 (s, 1H), 6.94 (d, J = 8.8 Hz, 1H), 6.50 (d, J = 2.6 Hz, 1H), 6.42 (dd, J = 8.4 Hz, J = 2.6 Hz, 1H), 4.70 (s, 1H), 4.41 (d, J = 3.9 Hz, 2H), 3.84 (t, J = 4.8 Hz, 4H), 3.78 (s, 3H), 3.09 (t, J = 4.8 Hz, 4H), 2.35 (s, 3H) ppm. ESI mass spectrum analysis m/z(relative intensity) 556 (M+H, 100).

Part B: Preparation of 1-(3-aminophenyl)-3-methyl-5-((2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole trifluoroacetate.

To 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-5-((2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole (100 mg, 0.18 mmol) was added trifluoroacetic acid (5 mL) and the resulting solution was refluxed for 4 hours. The solution was concentrated and purified on TLC plate with ethyl acetate to a viscous liquid (60 mg, 80%). <sup>1</sup>HNMR(CD<sub>3</sub>OD) δ: 7.58 (s, 1H), 7.53-7.48 (m, 3H), 7.06 (s, 1H), 6.91 (d, J = 8.4 Hz, 1H), 6.63 (d, J = 2.6 Hz, 1H), 6.47 (dd, J = 8.8 Hz, J = 2.6 Hz, 1H), 4.15 (s, 2H), 3.79 (t, J = 4.8 Hz, 4H), 3.76 (s, 3H), 3.09 (t, J = 4.8 Hz, 4H), 2.35 (s, 3H) ppm. ESI mass spectrum analysis m/z(relative intensity) 422 (M+H, 100).

## Example 241

1-(3-aminomethylphenyl)-3-methyl-5-[(4'-(3"-methyl-5"-oxo-3"-pyrazolin-2"-yl)-phenyl)aminocarbonyl]pyrazole trifluoroacetate

- 5 Part A: Preparation of 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-5-((4'-(3"-methyl-5"-oxo-3"-pyrazolin-2"-yl)-phenyl)aminocarbonyl)-pyrazole.

To a solution of 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-pyrazole-5-carboxylic acid (150 mg, 0.41 mmol) in DMF (5 mL) was added PyBrop (bromo-tris-pyrrolidino-phosphonium hexafluorophosphate, 233 mg, 0.5 mmol) and the resulting solution was stirred at room temperature for 10 min. To this solution was added N,N-dimethylpyridine (70 mg, 0.57 mmol) and stirred for an additional 10 min. 2-(4-aminophenyl)-3-methyl-3-pyrazolin-5-one (125 mg, 0.6 mmol) was added and the resulting mixture was stirred at 60°C for 24 hours. The mixture was diluted with EtOAc (100 mL), washed with 1N HCl (10 mL) and brine (5 x 10 mL), dried over MgSO<sub>4</sub>, and purified by column chromatography with EtOAc to afford the product (260 mg). ESI mass spectrum analysis m/z(relative intensity) 537.2 (M+1, 100).

- 25 Part B: Preparation of 1-(3-aminophenyl)-3-methyl-5-((2'-methoxy-4'-(N-morpholino)phenyl)aminocarbonyl)pyrazole trifluoroacetate.

To 1-(3-N-(benzyloxycarbonyl)aminophenyl)-3-methyl-5-((4'-(3"-methyl-5"-oxo-3"-pyrazolin-2"-yl)-phenyl)aminocarbonyl)-pyrazole (260 mg) was added trifluoroacetic acid (5 mL) and the resulting solution was refluxed for 2 hours. The solution was concentrated and purified on TLC plate with ethyl acetate to a viscous liquid (120 mg, 74.6% for two steps). <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ: 7.69 (d, J = 8.8 Hz, 2H), 7.55 (7.55 (bs, 1H), 7.52-7.46 (m, 3H), 7.35 (d, J = 8.8 Hz, 2H), 6.87 (s, 1H), 5.57 (s, 1H), 4.14 (s, 2H), 2.35 (s, 3H), 2.21 9 (s, 3H)ppm. ESI mass spectrum analysis m/z(relative intensity) 403.1 (M+H, 100).



## Example 242

1-[3-(aminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid salt

5

Part A: Preparation of 1,1-di(methylthio)ethylene.

In a 2 L flask fitted with mechanical stirrer, condenser, under argon, methyl magnesium bromide (3.0 M in Et<sub>2</sub>O, 84 mL, 252 mmol) was diluted to 1.0 M in THF (168 mL), keeping the pot  
10 temperature below 40°C. Carbon disulfide (22.6 mL, 376 mmol) in THF (23 mL) was added over 30 min., and the reaction was maintained at 40°C for 135 min. Heat was removed and the reaction was cooled to -72°C. Lithium diisopropylamide (2.0 M  
15 in heptane, THF, and ethylbenzene, 126 mL, 252 mmol) was added over 35 min., keeping the internal temperature below -60°C. The resulting thick, dark orange-red paste was maintained near -60°C for 160 min. Dimethyl sulfate (48 mL, 504 mmol) was added over 45 min., and the reaction was allowed to warm to  
20 room temperature over 70 min. The mechanical stirrer was turned off, and the reaction stood at room temperature for 17 h. The resulting mixture was diluted with Et<sub>2</sub>O (300 mL) and poured into aq. sodium bicarbonate (20%, 500 mL). An argon atmosphere was maintained for all manipulations. The layers  
25 were separated, and the organics were extracted with aq. sodium bicarbonate (25%, 200 mL), dried over MgSO<sub>4</sub>, filtered, and concentrated to about 100 mL. The resulting oil was distilled under vacuum (70°C head temperature, 10 Torr) to yield 25.37 g product contaminated with ethylbenzene, for an estimated yield  
30 of pure product (15.59 g, 52%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 5.24 (s, 2H), 2.36 (s, 6H)ppm.

Part B: Preparation of methyl 4,4-di(methylthio)-2-oxo-but-3-enoate.

35

A solution of 1,1'-di(methylthio)ethylene (19.73 g containing 9.95 g of compound, 83 mmol) in Et<sub>2</sub>O (125 mL) was cooled to -60°C under argon. Oxalyl chloride (5.6 mL, 64 mmol)

was added over 3 min., allowing the internal temperature to reach -55°C. The reaction was warmed to -15°C over 20 min., and dry methanol (20 mL, 494 mmol) was added over 2 min. The reaction continued to warm and stir at room temperature for 2 h. The resulting mixture was diluted with Et<sub>2</sub>O and filtered under argon to yield a yellow solid (8.28 g, 63%).  
<sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 6.84 (s, 1H), 3.87 (s, 3H), 2.57 (s, 3H), 2.55 (s, 3H) ppm.

10 Part C: Preparation of methyl 1-(3-cyanophenyl)-3-(methylthio)pyrazole-5-carboxylate.

A mixture of methyl 4,4-di(methylthio)-2-oxo-but-3-enoate (2.0 g, 9.7 mmol), triethylamine (1.5 mL, 10.7 mmol), and *m*-cyanophenylhydrazine hydrochloride (1.81 g, 10.7 mmol) were combined in dry methanol (20 mL) and heated at reflux for 47 h. The reaction was evaporated and chromatographed on silica gel (CH<sub>2</sub>Cl<sub>2</sub>, followed by 40% EtOAc / hexanes) to yield a partially purified intermediate (1.91 g), which was redissolved in acetonitrile (85 mL) and refluxed 23 h. The crude reaction mixture was chromatographed on silica gel in CH<sub>2</sub>Cl<sub>2</sub>, to yield desired pyrazole (780 mg, 29%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.78 (s, 1H), 7.70 (m, 2H), 7.57 (m, 1H), 6.95 (s, 1H), 3.83 (s, 3H), 2.57 (s, 3H) ppm.

25 Part D: Preparation of methyl 1-[3-(aminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylate.

To a solution of methyl 1-(3-cyanophenyl)-3-(methylthio)pyrazole-5-carboxylate (777 mg, 2.8 mmol) in dry DMF (50 mL), CoCl<sub>2</sub> (39 mg, 0.30 mmol) and NaBH<sub>4</sub> (158 mg, 4.2 mmol) were added. The initial solution was emerald green, then turned dark black. After stirring for 2 h., additional NaBH<sub>4</sub> (145 mg, 3.8 mmol) was added. After another 3 h., additional CoCl<sub>2</sub> (330 mg, 2.5 mmol) was added. The reaction continued stirring at room temperature for 17 h. Methanol (10 mL) was added and stirred 40 min. to quench the reaction. The reaction was concentrated to 30 mL and chromatographed on silica gel

(0%-100% EtOAc / hexanes followed by 10-30% MeOH / CHCl<sub>3</sub>) to yield the desired product (198 mg, 25%). <sup>1</sup>HNMR(CDCl<sub>3</sub>) δ: 7.41 (m, 3H), 7.30 (d, 1H, J = 7.3), 6.90 (s, 1H), 4.02 (bs, 1H), 3.78 (s, 3H), 3.49 (s, 2H), 2.54 (s, 3H)ppm.

5

Part E: Preparation of methyl 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylate.

10 Di-t-butyl dicarbonate (184 mg, 0.84 mmol) was added to a suspension of methyl 1-[3-(aminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylate (195 mg, 0.70 mmol) in dry THF (8 mL). After stirring 3 h., additional THF (5 mL) was added to aid solubility. The reaction was stirred an  
15 additional 16 h., and additional di-t-butyl dicarbonate (54 mg, 0.25 mmol) was added. After another 5 h., triethylamine (100 µL, 0.72 mmol) was added and stirred 2 h. The reaction was diluted with EtOAc and extracted twice with H<sub>2</sub>O. The aqueous were combined and extracted with EtOAc. The organics were  
20 combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, evaporated, and chromatographed on silica gel (30% EtOAc) to yield the desired product (228 mg, 86%). <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.37 (m, 4H), 6.91 (s, 1H), 4.87 (bs, 1H), 4.38 (d, 2H, J = 5.8), 3.79 (s, 3H), 2.56 (s, 3H), 1.46 (s, 9H)ppm.

25

Part F: Preparation of 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylic acid.

30 To a solution of methyl 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylate (50 mg, 0.13 mmol) in THF (2 mL) was added aq. LiOH (1.0 M, 160 µL, 0.16 mmol). The resulting solution was stirred for 19 h. Additional LiOH (30 µL, 0.03 mmol) was added and  
35 stirred for 3 h. The reaction was partitioned between H<sub>2</sub>O and Et<sub>2</sub>O / EtOAc. The aqueous extracts were neutralized with HCl (0.1 M, 1.0 mL) and ice. This aqueous solution was extracted once with Et<sub>2</sub>O / EtOAc. Additional HCl (0.1 M, 0.5 mL) was

added and further extracted with Et<sub>2</sub>O / EtOAc. A final pH of 3.5 was reached with additional HCl (0.1 M, 0.4 mL). This was extracted again with EtOAc. The organic extracts after acidification were combined, dried over MgSO<sub>4</sub>, filtered, and evaporated to yield the desired product (54 mg, 100%).  
5 <sup>1</sup>H NMR (CDCl<sub>3</sub>): 7.33 (m, 4H), 6.97 (s, 1H), 4.35 (bd, 2H, J = 4.4), 4.27 (bs, 1H), 2.55 (s, 3H), 1.45 (s, 9H) ppm.

Part G: Preparation of 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole.  
10

DMF (3 or 4 drops) was added to a mixture of 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylic acid (94 mg, 0.26 mmol) and oxalyl chloride (35 µL, 0.40 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (3 mL). The resulting solution was stirred for 55 min. and evaporated. After a few min. under high vacuum, the compound was redissolved in CH<sub>2</sub>Cl<sub>2</sub> (3 mL), and 4-amino-2'-methylsulfonyl-[1,1']-biphenyl hydrochloride (85 mg, 0.30 mmol) and 4-dimethylaminopyridine (85 mg, 0.70 mmol) were added and stirred 20 h. The reaction was diluted with H<sub>2</sub>O and extracted twice with EtOAc. The combined organics were extracted with aq. NaHCO<sub>3</sub>, followed by aq. HCl (0.1 M, cooled with ice). Solid NaCl was added to aid separation. The organic layer was removed, and the aqueous solution was extracted an additional 2 times with EtOAc. The organic extracts were combined, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered, and evaporated. The crude product was chromatographed on silica gel (50% EtOAc / hexanes) to yield the desired product (65 mg, 43%). ESI mass spectrum analysis m/z = 615 (M+Na)<sup>+</sup>.  
25  
30

Part H: Preparation of 1-[3-(aminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid salt.  
35

1-[3-(t-Butoxycarbonylaminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole (65 mg, 0.11 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub>,

(3 mL) and TFA (1 mL) and stirred 17 h. The reaction was evaporated and purified by prep. HPLC (10-90% MeCN / H<sub>2</sub>O / 0.5% TFA) to yield the desired product (37 mg, 55%). <sup>1</sup>HNMR(DMSO)-10.78 (s, 1H), 8.21 (bs, 2H), 8.08 (d, 1H, J = 7.7), 7.70 (m, 5H), 7.45 (m, 6H), 7.16 (s, 1H), 4.13 (bd, 2H, J = 4.8), 2.84 (s, 3H), 2.57 (s, 3H)ppm. ESI mass spectrum analysis m/z = 493 (M+H, 100).

#### Example 243

10      1-(3-aminomethyl-4-fluorophenyl)-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-methyl pyrazole.

To a mixture of 3-cyano-4-fluorophenylhydrazine tin chloride (10 g, 26.6 mmol) in acetic acid(150 mL) was added 1,1,1-trifluoro-2,4-pentanedione (4.09 g, 26.6 mmol). The reaction mixture was brought to reflux overnight. Acetic acid was removed on rotary evaporator under reduced pressure. Residue was partitioned between ethyl acetate (200 mL) and water (150 mL). Organic phase was separated and washed with water (3 x 100 mL), dried over sodium sulfate; filtered, concentrated and subjected to silica-gel flash chromatography(ethyl acetate:hexane, 1:10) to afford 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-methyl pyrazole(4.0 g). CI mass spectrum m/z (rel. intensity) 270 (M+H, 100).

30      Part B: Preparation of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-bromomethyl pyrazole.

To a solution of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-methyl pyrazole(4.0 g, 14.87 mmol) in carbon tetrachloride(50 mL) was added NBS(2.65 g, 14.87 mmol) and benzoyl peroxide(0.36 g, 1.48 mmol). The reaction mixture was brought to reflux overnight. Solvent was removed on rotary evaporator under reduced pressure. Residue was partitioned

between ethyl acetate(80 mL) and sodium bicarbonate(sat. 80 mL). Organic phase was separated and washed with water(60 mL); dried over sodium sulfate; filtered, concentrated and subjected to silica-gel flash chromatography (ethyl acetate : hexane, 1:10) to afford 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-bromomethyl pyrazole(2.5 g). CI mass spectrum m/z (rel. intensity) 348 (M+H, 100).

Part C: Preparation of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxymethyl pyrazole.

To a solution of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-bromomethyl pyrazole(2.5 g, 7.18 mmol) in DMSO (40 mL) was added copper(I) oxide(2.16 g, 15.08 mmol) and water (12 mL). The reaction mixture was stirred at 60 °C for 2 hours then cooled to RT and stirred at RT overnight. The next day, the mixture was filtered through celite, filter pad was washed with ethyl acetate(20 mL); the filtrate was partitioned between ethyl acetate(50 mL) and water(50 mL); organic phase was separated and washed with water(3 x 30 mL); dried over sodium sulfate; filtered, concentrated, flash chromatography (ethyl acetate : hexane, 1:6) to afford 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxymethyl pyrazole(1.7 g). CI mass spectrum m/z (rel. intensity) 286 (M+H, 100).

Part D: Preparation of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxycarbonylmethyl pyrazole.

To a solution of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxymethyl pyrazole(1.5 g, 5.26 mmol) in acetonitrile(30 mL) was added NaIO<sub>4</sub>(2.65 g, 11.05 mmol), catalytic amount of RuCl<sub>3</sub> and water(30 mL) at 0 °C. The reaction mixture was stirred at 0 °C to RT overnight. Acetonitrile was removed on rotary evaporator under reduced pressure. The residue was partitioned between ethylacetate(60 mL) and HCl(10%, 25 mL). Organic phase was separated and dried over sodium sulfate, filtered and concentrated to give 1-(3-

cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxycarbonylmethyl pyrazole(1.4 g). ESI mass spectrum m/z (rel. intensity) 298 (M-H, 100).

- 5 Part E 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole.

To a solution of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-hydroxycarbonylmethyl pyrazole(0.20 g, 0.67  
10 mmol) in methylene chloride(20 mL) was added ClCOCOC1(0.84 g, 6.7 mmol) and a drop of DMF. The reaction mixture was stirred at RT overnight. Methylene chloride and excess ClCOCOC1 was removed on rotary evaporator. The residue was redissolved in methylene chloride(20 mL) and to the solution was added 2'-  
15 methylsulfonyl-[1,1']-3-fluoro-4-amino-biphenyl (0.20 g, 0.67 mmol) and DMAP (0.25 g, 2.01 mmol). The mixture was stirred at RT overnight. The next day, methylene chloride was removed on rotary evaporator under reduced pressure. The residue was partitioned between ethyl acetate(30 mL) and Hcl(10%, 20 mL).  
20 Organic phase was separated and washed with water(2 x 20 mL), dried over sodium sulfate, filtered and concentrated to leave 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole(0.32 g). ESI mass spectrum m/z (rel. intensity) 569 (M+Na, 100).

25

Part F 1-(3-aminomethyl-4-fluorophenyl)-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole Trifluoroacetic acid salt.

30 To a solution of 1-(3-cyano-4-fluorophenyl)-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole(50 mg) in ethanol(20 mL) was added palladium(10% on activated carbon, 40 mg). The mixture was hydrogenated at 45 psi overnight. The next day, the reaction  
35 mixture was filtered through celite, filtrate was concentrated and the residue was purified on HPLC(RP gradient) to give 1-(3-aminomethyl-4-fluorophenyl)-3-trifluoromethyl-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-pyrazole( 40

mg) as Trifluoroacetic acid salt. ESI mass spectrum  $z$  (rel. intensity) 551 ( $M+H$ , 100).

Example 244

- 5 Ethyl 1-[3-(aminomethyl)-phenyl]-5-[3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl]aminocarbonylpyrazole-3-carboxylate, trifluoroacetic acid salt.

Part A. Preparation of ethyl 4-(2-furyl)-2,4-dioxobutanoate.

10

To a solution of sodium ethoxide (75 mL of a 21% solution in ethanol, 0.20 mol) in 300 mL of ethanol was added a mixture of 2-acetylfuran (20.0 g, 0.18 mol) and diethyloxalate (26.5 g, 0.18 mol) in 200 mL of tetrahydrofuran over 30 min. The  
15 resulting mixture was allowed to stir at room temperature for 18 h. The reaction mixture was filtered and the solids were washed with ether. The solids were dissolved in water and acidified with 10% HCl. The aqueous was extracted with ethyl acetate and the ethyl acetate layer was washed with brine,  
20 dried ( $MgSO_4$ ) and concentrated in vacuo to afford 21.9 g (57%) of the title compound.  $^1H$ NMR( $CDCl_3$ )  $\delta$ : 7.68 (d, 1H), 7.35 (d, 1H), 6.93 (s, 1H), 6.62 (dd, 1H), 4.39 (q, 2H), 1.40 (t, 3H)ppm.

- 25 Part B. Preparation of ethyl 1-[(3-cyano)phenyl]-5-[fur-2-yl]pyrazole-3-carboxylate.

To a solution of ethyl 4-(2-furyl)-2,4-dioxobutanoate (3.00 g, 14.3 mmol) in 50 mL of absolute ethanol was added 3-  
30 hydrazinobenzonitrile (2.09 g, 15.7 mmol) and *p*-toluenesulfonic acid (2.45 g, 14.3 mmol). This mixture was stirred at 80° C for 2 h. The reaction was concentrated in vacuo and the residue was taken up in ethyl acetate, filtered through a pad of silica gel and concentrated in vacuo. The residue was  
35 recrystallized from hexanes to afford 3.1 g (70%) of the title compound.  $^1H$ NMR( $CDCl_3$ )  $\delta$ : 7.80-7.70 (m, 4H), 7.58 (t, 1H), 7.42 (d, 1H), 7.16 (s, 1H), 6.42 (dd, 1H), 6.24 (d, 1H), 4.45 (q,



2H), 1.42 (t, 3H)ppm. ESI mass spectrum analysis m/z 308.1 (M+H)+.

Part C. Preparation of ethyl 1-[(3-cyano)phenyl]-5-[carboxy]pyrazole-3-carboxylate.

To a solution of ethyl 1-[(3-cyano)phenyl]-5-[fur-2-yl]pyrazole-3-carboxylate ( 1.00 g, 3.25 mmol) in 50 mL of a 2:3:2 mixture of acetonitrile / water / carbon tetrachloride was added sodium periodate (3.13 g, 14.64 mmol) and ruthenium trichloride hydrate (0.015 g, 0.071 mmol). The mixture was stirred at room temperature for 1 h and then was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was triturated with ether to afford 0.9 g (96%) of the title compound. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 8.15 (m, 1H), 7.99 (m, 1H), 7.91 (m, 1H), 7.87 (t, 1H), 7.38 (s, 1H), 4.30 (q, 2H), 1.27 (t, 3H)ppm. ESI mass spectrum analysis: (AP+) m/z 286.1 (M+H)+.

Part D. Preparation of ethyl 1-(3-cyanophenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate.

To a solution ethyl 1-[(3-cyano)phenyl]-5-[carboxy]pyrazole-3-carboxylate ( 0.49 g, 1.72 mmol) in 10 mL of benzene was added oxalyl chloride (0.22 mL, 2.58 mmol) and about 3 drops of dimethylformamide. This solution was allowed to stir at room temperature for 6 h and then was concentrated in vacuo. The residue was dissolved in methylene chloride and then there was added (3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)amine (0.52 g, 1.72 mmol) and 4-dimethylaminopyridine (0.63 g, 5.17 mmol). The resulting mixture was stirred at room temperature for 18 h. The reaction was diluted with ethyl acetate, washed with 10% aq HCl, saturated aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with

2:1 hexanes/ethyl acetate) to afford the 0.70 g (76%) of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.32 (t, 1H), 8.22 (dd, 1H), 8.07 (broad d, 1H), 7.87 (m, 1H), 7.79 (m, 2H), 7.70-7.58 (m, 3H), 7.45 (s, 1H), 7.36 (m, 2H), 7.20 (d, 1H), 4.49 (q, 2H), 2.73 (s, 3H), 1.45 (t, 3H)ppm. ESI mass spectrum analysis m/z 533.2 (M+H)+.

Part E. Preparation of ethyl 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate, trifluoroacetic acid salt.

To a solution of ethyl 1-[(3-cyano)phenyl]-5-[(3-fluoro)-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate (0.20 g, 0.38 mmol) in 100 mL of absolute ethanol was added 2 mL of trifluoroacetic acid and 50 mg of 10% palladium on carbon catalyst. This mixture was stirred under 50 psi of hydrogen on a Parr apparatus for 24 h. The mixture was filtered through a pad of celite and concentrated *in vacuo*. The residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 130 mg (53%) of the title compound as a white powder. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 9.76 (s, 1H), 8.64 (broad s, 3H), 7.94 (d, 1H), 7.67 (m, 1H), 7.50-7.37 (m, 5H), 7.28 (m, 2H), 7.12 (d, 1H), 7.05 (dd, 1H), 6.94 (d, 1H), 4.21 (q, 2H), 3.88 (broad s, 2H), 2.51 (s, 3H), 1.19 (t, 3H)ppm. ESI mass spectrum analysis m/z 537.2 (M+H)+.

#### Example 245

1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylic acid, trifluoroacetic acid salt.

To a solution of ethyl 1-[3-(aminomethyl)-phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate, trifluoroacetic acid salt (0.03 g, 0.05 mmol) in 5 mL of 1:1 ethanol/water was added

potassium hydroxide (0.013 g, 0.23 mmol). This mixture was stirred at room temperature for 3 h and then was acidified by the addition of several drops of trifluoroacetic acid. The reaction was concentrated *in vacuo* and the residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 15 mg (52%) of the title compound as a white powder. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>) δ: 10.60 (s, 1H), 8.19 (broad s, 3H), 8.06 (d, 1H), 7.75 (m, 1H), 7.69-7.51 (m, 5H), 7.50 (m, 2H), 7.39 (d, 1H), 7.34 (dd, 1H), 7.21 (d, 1H), 4.11 (broad s, 2H), 2.90 (s, 3H)ppm. ESI mass spectrum analysis m/z 509.2 (M+H)+.

#### Example 246

1-[3-(aminomethyl)phenyl]-3-[aminocarbonyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

Part A. Preparation of ethyl 1-[3-(N-(tert-butyloxycarbonyl)aminomethyl)-phenyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate.

To a solution of ethyl 1-[3-(aminomethyl)-phenyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate from Example 244 (0.26 g, 0.40 mmol) in 10 mL of methylene chloride was added di-tert-butyl dicarbonate (0.09 g, 0.40 mmol) and 4-dimethylaminopyridine (0.15 g, 1.20 mmol). The resulting mixture was allowed to stir at room temperature for 18 h. The reaction was diluted with ethyl acetate and then was washed with 10% aq HCl, sat'd aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated *in vacuo*. The residue was purified by flash chromatography (elution with 2:1 hexanes/ethyl acetate) to afford the 0.24 g (80%) of the title compound.

<sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 8.28 (t, 1H), 8.14 (d, 1H), 7.89 (broad s, 1H), 7.56 (m, 2H), 7.45-7.35 (m, 4H), 7.30-7.20 (m, 3H), 7.11 (d, 1H), 4.86 (broad s, 1H), 4.40 (q, 2H), 4.33 (m, 2H), 2.65 (s,

3H), 1.40 (t, 3H), 1.37 (s, 9H)ppm. ESI (-ve) mass spectrum analysis m/z (relative intensity) m/z 635.2 (M-H, 100).

Part B. Preparation of 1-[3-(aminomethyl)-phenyl]-3-  
5 [aminocarbonyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole, trifluoroacetic acid salt.

To a solution of ethyl 1-[3-(N-(tert-  
butyloxycarbonyl)aminomethyl)-phenyl]-5-[3-fluoro-(2'-  
10 methylsulfonyl-[1,1']-biphenyl-4-yl)aminocarbonyl]pyrazole-3-  
carboxylate (0.24 g, 0.38 mmol) in 20 mL of 1:1 tetrahydrofuran / water was added potassium hydroxide (0.08 g, 1.5 mmol). The resulting mixture was stirred at 60° C for 1 h and then was cooled and concentrated in vacuo. The residue was diluted with  
15 water and extracted with 1:1 hexane/ethyl acetate. The organics were discarded. The aqueous layer was acidified with aq HCl and extracted with ethyl acetate. The extracts were washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo.  
The residue was dissolved in 10 mL of acetonitrile, cooled to  
20 0° C and then there was added triethylamine (0.10 mL, 0.71 mmol) and iso-butyl chloroformate (0.067 mL, 0.52 mmol). This mixture was allowed to stir for 30 min and then there was added ammonia (0.95 mL of a 2M solution in methanol, 1.88 mmol) and the reaction was allowed to stir with warming to room  
25 temperature for 18 h. The reaction mixture was diluted with ethyl acetate and then was washed with 10% aq HCl, sat'd aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was dissolved in 5 mL of trifluoroacetic acid and stirred at room temperature for 2 h and then was  
30 concentrated in vacuo. The residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 115 mg (40%) of the title compound as a white powder. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 9.53 (s, 1H), 8.78 (broad s, 3H), 8.04 (d, 1H), 7.86 (m, 1H), 7.64 (s, 1H),  
35 7.52 (m, 1H), 7.42 (m, 2H), 7.37 (m, 3H), 7.20 (d, 1H), 7.17 (m, 2H), 7.04 (d, 1H), 6.15 (broad s, 1H), 3.99 (broad s, 2H),

2.60 (s, 3H)ppm. ESI (+ve) mass spectrum analysis m/z (relative intensity) (ESI) m/z 508.2 (M+H, 100).

#### Example 247

5        **Ethyl 1-[3-(aminomethyl)-phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-4-carboxylate, trifluoroacetic acid salt.**

10      **Part A. Preparation of N-(3-cyanophenyl)-trifluoroacetohydrazonoyl bromide.**

To a solution of 3-hydrazinobenzonitrile HCl salt (1.3 g, 7.66 mmol) in 20 mL of absolute ethanol was added  
15      trifluoroacetaldehyde ethyl hemiacetal (1.33 g, 9.19 mmol). The resulting mixture was allowed to stir at 80° C for 18 h and then the reaction was cooled and concentrated in vacuo. The residue was dissolved in 10 mL of dimethylformamide and then there was added N-bromosuccinimide (1.36 g, 7.66 mmol). The  
20      solution was allowed to stir at room temperature for 18h. The reaction was diluted with ethyl acetate, washed with water, sat'd aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo to yield 2.1 g (95%) of the title compound which was sufficiently pure to be used without  
25      purification. <sup>1</sup>HNMR(CDCl<sub>3</sub>): 8.16 (broad s, 1H), 7.47-7.30 (m, 4H)ppm.

**Part B. Preparation of ethyl 3-(2-furyl)-3-oxopropanoate.**

30        To a suspension of hexane-washed sodium hydride (3.5 g of 60% dispersion in mineral oil, 90.8 mmol) in 200 mL of tetrahydrofuran was added diethyl carbonate (10.7 g, 90.8 mmol) and 2-acetylfuran (5.0 g, 45.4 mmol). The resulting mixture was stirred at 70° C for 1h and then was cooled to room  
35      temperature and quenched by the slow addition of 10% aq HCl. The tetrahydrofuran was removed in vacuo and the aqueous was extracted with ethyl acetate. The organics were washed with

water and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo to yield 6.9 g (83%) of the title compound which was sufficiently pure to be used without purification. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.61 (t, 1H), 7.27 (dd, 1H), 6.57 (dd, 1H), 4.20 (q, 2H), 3.84 (s, 2H), 1.25 (t, 3H)ppm.

Part C. Preparation of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[furyl-2-yl]pyrazole-4-carboxylate.

To a solution of ethyl 3-(2-furyl)-3-oxopropanoate (1.87 g, 10.26 mmol) in 20 mL of absolute ethanol was added sodium ethoxide (2.6 mL of a 21% solution in ethanol, 6.84 mmol). Then there was added N-(3-cyanophenyl)-trifluoroacetohydrazonoyl bromide (1.0 g, 3.42 mmol) in absolute ethanol. The resulting mixture was stirred at room temperature for 3 h and then was diluted with ether. The layers were separated and the organics were washed with water, sat'd sodium carbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 4:1 hexanes/ethyl acetate) to afford 0.80 g (63%) of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.71 (m, 1H), 7.60 (m, 1H), 7.53 (m, 2H), 7.44 (d, 1H), 6.95 (d, 1H), 6.55 (dd, 1H), 4.33 (q, 2H), 1.32 (t, 3H)ppm.

Part D. Preparation of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[carboxy]pyrazole-4-carboxylate.

To a solution of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[furyl-2-yl]pyrazole-4-carboxylate (0.75 g, 2.0 mmol) in 30 mL of a 2:3:2 mixture of acetonitrile/water/carbon tetrachloride was added sodium periodate (1.92 g, 9.0 mmol) and ruthenium trichloride hydrate (0.008 g, 0.04 mmol). The mixture was stirred at room temperature for 18 h and then was concentrated in vacuo. The residue was dissolved in ethyl acetate, washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. This residue was dissolved in 1:1 hexanes/ethyl acetate and extracted with sat'd

aq sodium carbonate. The aqueous layer was acidified with HCl and then was extracted with ethyl acetate. These ethyl acetate extracts were washed with brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo to afford 0.40 g (56%) of the title compound which was sufficiently pure to be used without purification.

<sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 7.82 (m, 1H), 7.71 (d, 1H), 7.64 (m, 2H), 4.55 (q, 2H), 1.47 (t, 3H)ppm. ESI (-ve) mass spectrum analysis m/z (relative intensity) 352.1 (M-H, 100).

10 Part E. Preparation of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-4-carboxylate.

To a solution of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[carboxy]pyrazole-4-carboxylate ( 0.33 g, 0.93 mmol) in 10 mL of methylene chloride was added oxalyl chloride (0.12 mL, 1.4 mmol) and about 3 drops of dimethylformamide. This solution was allowed to stir at room temperature for 6 h and then was concentrated in vacuo. The residue was dissolved in methylene chloride and then there was added 4-dimethylaminopyridine (0.34 g, 2.79 mmol) and (2-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)amine hydrochloride (0.28 g, 0.93 mmol). The resulting mixture was stirred at room temperature for 18 h. The reaction was diluted with ethyl acetate, washed with 10% aq HCl, sat'd aq sodium bicarbonate and brine, dried (MgSO<sub>4</sub>) and concentrated in vacuo. The residue was purified by flash chromatography (elution with 2:1 hexanes/ethyl acetate) to afford the 0.25 g (45%) of the title compound. <sup>1</sup>HNMR(CDCl<sub>3</sub>)δ: 11.27 (s, 1H), 8.29 (t, 1H), 8.21 (d, 1H), 7.79 (m, 2H), 7.67-7.52 (m, 4H), 7.40-7.30 (m, 2H), 7.18 (d, 1H), 4.51 (q, 2H), 2.73 (s, 3H), 1.45 (t, 3H)ppm. ESI (+ve) mass spectrum analysis m/z (relative intensity) 623.1 (M+Na, 100).

35 Part F. Preparation of ethyl 1-[3-(aminomethyl)-phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-

yl)aminocarbonyl]pyrazole-4-carboxylate, trifluoroacetic acid salt.

To a solution of ethyl 1-[(3-cyano)phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-4-carboxylate (0.13 g, 0.22 mmol) in 20 mL of absolute ethanol was added conc. HCl (0.018 mL, 0.22 mmol) and 20 mg of 10% palladium on carbon catalyst. This mixture was stirred under 1 atm of hydrogen for 18h. The mixture was filtered through a pad of celite and concentrated in vacuo. The residue was purified by prep HPLC (C18 reverse phase column, elution with a H<sub>2</sub>O/CH<sub>3</sub>CN gradient with 0.5% TFA) and lyophilized to afford 35 mg (21%) of the title compound as a white powder. <sup>1</sup>HNMR(DMSO-d<sub>6</sub>)δ: 11.22 (s, 1H), 8.21 (broad s, 3H), 8.06 (dd, 1H), 7.87 (t, 1H), 7.80-7.40 (m, 6H), 7.38 (m, 2H), 7.22 (dd, 1H), 4.26 (q, 2H), 4.13 (broad q, 2H), 2.91 (s, 3H), 1.14 (t, 3H)ppm. ESI (+ve) mass spectrum analysis m/z (relative intensity) (AP+) 605.2 (M+H, 100).

#### Example 248

1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole, trifluoroacetic acid salt

Part A: Preparation of 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole.

DMF (3 drops) was added to 1-[3-(t-butoxycarbonylaminomethyl)phenyl]-3-(methylthio)pyrazole-5-carboxylic acid (553 mg, 1.5 mmol), and oxalyl chloride (260 μL, 3.0 mmol) in dry CH<sub>2</sub>Cl<sub>2</sub> (30 mL). The resulting solution was stirred at room temperature for 1 h. and evaporated. The resulting solid was redissolved in dry CH<sub>2</sub>Cl<sub>2</sub> (30 mL), and 4-dimethylaminopyridine (585 mg, 4.8 mmol) was added. After stirring 4 min., 4-amino-3-fluoro-2'-methylsulfonyl-[1,1']-



biphenyl, hydrochloride (530 mg, 1.8 mmol) was added portionwise over 5 min., and stirred 22 h. The reaction was extracted once with sat.  $\text{NaHCO}_3$ , then once with a cooled solution of 0.1 M  $\text{HCl}$ . The organic layer was dried over  $\text{MgSO}_4$ , filtered, and evaporated. The crude product was chromatographed on silica gel (40-50%  $\text{EtOAc}$  / hexanes) to yield the desired product (376 mg, 41%).  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 8.38 (bt, 1H), 8.21 (dd, 1H,  $J = 7.7$ ,  $J' = 1.1$ ), 7.81 (bs, 1H), 7.65 (td, 1H,  $J = 7.4$ ,  $J' = 1.4$ ), 7.58 (td, 1H,  $J = 7.7$ ,  $J' = 1.5$ ), 7.43 (m, 4H), 7.32 (m, 2H), 7.17 (d, 1H,  $J = 8.8$ ), 6.84 (s, 1H), 4.90 (bs, 1H), 4.39 (d, 2H,  $J = 6.3$ ), 2.72 (s, 3H), 2.60 (s, 3H), 1.45 (s, 9H)ppm.

Part B: Preparation of 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole.

1-[3-(t-Butoxycarbonylaminoethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole (287 mg, 0.47 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  (5 mL) and TFA (5 mL) and stirred at room temperature for 16h. The reaction was evaporated and purified by prep. HPLC (10-70%  $\text{MeCN}/\text{H}_2\text{O}/0.05\%$  TFA) to yield the desired product (271 mg, 92%).  $^1\text{H NMR}(\text{DMSO}-d_6)\delta$ : 10.60 (s, 1H), 8.25 (bs, 2H), 8.13 (d, 1H,  $J = 8.1$ ), 7.82 (td, 1H,  $J = 7.3$ ,  $J' = 1.5$ ), 7.74 (m, 3H), 7.48 (m, 5H), 7.28 (d, 1H,  $J = 8.4$ ), 7.23 (s, 1H), 4.16 (d, 2H,  $J = 5.8$ ), 2.97 (s, 3H), 2.61 (s, 3H)ppm. APcI mass spectrum analysis  $m/z = 511$  ( $M+H$ , 100).

30

#### Example 249

1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole, trifluoroacetic acid salt

35

MCPBA (110 mg, 57-86%) was added to 1-[3-(t-butoxycarbonylaminoethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole (89 mg, 0.15 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and

stirred at room temperature for 6 h. The reaction was extracted once with sat.  $\text{Na}_2\text{SO}_3$ , then once with sat.  $\text{NaHCO}_3$ . The organic layer was dried over  $\text{Na}_2\text{SO}_4$ , filtered, and evaporated. The crude product was redissolved in  $\text{CH}_2\text{Cl}_2$  (1.5 mL) and TFA (1.5 mL) and stirred at room temperature for 5 h. The resulting solution was evaporated and purified by prep. HPLC (10-70% MeCN/ $\text{H}_2\text{O}$ /0.05% TFA) to yield the desired product.  $^1\text{H}$ NMR( $\text{DMSO}-d_6$ )  $\delta$ : 10.75 (s, 1H), 8.20 (bs, 3H), 8.06 (dd, 1H,  $J = 8.0$ ,  $J' = 1.5$ ), 7.70 (m, 5H), 7.56 (m, 3H), 7.38 (m, 2H), 7.20 (dd,  $J = 8.1$  and 1.7 Hz, 1H), 4.11 (d, 2H,  $J = 5.5$ ), 3.36 (s, 3H), 2.91 (s, 3H) ppm. ESI mass spectrum analysis  $m/z$  (relative intensity) 543 ( $M+H$ , 100).

#### Example 250

1- $[3-(\text{aminomethyl})\text{phenyl}]-5-[(4-(5-(\text{methoxyaminocarbonyl})\text{imidazol-1-yl})\text{phen-1-yl})\text{aminocarbonyl}]-3\text{-trifluoromethylpyrazole, trifluoroacetic acid salt.}$

Part A: A solution of 4-amino-nitrobenzene (5.3 g, 38.4 mmol) in ethyl alcohol (50 mL) was treated with *n*-butyl glyoxylate (10.0 g, 76.9 mmol). After stirring at reflux for 18h, the reaction mixture was concentrated at reduced pressure. The residue was carried to the next step without purification. To the solution of the imine (10.0 g, 40.0 mmol) in methyl alcohol (50 mL) was added potassium carbonate (11.0 g, 80.0 mmol) and tosylmethyl isocyanate (11.7 g, 60.0 mmol). The solution was stirred for 1h at rt, then the solvent was removed under reduced pressure. The residue was treated with saturated sodium chloride solution and the mixture was extracted with methylene chloride. The organic extract was concentrated and triturated with methyl alcohol. The precipitate was collected and dried to afford an imidazole (5.9 g, 59%, 2 steps). Reduction to the aniline was accomplished in MeOH and 10% of Pd/C at 50 psi over 18h. MS (ESI)  $m/z$  (rel. intensity), 216 ( $M^+ + H$ , 100).

Part B: The product from part A was then coupled to 1-(3-cyanophenyl)-3-trifluoromethylpyrazole carboxylic acid via the

acid chloride methodology previously described. The product was purified via silica gel column chromatography (hexane:ethyl acetate, 4:3) to afford pure coupled product. ESI mass spectrum analysis m/z (relative intensity) 481 ( $M^+ + H$ , 100).

5  
Part C: The product from part B (200 mg, 0.4 mmol) in THF (3 mL) was treated with 1N NaOH (0.8 mL, 0.8 mmol). The resultant reaction mixture was stirred for 18h at rt, then acidified to pH 7 with 1N HCl, extracted with ethyl acetate, dried over  
10 magnesium sulfate and concentrated. The resultant acid (100 mg, 0.2 mmol) was dissolved in THF (5 mL), treated with DIEA (0.001 mL, 0.6 mmol), methoxylamine hydrochloride (0.030 g, 0.36 mmol) and TBTU (83 mg, 0.2 mmol) and stirred for 18h at rt. The residue was treated with water and the mixture was  
15 extracted with ethyl acetate, dried over sodium sulphate and concentrated. Purification by silica gel flash chromatography (methanol/methylene chloride, 1:9) afforded the methoxy hydroxamate intermediate (60 mg, 56%). ESI mass spectrum analysis m/z (rel. intensity), 496 ( $M^+ + H$ , 100). Reduction of  
20 the nitrile to the benzylamine was then accomplished via standard conditions.  $^1\text{H NMR}(\text{CD}_3\text{OD})\delta$ : 3.74 (s, 3H), 4.21 (s, 2H), 7.43 (s, 1H), 7.46 (m, 2H), 7.60 (m, 3H), 7.78 (m, 2H), 7.80 (m, 3H) ppm. ESI mass spectrum analysis m/z (relative intensity) 442 ( $M^+ + H$ , 100).

25

#### Example 251

1-(3-aminomethylphenyl)-5-[(4-(5-methyl-1,2,3-triazol-1-yl)phen-1-yl)aminocarbonyl]-3-trifluoromethylpyrazole,  
trifluoroacetic acid salt.


30

Part A: A solution of 4-tert-butyl-[1-(4-nitrophenyl)]5-methyl-1,2,3-tiazol-1-yl-carboxylate (Maybridge Chemical Company, 0.5 g, 1.6 mmol) in TFA (10 mL) was refluxed over 18h. The reaction mixture was concentrated at reduced pressure. The  
35 residue was then reduced to the aniline via standard conditions without purification.  $^1\text{H NMR}(\text{CDCl}_3)\delta$ : 2.36 (s, 3H), 6.83 (d, J =

8.8Hz, 2H), 7.23 (d, J = 8.8Hz, 2H), 7.80 (s, 1H)ppm. ESI mass spectrum analysis m/z (relative intensity) 175 (M<sup>+</sup>+H, 100).

- Part B: The intermediate was then coupled to 1-(3-
- 5 cyanophenyl)-3-trifluoromethylpyrazole carboxylic acid via the acid chloride methodology previously described followed by reduction of the nitrile to the benzylamine and purification via HPLC under reverse phase techniques and lyophilization to afford the title compound as a colorless solid. <sup>1</sup>HNMR(CD<sub>3</sub>OD)δ:
- 10 2.35 (s, 3H), 4.22 (s, 2H), 7.51 (d, J = 9.5 Hz, 2H), 7.55 (s, 1H), 7.60 (m, 3H), 7.65 (s, 1H), 7.71 (s, 1H), 7.89 (d, J = 9.2 Hz, 2H)ppm. ESI mass spectrum analysis m/z (relative intensity) 500 (M<sup>+</sup>+H, 100).


Table 1a



Ex	Ring M	Z	R <sup>1a</sup>	A'	A''	B	MS
1	pyrrole-a	CONH	H	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	460.3
2	pyrrole-a	CONH	H	CH	CH	2-t-Bu-HNSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	516.4
3	pyrrole-a	CONH	Br	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	538.2
4	pyrrole-a	CONH	H	N	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	461.3
5	pyrrole-b	CONH	benzyl	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	550.3
6	pyrrole-b	CONH	benzyl	CH	CH	2-t-Bu-HNSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	606.5
7	imidazole-b	CONH	H	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	461.1
8	imidazole-b	CONH	H	CH	CH	2-t-Bu-HNSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	517.2
9	imidazole-a	CONH	H	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	461.3
10	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	475.2
11	pyrazole	NHCO	CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	475.2
12	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-(5'-CF <sub>3</sub> -tetrazo-1'-yl)C <sub>6</sub> H <sub>4</sub>	532.4
13	4-Cl-pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-t-Bu-NHSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	509.1
14	pyrazole	CONH	CF <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	529.0
15	4-CH <sub>3</sub> O-pyrazole	CONH	CF <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	559.4
16	pyrazole	CONH	CH <sub>3</sub>	CH	CH	1-imidazolyl	386.2
17	pyrazole	CONH	CH <sub>3</sub>	CH	CH	-O-2'-CH <sub>3</sub> SO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	490.3
18	pyrazole	COCH <sub>2</sub>	CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	474.2
19	1,2,3-triazole	CONH	H	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	463.1
20	tetrazole	CONH	-	CH	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	452.2

21	tetrazole	SCH <sub>2</sub>	-	C-Cl	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	500.2
22	tetrazole	SOCH <sub>2</sub>	-	C-Cl	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	516.2
23	tetrazole	SO <sub>2</sub> CH <sub>2</sub>	-	C-Cl	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	532.2
24	tetrazole	CONH	-	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	463.3
25	pyrazole	CONH	CH <sub>3</sub>	N	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	476.3
26	pyrazole	CONH	CH <sub>3</sub>	N	N	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	477.2
27	pyrazole	CONH	CH <sub>3</sub>	C-Cl	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	509.3
28	pyrazole	CONH	CH <sub>3</sub>	C-F	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	493.2
29	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -4-F-C <sub>6</sub> H <sub>3</sub>	493.3
30	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	464.3
31	pyrazole	CONH	CH <sub>3</sub>	C-Cl	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	498.3
32	pyrazole	CONH	CH <sub>3</sub>	C-F	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	482.2
33	pyrazole	CONH	CH <sub>3</sub>	N	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	465.3
34	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-F-C <sub>6</sub> H <sub>4</sub>	414.3
35	pyrazole	CONH	CH <sub>3</sub>	C-Cl	CH	2-F-C <sub>6</sub> H <sub>4</sub>	448.0
36	pyrazole	CONH	CH <sub>3</sub>	CH	CH	2-CH <sub>3</sub> SO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	474.3
37	pyrazole	CONCH <sub>3</sub>	CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	489.3
38	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	517.4
39	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	N	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	518.2
40	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	N	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	506.3
41	pyrazole	CONH	CF <sub>3</sub>	CH	CH	2-CH <sub>3</sub> SO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	528.2
42	pyrazole	CONH	CF <sub>3</sub>	CH	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	518.2
43	4-CH <sub>3</sub> O-pyrazole	CONH	CF <sub>3</sub>	CH	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	548.3
44	pyrazole	CONH	CH <sub>3</sub>	CH	CH	CF <sub>3</sub>	388.2
45	imidazole-a	CONH	4-CH <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	475.3
46	1,2,3-triazole	CONH	H	N	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	463.3
47	1,2,3-triazole	CONH	H	CH	CH	2-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	451.3
48	1,2,4-triazole	CONH	CF <sub>3</sub>	CH	CH	2-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	530.3

Table 1b



5 Unless otherwise indicated, D is at the meta position and is amidino (AM) and R is absent.

Ex	M	Z	R <sub>1a</sub>	A-B	MS
49	pyrazole	CONH	methyl	4-(4'-chlorophenyl)-thiazol-2-yl	437.1
50	pyrazole	CONH	methyl	2'-CF <sub>3</sub> S-biphenyl	496.1
51	pyrazole	CONH	methyl	2'-CF <sub>3</sub> S(O)-biphenyl	512
52	pyrazole	CONH	methyl	2'-CF <sub>3</sub> S(O) <sub>2</sub> -biphenyl	528.1
53	pyrazole	CONH	methyl	4-carboxymethyl-C <sub>6</sub> H <sub>4</sub>	378.2
54	pyrazole	CONH	methyl	4-N,N-(CH <sub>3</sub> ) <sub>2</sub> NC(O)-C <sub>6</sub> H <sub>4</sub>	391
55	pyrazole	CONH	methyl	4-N,N-(CH <sub>3</sub> ) <sub>2</sub> NS(O) <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	426
56	pyrazole	CONH	methyl	4-t-Bu-HNSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	455
57	pyrazole	CONH	methyl	4-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub>	381.3
58	pyrazole	CONH	methyl	4-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub>	388.1
59	pyrazole	CONH	methyl	4-benzylsulfonyl-piperidyl	481.2
60	pyrazole	CONCH <sub>3</sub>	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	489.2
61	pyrazole	CONH	methyl	4'-F-biphenyl	493.1
62	pyrazole	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )-pyridin-2-yl	476.1
63	pyrazole (D= -CN)	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )-pyridin-2-yl	459.1
64	pyrazole	CONH	methyl	2'-CF <sub>3</sub> -biphenyl	464.2
65	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	476.1
66	pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -3-chlorobiphenyl	509.1
67	pyrazole	CONH	methyl	2'-CF <sub>3</sub> -3-chlorobiphenyl	498.1
68	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	517.2
69	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	2'-CF <sub>3</sub> -biphenyl	507.2
70	pyrazole	CONH	C <sub>4</sub> H <sub>9</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	518.2

71	4-CH <sub>3</sub> O-pyrazole	CONH	CF <sub>3</sub>	2'-CF <sub>3</sub> -biphenyl	548.2
72	pyrazole	CONH	CF <sub>3</sub>	2'-CF <sub>3</sub> -biphenyl	518.1
73	pyrazole	CONH	CF <sub>3</sub>	2'-SO <sub>2</sub> CH <sub>3</sub> -biphenyl	528.1
74	pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -3-Br-biphenyl	553.1
75	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -3-Br-biphenyl	554.1
76	pyrazole	COCH <sub>2</sub>	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	474.2
77	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	477.1
78	pyrazole	CONH	CF <sub>3</sub>	5-(2'-t-Bu-HNSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	587.2
79	pyrazole	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	531.1
80	pyrazole (D=CONH <sub>2</sub> )	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	532.1
81	pyrazole (D=-CN)	CONH	CF <sub>3</sub>	4'-imidazol-1-yl-C <sub>6</sub> H <sub>4</sub>	440.1
82	pyrazole	CONH	CF <sub>3</sub>	4'-morpholin-1-yl-C <sub>6</sub> H <sub>4</sub>	459.2
83	pyrazole (D=CONH <sub>2</sub> )	CONH	CF <sub>3</sub>	4'-morpholin-1-yl-C <sub>6</sub> H <sub>4</sub>	460.1
84	pyrazole	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	530.1
85	pyrazole (D=CONH <sub>2</sub> )	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	531.1
86	pyrazole	CONH	CF <sub>3</sub>	4'-(3-methyltetrazol-1-yl)C <sub>6</sub> H <sub>4</sub>	456.2
87	pyrazole	NHSO <sub>2</sub>	methyl	2'-naphthyl	406.1
88	pyrazole	NHSO <sub>2</sub>	methyl	2'-(4-bromo-C <sub>6</sub> H <sub>4</sub> )	434.0
89	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	462.2
90	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	516.1
91	pyrazole	CONH	methyl	5-(2'-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyrid-2-yl	465.2
92	pyrazole	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	477.2
93	pyrazole	CONH	methyl	2'-F-biphenyl	414.2
94	pyrazole	CONH	methyl	3-Cl-2'-F-biphenyl	448.1
95	pyrazole	CONH	methyl	3-F-2'-F-biphenyl	482.2
96	pyrazole	CONH	methyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	493.1
97	pyrazole	CONH	methyl	5-(2'-F-C <sub>6</sub> H <sub>4</sub> )pyrid-2-yl	415.2
98	pyrazole	CONH	methyl	5-(2'-t-Bu-NHSO <sub>2</sub> -phenyl)pyrimidin-2-yl	533.2
99	pyrazole	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )-([1,6]-dihydropyrimidin-2-yl)	579.2



100	pyrazole	CONH	methyl	4-pyrid-3'-yl-C <sub>6</sub> H <sub>4</sub>	379.2
101	pyrazole	CONH	methyl	2-(2'-pyridyl)ethyl	349.2
102	pyrazole	CONH	methyl	3-(C <sub>6</sub> H <sub>4</sub> )propyl	362.2
103	pyrazole	CONH	methyl	4-(pyrid-2'-yl)C <sub>6</sub> H <sub>4</sub>	397.2
104	pyrazole	CONH	methyl	4-(i-propoxy)C <sub>6</sub> H <sub>4</sub>	378.2
105	pyrazole	CONH	methyl	5-(2'-CF <sub>3</sub> -phenyl)pyrimidin-2-yl	466.2
106	pyrazole	CONH	methyl	4-(piperidino-SO <sub>2</sub> )C <sub>6</sub> H <sub>4</sub>	467.2
107	pyrazole	CONH	methyl	4-(piperidino-CO)C <sub>6</sub> H <sub>4</sub>	431.1
108	pyrazole (R=F)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	493
109	pyrazole (D=CONH <sub>2</sub> ) (R=F)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	494.1
110	3-pyrazole	CONH	1-methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	475.3
111	pyrazole	CONH	methyl	4-(pyrazol-4'-yl)C <sub>6</sub> H <sub>4</sub>	386.3
112	pyrazole	CONH	methyl	5-(2'-SO <sub>2</sub> CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyrid-2-yl	475.2
113	pyrazole	CONH	methyl	5-(2'-SO <sub>2</sub> CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimid-2-yl	476.2
114	pyrazole (D= -CN)	CONH	methyl	5-(2'-SO <sub>2</sub> CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimid-2-yl	459.0
115	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	5-(2'-SO <sub>2</sub> CH <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyrimid-2-yl	477.1
116	pyrazole (D= N-NH <sub>2</sub> -AM)	CONH	methyl	2'-t-Bu-NHSO <sub>2</sub> -biphenyl	490.2
117	pyrazole (D= N-NH <sub>2</sub> -AM)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	546.2
118	pyrazole (D=N-Me-N-HO-AM)	CONH	methyl	2'-t-Bu-NHSO <sub>2</sub> -biphenyl	561.2
119	pyrazole (D=N-Me-AM)	CONH	methyl	2'-t-Bu-NHSO <sub>2</sub> -biphenyl	545.2
120	pyrazole (D=N-Me-AM)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	489.2
121	tetrazole	CONH	-	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	464.2
122	tetrazole (D=CONH <sub>2</sub> )	CONH	-	5-(2'-H <sub>2</sub> NSO <sub>2</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	465.1
123	tetrazole	CONH	-	5-(2'-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	453.2
124	tetrazole	CONH	-	4-Br-C <sub>6</sub> H <sub>4</sub>	386.0
125	tetrazole (D=CONH <sub>2</sub> )	CONH	-	5-(2'-CF <sub>3</sub> -C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	454.1

126	tetrazole	CH <sub>2</sub>	-	2'-CF <sub>3</sub> -biphenyl	423.2
127	1-(3-AM-phenyl)-methyl-pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	489
128	1-(4-AM-phenyl)-methyl-pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	489
129	imidazole-a	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	461
130	imidazole-a	CONH	4-methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	475.2
131	imidazole-a	COHN	5-Cl, 4-Me	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	509.1
132	imidazole-c	CONH	2-methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	475.1
133	pyrazole	CONH	methyl	4'-(N-benzimidazol-1-yl)C <sub>6</sub> H <sub>4</sub>	436.2
134	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	4'-(N-benzimidazol-1-yl)C <sub>6</sub> H <sub>4</sub>	437.2
135	pyrazole	CONH	methyl	4'-(2'-methylimidazol-1-yl)C <sub>6</sub> H <sub>4</sub>	400.2
136	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	4'-(2'-methylimidazol-1-yl)C <sub>6</sub> H <sub>4</sub>	401.2
137	pyrazole	CONH	methyl	4'-(1,2,4-triazol-2-yl)C <sub>6</sub> H <sub>4</sub>	387.2
138	pyrazole	CONH	methyl	4'-cyclohexyl-C <sub>6</sub> H <sub>4</sub>	402.2
139	pyrazole	CONH	methyl	biphenyl	396.2
140	pyrazole	CONH	methyl	4'-morpholino-C <sub>6</sub> H <sub>4</sub>	405.2
141	pyrazole	CONH	methyl	4'-(2-CF <sub>3</sub> -tetrazol-1-yl)C <sub>6</sub> H <sub>4</sub>	456.2
142	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	4'-(2-CF <sub>3</sub> -tetrazol-1-yl)C <sub>6</sub> H <sub>4</sub>	443.2
143	pyrazole	CONH	methyl	4-(CH <sub>3</sub> ) <sub>2</sub> NC(O)NH-C <sub>6</sub> H <sub>4</sub>	406.2
144	pyrazole	CONH	methyl	4-(CH <sub>3</sub> ) <sub>2</sub> N-C <sub>6</sub> H <sub>4</sub>	391.2
145	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	4-(CH <sub>3</sub> ) <sub>2</sub> N-C <sub>6</sub> H <sub>4</sub>	392.2
146	pyrazole	CONH	methyl	4-tetrazol-1-yl-C <sub>6</sub> H <sub>4</sub>	388.2
147	pyrazole (D=CONH <sub>2</sub> )	CONH	methyl	4-tetrazol-1-yl-C <sub>6</sub> H <sub>4</sub>	389.2
148	pyrazole	CONH	methyl	4-(N-acetylpiperazin-1-yl)C <sub>6</sub> H <sub>4</sub>	446.2
149	pyrazole	CONH	methyl	4-(N-t-butoxycarbonylpiperazin-1-yl)C <sub>6</sub> H <sub>4</sub>	504.3
150	pyrazole	CONH	methyl	4-(piperazin-1-yl)C <sub>6</sub> H <sub>4</sub>	404.2
151	pyrazole	CONH	CF <sub>3</sub>	4-cyclohexylphenyl	456.2
152	pyrazole	CONH	methyl	4-(N-morpholino)-3-chloro-C <sub>6</sub> H <sub>4</sub>	439.2

153	pyrazole	CONH	CH <sub>3</sub> S	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	507.1
154	pyrazole	CONH	CH <sub>3</sub> SO	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	523.1
155	pyrazole	CONH	CH <sub>3</sub> SO <sub>2</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	539.1
156	tetrazole (D=CONH <sub>2</sub> )	CH <sub>2</sub>	-	2'-CF <sub>3</sub> -biphenyl	424.1
157	tetrazole (D=CONH <sub>2</sub> )	CH <sub>2</sub>	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	435.1
158	pyrazole	CONH	methyl	4-cyclopentyloxyphenyl	404.2
159	pyrazole	CONH	methyl	3-(pyrid-2-yl-NHCH <sub>2</sub> ) C <sub>6</sub> H <sub>4</sub>	426.2
160	pyrazole	CONH	methyl	4-(N-imidazolyl)phenyl	386.2
161	pyrazole	CONH	CF <sub>3</sub>	4-(N-morpholino)-3-Cl- C <sub>6</sub> H <sub>4</sub>	493.1
162	pyrazole	CONH	methyl	4-(N-pyrrolidino- carbonyl)-3-Cl-C <sub>6</sub> H <sub>4</sub>	451.2
163	pyrazole	CONH	methyl	4-(N-morpholino- carbonyl)-3-Cl-C <sub>6</sub> H <sub>4</sub>	433.2
164	pyrazole D= -CN	CONH	CF <sub>3</sub>	4-(N-imidazolyl)phenyl	423.2
165	pyrazole	CONH	CF <sub>3</sub>	4-(N-imidazolyl)phenyl	440.2
166	pyrazole	CONH	CF <sub>3</sub>	4-(N-methyltetrazolon- 1-yl)phenyl	472.1
167	pyrazole (D=CONH <sub>2</sub> )	COCH <sub>2</sub>	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	433.2
168	pyrazole	CONH	methyl	4-N-pyrrolidino- methylphenyl	403.2
169	pyrazole (D= NH <sub>2</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	448.1
170	pyrazole (D= 2-NH <sub>2</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	448.1
171	pyrazole (D= NH <sub>2</sub> ) (R= 4-Cl)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	482.0
172	pyrazole (D= NH <sub>2</sub> ) (R= 4-F)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	466.0
173	pyrazole (D= NH <sub>2</sub> ) (R= 4-OMe)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	478.1
174	tetrazole (D= NH <sub>2</sub> ) (R= 4-Cl)	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	470.0
175	tetrazole (D= NH <sub>2</sub> ) (R= 4-Cl)	CONH	-	5-(2'-H <sub>2</sub> NSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	471.2
176	tetrazole (D= NH <sub>2</sub> ) (R= 4-OMe)	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	466.0
177	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )pyridin-2-yl	463.3

178	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R= 4-CH <sub>3</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	476
179	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R= 4-F)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	480
180	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	4-(N-pyrrolidino- carbonyl)C <sub>6</sub> H <sub>4</sub>	458.2
181	pyrazole (D=*)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	547.2
182	pyrazole (D= **)	CONH	methyl	2'-t-Bu-NHSO <sub>2</sub> -biphenyl	603.2
183	pyrazole (D= **)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	547.2
184	pyrazole (D= ***)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	631.2
185	1-(pyrid- 2-yl)- pyrazole	CONH	methyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	452
186	1-(6-Br- pyrid-2- yl)- pyrazole	CONH	methyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	530
187	tetrazole (D=3-NH <sub>2</sub> ) (R=4-Cl)	CONH	-	3-Cl-2'-H <sub>2</sub> NSO <sub>2</sub> - biphenyl	504.0
188	tetrazole (D=3-NH <sub>2</sub> ) (R=4-Cl)	CONH	-	4-(N-pyrrolidino- carbonyl)C <sub>6</sub> H <sub>4</sub>	430
189	tetrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	450.2
190	1,3,4- triazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	H	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	467.9
191	imidazole- d (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	448.2
192	imidazole- d (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	-	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	447
193	imidazole- d	CONH	-	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	461.2
194	pyrazole (D= CH <sub>2</sub> NHCH <sub>3</sub> )	CONH	methyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	494.1
195	pyrazole (D= CH <sub>2</sub> NHCH <sub>3</sub> )	CONH	methyl	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	492.2
196	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	3-CF <sub>3</sub> 4-OCH <sub>3</sub>	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	545.1
197	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	2-F-4-(N-pyrrolidino- carbonyl)C <sub>6</sub> H <sub>4</sub>	476.2

198	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-F-4-(N-pyrrolidino- carbonyl)C <sub>6</sub> H <sub>4</sub>	476.2
199	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	515.1
200	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	534.1
201	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )[1,6- dihydro]pyrimidin-2-yl	520.1
202	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	5-(2'-H <sub>2</sub> NSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	518.1
203	pyrazole (D=CH(CH <sub>3</sub> )- NH <sub>2</sub> )	CONH	CF <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	530.1
204	pyrazole (D=C(=NH)- N- morpholino)	CONH	CF <sub>3</sub>	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	616.9
205	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CH(OH) CH <sub>2</sub>	CF <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	517.2
206	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	532.9
207	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	5-(2'-H <sub>3</sub> CSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	517.1
208	pyrazole	CONH	CF <sub>3</sub>	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	546
209	pyrazole	CONH	CF <sub>3</sub>	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	547.1
210	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	COCH <sub>2</sub>	CF <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	514.8
211	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CH <sub>2</sub> SO <sub>2</sub> - CH <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	540.1
212	pyrazole	CONH	CH <sub>2</sub> NH- SO <sub>2</sub> CH <sub>3</sub>	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	568.1
213	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CH <sub>2</sub> NH- SO <sub>2</sub> CH <sub>3</sub>	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	572.1
214	pyrazole (D=CH(=NH) NHCO <sub>2</sub> CH <sub>3</sub> )	CONH	methyl	5-(2'-H <sub>2</sub> NSO <sub>2</sub> - C <sub>6</sub> H <sub>4</sub> )pyrimidin-2-yl	535.1
215	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	461.2
216	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-CH <sub>3</sub> -2'-H <sub>3</sub> CSO <sub>2</sub> - biphenyl	530.2
217	1,2,3- triazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	-	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	466.1
218	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R=4-CH <sub>3</sub> )	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	476.2
219	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R=4-F)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	480.2

220	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R=4-Cl)	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	497.1
221	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R=4-F)	CONH	CF <sub>3</sub>	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	551.9
222	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	480
223	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	479
224	pyrazole	CONH	methyl	3-F-4-(N-morpholino)phenyl	423.2
225	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	3-F-4-(N-morpholino)phenyl	410.2
226	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-F-4-(2'-CH <sub>3</sub> -imidazol-1-yl)phenyl	459.2
227	pyrazole (D=CN)	CH <sub>2</sub> O	methyl	biphenyl	420
228	pyrazole	CH <sub>2</sub> O	methyl	biphenyl	437.2
229	pyrazole (D=CONH <sub>2</sub> )	CH <sub>2</sub> O	methyl	biphenyl	438.2
230	pyrazole	CONH	CF <sub>3</sub>	2-F-4-(N-morpholino)phenyl	477.2
231	pyrazole (D=CONH <sub>2</sub> )	CONH	CF <sub>3</sub>	2-F-4-(N-morpholino)phenyl	478.1
232	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	3-CF <sub>3</sub> -4-(N-morpholino)phenyl	514
233	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	ethyl	3-F-2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	493.9
234	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	ethyl	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	493
235	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	ethyl	2-F-4-(2'-H <sub>3</sub> CSO <sub>2</sub> -imidazolyl)phenyl	465.2
236	1-(6-NH <sub>2</sub> CH <sub>2</sub> -pyrid-2-yl)-pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	462.9
237	1-(6-C(=NH <sub>2</sub> )NOH-pyrid-2-yl)-pyrazole	CONH	methyl	2'-t-BuHNSO <sub>2</sub> -biphenyl	548.1
238	1-(6-AM-pyrid-2-yl)-pyrazole	CONH	methyl	2'-H <sub>2</sub> NSO <sub>2</sub> -biphenyl	476.2
239	1-(6-AM-pyrid-2-yl)-pyrazole	CONH	methyl	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	493.9

240	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	2-CH <sub>3</sub> O-4-(N-morpholino)phenyl	422.2
241	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	methyl	4-(3'-CH <sub>3</sub> -5'-oxo-3'-pyrazolin-2'-yl)phenyl	403.1
242	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	SCH <sub>3</sub>	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	493
243	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> ) (R=4-F)	CONH	CF <sub>3</sub>	2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	551
244	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CO <sub>2</sub> Et	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	537.2
245	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	COOH	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	509.2
246	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CONH <sub>2</sub>	2-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	537.2
247	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	3-CF <sub>3</sub> 4-CO <sub>2</sub> Et	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	605.2
248	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	SCH <sub>3</sub>	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	511
249	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	SO <sub>2</sub> CH <sub>3</sub>	3-F-2'-H <sub>3</sub> CSO <sub>2</sub> -biphenyl	543
250	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	4-((5-CH <sub>3</sub> ONHC(O))imidazol-1-yl)phenyl	442
251	pyrazole (D=CH <sub>2</sub> NH <sub>2</sub> )	CONH	CF <sub>3</sub>	4-(5-CH <sub>3</sub> -1,2,3-triazol-1-yl)phenyl	500


\*D=Ethylcarboxyamidino.

\*\*D=1'-imino-1'-N-morpholino)methyl.


\*\*\*D=N-((5-methyl-2-oxo-1,3-dioxol-4-yl)methoxycarbonyl)amidino.

- 5 The following tables contain representative examples of the present invention. Each entry in each table is intended to be paired with each formulae at the start of the table. For example, in Table 2, example 1 is intended to be paired with each of formulae a-nn and in Table 3, example 1 is
- 10 intended to be paired with each of formulae a-nn.


The following groups are intended for group A in the following tables.




2-pyridyl




3-pyridyl




2-pyrimidyl




5-pyrimidyl



2-Cl-phenyl




2-F-phenyl





2,6-diF-phenyl





Table 2














oo




pp




qq




rr




ss



tt



uu



vv

Ex #	R <sup>1a</sup>	A	B
1	CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
2	CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
3	CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
4	CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
5	CH <sub>3</sub>	phenyl	4-morpholino
6	CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
8	CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
9	CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
10	CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
14	CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH <sub>3</sub>	2-pyridyl	4-morpholino
16	CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
18	CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
19	CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
20	CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
24	CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH <sub>3</sub>	3-pyridyl	4-morpholino

26	CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
28	CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
29	CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
30	CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH <sub>3</sub>	2-pyrimidyl	4-morpholino
36	CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
38	CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH <sub>3</sub>	5-pyrimidyl	4-morpholino
46	CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
47	CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
48	CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
56	CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
58	CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH <sub>3</sub>	2-F-phenyl	4-morpholino
66	CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
68	CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
69	CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
70	CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl

74	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
76	CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
82	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
83	CH <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
84	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
85	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
86	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
87	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
88	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
89	CH <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
90	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
94	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
96	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
97	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
98	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
99	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
100	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
104	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
106	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
107	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
108	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
109	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
110	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
116	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
117	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
118	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl

122	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
123	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
126	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
127	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
128	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
133	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
136	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
137	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
138	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
143	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
146	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
147	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
148	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
149	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
150	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
156	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
157	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
162	CF <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
163	CF <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
164	CF <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
165	CF <sub>3</sub>	phenyl	4-morpholino
166	CF <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
167	CF <sub>3</sub>	phenyl	4-morpholinocarbonyl
168	CF <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
169	CF <sub>3</sub>	phenyl	5-methyl-1-imidazolyl



170	CF <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF <sub>3</sub>	2-pyridyl	2-(methylaninosulfonyl)phenyl
173	CF <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
174	CF <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF <sub>3</sub>	2-pyridyl	4-morpholino
176	CF <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
177	CF <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
178	CF <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
179	CF <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
180	CF <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF <sub>3</sub>	3-pyridyl	2-(methylaninosulfonyl)phenyl
183	CF <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
184	CF <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
185	CF <sub>3</sub>	3-pyridyl	4-morpholino
186	CF <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
187	CF <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
188	CF <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
189	CF <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
190	CF <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF <sub>3</sub>	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
193	CF <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF <sub>3</sub>	2-pyrimidyl	4-morpholino
196	CF <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
197	CF <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
198	CF <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF <sub>3</sub>	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
203	CF <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF <sub>3</sub>	5-pyrimidyl	4-morpholino
206	CF <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
207	CF <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
208	CF <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF <sub>3</sub>	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
213	CF <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF <sub>3</sub>	2-Cl-phenyl	4-morpholino
216	CF <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
217	CF <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl

218	CF <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
223	CF <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF <sub>3</sub>	2-F-phenyl	4-morpholino
226	CF <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
227	CF <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
228	CF <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
229	CF <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
230	CF <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
233	CF <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholino
236	CF <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
237	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
242	SCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
243	SCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
244	SCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
245	SCH <sub>3</sub>	phenyl	4-morpholino
246	SCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
247	SCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
248	SCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
249	SCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
250	SCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
253	SCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH <sub>3</sub>	2-pyridyl	4-morpholino
256	SCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
257	SCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
258	SCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
259	SCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
260	SCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
263	SCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH <sub>3</sub>	3-pyridyl	4-morpholino

266	SCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
267	SCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
268	SCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
269	SCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
270	SCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH <sub>3</sub>	2-pyrimidyl	2-(methyaminosulfonyl)phenyl
273	SCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH <sub>3</sub>	2-pyrimidyl	4-morpholino
276	SCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
277	SCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
278	SCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
281	SCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH <sub>3</sub>	5-pyrimidyl	2-(methyaminosulfonyl)phenyl
283	SCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH <sub>3</sub>	5-pyrimidyl	4-morpholino
286	SCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
287	SCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
288	SCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH <sub>3</sub>	2-Cl-phenyl	2-(methyaminosulfonyl)phenyl
293	SCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
296	SCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
297	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH <sub>3</sub>	2-F-phenyl	2-(methyaminosulfonyl)phenyl
303	SCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH <sub>3</sub>	2-F-phenyl	4-morpholino
306	SCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
307	SCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
308	SCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
313	SCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl

314	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
315	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
316	SCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
317	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
322	SOCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
323	SOCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
324	SOCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
325	SOCH <sub>3</sub>	phenyl	4-morpholino
326	SOCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
327	SOCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
328	SOCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
329	SOCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
330	SOCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH <sub>3</sub>	2-pyridyl	4-morpholino
336	SOCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
337	SOCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
338	SOCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH <sub>3</sub>	3-pyridyl	4-morpholino
346	SOCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
347	SOCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
348	SOCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholino
356	SOCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
357	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl

362	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
363	SOCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholino
366	SOCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
367	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
376	SOCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
377	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
383	SOCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH <sub>3</sub>	2-F-phenyl	4-morpholino
386	SOCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
387	SOCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
388	SOCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
393	SOCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
396	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
397	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
402	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
403	SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
404	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
405	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
406	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
407	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
408	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
409	SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl

410	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
413	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
414	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
416	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
417	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
418	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
419	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
420	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
423	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
424	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
425	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
426	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
427	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
428	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
429	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
430	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
436	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
437	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
438	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
446	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
447	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
448	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
456	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
457	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl

458	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
466	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
467	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
468	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
469	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
470	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
473	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
476	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
477	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
482	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
483	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
484	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
485	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
486	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
487	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
488	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
489	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
490	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl

493	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
494	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
496	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
497	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
498	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
499	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
500	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
501	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
503	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
504	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
506	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
507	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
508	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
509	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
510	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
516	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl



517	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
518	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
525	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
526	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
527	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
528	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
536	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
537	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
538	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl

541	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
546	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
547	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
548	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
549	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
550	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
556	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
557	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl

570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl
589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl

625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl
644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl

680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl
699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
722	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
723	CO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
724	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
725	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
726	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
727	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
728	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
729	CO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
730	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl

733	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
734	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
736	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
737	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
738	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
739	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
740	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
744	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
746	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
747	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
748	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
749	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
750	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
756	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
757	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
758	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
766	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
767	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
768	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
776	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
777	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
778	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl

781	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaninosulfonyl)phenyl
783	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
786	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
787	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
788	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
789	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
790	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
793	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
796	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
797	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
798	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
802	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylaninosulfonyl)phenyl
803	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
804	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
805	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholino
806	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
807	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
808	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
809	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
810	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylaninosulfonyl)phenyl
813	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
814	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholino
816	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
817	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
818	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
819	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
820	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylaninosulfonyl)phenyl
823	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
824	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholino
826	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
827	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
828	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl


829	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
830	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholino
836	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
837	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
838	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholino
846	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
847	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
848	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
856	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
857	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
858	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholino
866	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
867	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
868	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
869	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
870	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
876	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl





877	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH <sub>2</sub>	phenyl	2-(aminosulfonyl)phenyl
882	CONH <sub>2</sub>	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH <sub>2</sub>	phenyl	1-pyrrolidinocarbonyl
884	CONH <sub>2</sub>	phenyl	2-(methylsulfonyl)phenyl
885	CONH <sub>2</sub>	phenyl	4-morpholino
886	CONH <sub>2</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
887	CONH <sub>2</sub>	phenyl	4-morpholinocarbonyl
888	CONH <sub>2</sub>	phenyl	2-methyl-1-imidazolyl
889	CONH <sub>2</sub>	phenyl	5-methyl-1-imidazolyl
890	CONH <sub>2</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH <sub>2</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH <sub>2</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH <sub>2</sub>	2-pyridyl	1-pyrrolidinocarbonyl
894	CONH <sub>2</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH <sub>2</sub>	2-pyridyl	4-morpholino
896	CONH <sub>2</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
897	CONH <sub>2</sub>	2-pyridyl	4-morpholinocarbonyl
898	CONH <sub>2</sub>	2-pyridyl	2-methyl-1-imidazolyl
899	CONH <sub>2</sub>	2-pyridyl	5-methyl-1-imidazolyl
900	CONH <sub>2</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH <sub>2</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH <sub>2</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH <sub>2</sub>	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH <sub>2</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH <sub>2</sub>	3-pyridyl	4-morpholino
906	CONH <sub>2</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
907	CONH <sub>2</sub>	3-pyridyl	4-morpholinocarbonyl
908	CONH <sub>2</sub>	3-pyridyl	2-methyl-1-imidazolyl
909	CONH <sub>2</sub>	3-pyridyl	5-methyl-1-imidazolyl
910	CONH <sub>2</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH <sub>2</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH <sub>2</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH <sub>2</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH <sub>2</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH <sub>2</sub>	2-pyrimidyl	4-morpholino
916	CONH <sub>2</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
917	CONH <sub>2</sub>	2-pyrimidyl	4-morpholinocarbonyl
918	CONH <sub>2</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH <sub>2</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH <sub>2</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH <sub>2</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH <sub>2</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH <sub>2</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH <sub>2</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl











925	CONH <sub>2</sub>	5-pyrimidyl	4-morpholino
926	CONH <sub>2</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
927	CONH <sub>2</sub>	5-pyrimidyl	4-morpholinocarbonyl
928	CONH <sub>2</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH <sub>2</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH <sub>2</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH <sub>2</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH <sub>2</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholino
936	CONH <sub>2</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
937	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH <sub>2</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH <sub>2</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH <sub>2</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH <sub>2</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
942	CONH <sub>2</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH <sub>2</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH <sub>2</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH <sub>2</sub>	2-F-phenyl	4-morpholino
946	CONH <sub>2</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
947	CONH <sub>2</sub>	2-F-phenyl	4-morpholinocarbonyl
948	CONH <sub>2</sub>	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH <sub>2</sub>	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH <sub>2</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH <sub>2</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH <sub>2</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholino
956	CONH <sub>2</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
957	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH <sub>2</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH <sub>2</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH <sub>2</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


Table 3














oo




pp




qq




rr




ss



tt



uu




vv


Ex #	A	B
1	phenyl	2-(aminosulfonyl)phenyl
2	phenyl	2-(methylaminosulfonyl)phenyl
3	phenyl	1-pyrrolidinocarbonyl
4	phenyl	2-(methylsulfonyl)phenyl
5	phenyl	4-morpholino
6	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	phenyl	4-morpholinocarbonyl
8	phenyl	2-methyl-1-imidazolyl
9	phenyl	5-methyl-1-imidazolyl
10	phenyl	2-methylsulfonyl-1-imidazolyl
11	2-pyridyl	2-(aminosulfonyl)phenyl
12	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	2-pyridyl	1-pyrrolidinocarbonyl
14	2-pyridyl	2-(methylsulfonyl)phenyl
15	2-pyridyl	4-morpholino
16	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	2-pyridyl	4-morpholinocarbonyl
18	2-pyridyl	2-methyl-1-imidazolyl
19	2-pyridyl	5-methyl-1-imidazolyl
20	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	3-pyridyl	2-(aminosulfonyl)phenyl
22	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	3-pyridyl	1-pyrrolidinocarbonyl
24	3-pyridyl	2-(methylsulfonyl)phenyl
25	3-pyridyl	4-morpholino
26	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	3-pyridyl	4-morpholinocarbonyl
28	3-pyridyl	2-methyl-1-imidazolyl
29	3-pyridyl	5-methyl-1-imidazolyl
30	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	2-pyrimidyl	2-(aminosulfonyl)phenyl


32	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	2-pyrimidyl	1-pyrrolidinocarbonyl
34	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	2-pyrimidyl	4-morpholino
36	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	2-pyrimidyl	4-morpholinocarbonyl
38	2-pyrimidyl	2-methyl-1-imidazolyl
39	2-pyrimidyl	5-methyl-1-imidazolyl
40	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	5-pyrimidyl	1-pyrrolidinocarbonyl
44	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	5-pyrimidyl	4-morpholino
46	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
47	5-pyrimidyl	4-morpholinocarbonyl
48	5-pyrimidyl	2-methyl-1-imidazolyl
49	5-pyrimidyl	5-methyl-1-imidazolyl
50	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	2-Cl-phenyl	4-morpholino
56	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	2-Cl-phenyl	4-morpholinocarbonyl
58	2-Cl-phenyl	2-methyl-1-imidazolyl
59	2-Cl-phenyl	5-methyl-1-imidazolyl
60	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	2-F-phenyl	2-(aminosulfonyl)phenyl
62	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	2-F-phenyl	1-pyrrolidinocarbonyl
64	2-F-phenyl	2-(methylsulfonyl)phenyl
65	2-F-phenyl	4-morpholino
66	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	2-F-phenyl	4-morpholinocarbonyl
68	2-F-phenyl	2-methyl-1-imidazolyl
69	2-F-phenyl	5-methyl-1-imidazolyl
70	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	2,6-diF-phenyl	4-morpholino
76	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	2,6-diF-phenyl	4-morpholinocarbonyl
78	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl




Table 4









Ex #	R <sup>1a</sup>	A	B
1	CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
2	CH <sub>3</sub>	phenyl	2-(methylaninosulfonyl)phenyl
3	CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
4	CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
5	CH <sub>3</sub>	phenyl	4-morpholino
6	CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
8	CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
9	CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
10	CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH <sub>3</sub>	2-pyridyl	2-(methylaninosulfonyl)phenyl
13	CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
14	CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH <sub>3</sub>	2-pyridyl	4-morpholino
16	CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
18	CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
19	CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
20	CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH <sub>3</sub>	3-pyridyl	2-(methylaninosulfonyl)phenyl
23	CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
24	CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH <sub>3</sub>	3-pyridyl	4-morpholino
26	CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
28	CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
29	CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
30	CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH <sub>3</sub>	2-pyrimidyl	2-(methylaninosulfonyl)phenyl
33	CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH <sub>3</sub>	2-pyrimidyl	4-morpholino
36	CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
38	CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH <sub>3</sub>	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
43	CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH <sub>3</sub>	5-pyrimidyl	4-morpholino
46	CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl

47	CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
48	CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
56	CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
58	CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH <sub>3</sub>	2-F-phenyl	4-morpholino
66	CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
68	CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
69	CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
70	CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
76	CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
82	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
83	CH <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
84	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
85	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
86	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
87	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
88	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
89	CH <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
90	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
94	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl

95	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
96	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
97	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
98	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
99	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
100	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
104	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
106	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
107	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
108	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
109	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
110	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
116	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
117	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
118	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
122	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
123	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
126	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
127	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
128	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
133	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
136	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
137	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
138	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl

143	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
146	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
147	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
148	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
149	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
150	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
156	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
157	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
162	CF <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
163	CF <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
164	CF <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
165	CF <sub>3</sub>	phenyl	4-morpholino
166	CF <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
167	CF <sub>3</sub>	phenyl	4-morpholinocarbonyl
168	CF <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
169	CF <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
170	CF <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
174	CF <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF <sub>3</sub>	2-pyridyl	4-morpholino
176	CF <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
177	CF <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
178	CF <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
179	CF <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
180	CF <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
184	CF <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
185	CF <sub>3</sub>	3-pyridyl	4-morpholino
186	CF <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
187	CF <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
188	CF <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
189	CF <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
190	CF <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl



191	CF <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
193	CF <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF <sub>3</sub>	2-pyrimidyl	2-(methysulfonyl)phenyl
195	CF <sub>3</sub>	2-pyrimidyl	4-morpholino
196	CF <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
197	CF <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
198	CF <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF <sub>3</sub>	2-pyrimidyl	2-methysulfonyl-1-imidazolyl
201	CF <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
203	CF <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF <sub>3</sub>	5-pyrimidyl	2-(methysulfonyl)phenyl
205	CF <sub>3</sub>	5-pyrimidyl	4-morpholino
206	CF <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
207	CF <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
208	CF <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF <sub>3</sub>	5-pyrimidyl	2-methysulfonyl-1-imidazolyl
211	CF <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
213	CF <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF <sub>3</sub>	2-Cl-phenyl	2-(methysulfonyl)phenyl
215	CF <sub>3</sub>	2-Cl-phenyl	4-morpholino
216	CF <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
217	CF <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
218	CF <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF <sub>3</sub>	2-Cl-phenyl	2-methysulfonyl-1-imidazolyl
221	CF <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
223	CF <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF <sub>3</sub>	2-F-phenyl	2-(methysulfonyl)phenyl
225	CF <sub>3</sub>	2-F-phenyl	4-morpholino
226	CF <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
227	CF <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
228	CF <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
229	CF <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
230	CF <sub>3</sub>	2-F-phenyl	2-methysulfonyl-1-imidazolyl
231	CF <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
233	CF <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF <sub>3</sub>	2,6-diF-phenyl	2-(methysulfonyl)phenyl
235	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholino
236	CF <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
237	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl

239	CF <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
242	SCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
243	SCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
244	SCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
245	SCH <sub>3</sub>	phenyl	4-morpholino
246	SCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
247	SCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
248	SCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
249	SCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
250	SCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
253	SCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH <sub>3</sub>	2-pyridyl	4-morpholino
256	SCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
257	SCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
258	SCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
259	SCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
260	SCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
263	SCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH <sub>3</sub>	3-pyridyl	4-morpholino
266	SCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
267	SCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
268	SCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
269	SCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
270	SCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
273	SCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH <sub>3</sub>	2-pyrimidyl	4-morpholino
276	SCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
277	SCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
278	SCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
281	SCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
283	SCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH <sub>3</sub>	5-pyrimidyl	4-morpholino
286	SCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl

287	SCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
288	SCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
293	SCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
296	SCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
297	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
303	SCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH <sub>3</sub>	2-F-phenyl	4-morpholino
306	SCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
307	SCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
308	SCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
313	SCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
314	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
315	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
316	SCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
317	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
322	SOCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
323	SOCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
324	SOCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
325	SOCH <sub>3</sub>	phenyl	4-morpholino
326	SOCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
327	SOCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
328	SOCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
329	SOCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
330	SOCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl

335	SOCH <sub>3</sub>	2-pyridyl	4-morpholino
336	SOCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
337	SOCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
338	SOCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH <sub>3</sub>	3-pyridyl	4-morpholino
346	SOCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
347	SOCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
348	SOCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholino
356	SOCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
357	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
362	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
363	SOCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholino
366	SOCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
367	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
376	SOCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
377	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl

383	SOCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH <sub>3</sub>	2-F-phenyl	4-morpholino
386	SOCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
387	SOCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
388	SOCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
393	SOCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
396	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
397	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
402	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
403	SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
404	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
405	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
406	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
407	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
408	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
409	SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
410	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
413	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
414	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
416	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
417	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
418	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
419	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
420	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
423	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
424	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
425	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
426	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
427	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
428	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
429	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
430	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl

431	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
436	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
437	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
438	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
446	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
447	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
448	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
456	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
457	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
458	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
466	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
467	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
468	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
469	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
470	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
473	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
476	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
477	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl

479	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
482	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
483	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
484	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
485	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
486	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
487	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
488	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
489	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
490	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
493	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
494	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
496	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
497	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
498	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
499	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
500	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
501	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
503	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl

504	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
506	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
507	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
508	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
509	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
510	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
516	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
517	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
518	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
525	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
526	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
527	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl



528	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
536	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
537	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
538	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
541	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
546	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
547	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
548	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
549	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
550	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl

552	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
553	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
556	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
557	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methylaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl
570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methylaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methylaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl
589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino

596	Cl	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl
625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl
644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl

651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl
680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl
699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino

706	F	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
722	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
723	CO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
724	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
725	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
726	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
727	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
728	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
729	CO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
730	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
733	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
734	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
736	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
737	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
738	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
739	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
740	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
744	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
746	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
747	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
748	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
749	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
750	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino

756	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
757	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
758	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
766	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
767	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
768	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
776	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
777	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
778	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
781	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
786	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
787	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
788	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
789	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
790	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
796	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
797	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
798	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
802	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
803	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl

804	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
805	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholino
806	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
807	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
808	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
809	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
810	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
814	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholino
816	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
817	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
818	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
819	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
820	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
824	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholino
826	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
827	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
828	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
829	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
830	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholino
836	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
837	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
838	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholino
846	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
847	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
848	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl


852	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methysulfonyl)phenyl
855	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
856	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
857	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
858	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methysulfonyl)phenyl
865	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholino
866	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
867	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
868	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
869	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
870	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methysulfonyl)phenyl
875	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
876	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
877	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH <sub>2</sub>	phenyl	2-(aminosulfonyl)phenyl
882	CONH <sub>2</sub>	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH <sub>2</sub>	phenyl	1-pyrrolidinocarbonyl
884	CONH <sub>2</sub>	phenyl	2-(methysulfonyl)phenyl
885	CONH <sub>2</sub>	phenyl	4-morpholino
886	CONH <sub>2</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
887	CONH <sub>2</sub>	phenyl	4-morpholinocarbonyl
888	CONH <sub>2</sub>	phenyl	2-methyl-1-imidazolyl
889	CONH <sub>2</sub>	phenyl	5-methyl-1-imidazolyl
890	CONH <sub>2</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH <sub>2</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH <sub>2</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH <sub>2</sub>	2-pyridyl	1-pyrrolidinocarbonyl
894	CONH <sub>2</sub>	2-pyridyl	2-(methysulfonyl)phenyl
895	CONH <sub>2</sub>	2-pyridyl	4-morpholino
896	CONH <sub>2</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
897	CONH <sub>2</sub>	2-pyridyl	4-morpholinocarbonyl
898	CONH <sub>2</sub>	2-pyridyl	2-methyl-1-imidazolyl
899	CONH <sub>2</sub>	2-pyridyl	5-methyl-1-imidazolyl



900	CONH <sub>2</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH <sub>2</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH <sub>2</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH <sub>2</sub>	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH <sub>2</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH <sub>2</sub>	3-pyridyl	4-morpholino
906	CONH <sub>2</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
907	CONH <sub>2</sub>	3-pyridyl	4-morpholinocarbonyl
908	CONH <sub>2</sub>	3-pyridyl	2-methyl-1-imidazolyl
909	CONH <sub>2</sub>	3-pyridyl	5-methyl-1-imidazolyl
910	CONH <sub>2</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH <sub>2</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH <sub>2</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH <sub>2</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH <sub>2</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH <sub>2</sub>	2-pyrimidyl	4-morpholino
916	CONH <sub>2</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
917	CONH <sub>2</sub>	2-pyrimidyl	4-morpholinocarbonyl
918	CONH <sub>2</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH <sub>2</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH <sub>2</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH <sub>2</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH <sub>2</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH <sub>2</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH <sub>2</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
925	CONH <sub>2</sub>	5-pyrimidyl	4-morpholino
926	CONH <sub>2</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
927	CONH <sub>2</sub>	5-pyrimidyl	4-morpholinocarbonyl
928	CONH <sub>2</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH <sub>2</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH <sub>2</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH <sub>2</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH <sub>2</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholino
936	CONH <sub>2</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
937	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH <sub>2</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH <sub>2</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH <sub>2</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH <sub>2</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
942	CONH <sub>2</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH <sub>2</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH <sub>2</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH <sub>2</sub>	2-F-phenyl	4-morpholino
946	CONH <sub>2</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
947	CONH <sub>2</sub>	2-F-phenyl	4-morpholinocarbonyl

948	CONH <sub>2</sub>	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH <sub>2</sub>	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH <sub>2</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH <sub>2</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH <sub>2</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholino
956	CONH <sub>2</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
957	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH <sub>2</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH <sub>2</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH <sub>2</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl


Table 5





Ex #	A	B
1	phenyl	2-(aminosulfonyl)phenyl
2	phenyl	2-(methylaminosulfonyl)phenyl
3	phenyl	1-pyrrolidinocarbonyl
4	phenyl	2-(methylsulfonyl)phenyl
5	phenyl	4-morpholino
6	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	phenyl	4-morpholinocarbonyl
8	phenyl	2-methyl-1-imidazolyl
9	phenyl	5-methyl-1-imidazolyl
10	phenyl	2-methylsulfonyl-1-imidazolyl
11	2-pyridyl	2-(aminosulfonyl)phenyl
12	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	2-pyridyl	1-pyrrolidinocarbonyl
14	2-pyridyl	2-(methylsulfonyl)phenyl
15	2-pyridyl	4-morpholino
16	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	2-pyridyl	4-morpholinocarbonyl
18	2-pyridyl	2-methyl-1-imidazolyl
19	2-pyridyl	5-methyl-1-imidazolyl
20	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	3-pyridyl	2-(aminosulfonyl)phenyl
22	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	3-pyridyl	1-pyrrolidinocarbonyl
24	3-pyridyl	2-(methylsulfonyl)phenyl
25	3-pyridyl	4-morpholino
26	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	3-pyridyl	4-morpholinocarbonyl
28	3-pyridyl	2-methyl-1-imidazolyl
29	3-pyridyl	5-methyl-1-imidazolyl
30	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	2-pyrimidyl	1-pyrrolidinocarbonyl
34	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	2-pyrimidyl	4-morpholino
36	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	2-pyrimidyl	4-morpholinocarbonyl
38	2-pyrimidyl	2-methyl-1-imidazolyl
39	2-pyrimidyl	5-methyl-1-imidazolyl
40	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	5-pyrimidyl	2-(methylaminosulfonyl)phenyl

43	5-pyrimidyl	1-pyrrolidinocarbonyl
44	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	5-pyrimidyl	4-morpholino
46	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
47	5-pyrimidyl	4-morpholinocarbonyl
48	5-pyrimidyl	2-methyl-1-imidazolyl
49	5-pyrimidyl	5-methyl-1-imidazolyl
50	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	2-Cl-phenyl	4-morpholino
56	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	2-Cl-phenyl	4-morpholinocarbonyl
58	2-Cl-phenyl	2-methyl-1-imidazolyl
59	2-Cl-phenyl	5-methyl-1-imidazolyl
60	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	2-F-phenyl	2-(aminosulfonyl)phenyl
62	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	2-F-phenyl	1-pyrrolidinocarbonyl
64	2-F-phenyl	2-(methylsulfonyl)phenyl
65	2-F-phenyl	4-morpholino
66	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	2-F-phenyl	4-morpholinocarbonyl
68	2-F-phenyl	2-methyl-1-imidazolyl
69	2-F-phenyl	5-methyl-1-imidazolyl
70	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	2,6-diF-phenyl	4-morpholino
76	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	2,6-diF-phenyl	4-morpholinocarbonyl
78	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

Table 6







For each example, DE is:

- (A) pyridin-4-yl-CH<sub>2</sub>,  
 (B) 2-amino-pyrimidin-4-yl,  
 (C) 6-amino-pyridin-2-yl,  
 (D) 3-amidino-4-F-phenyl, or  
 (E) N-amidino-3-piperidinyl.

5

Ex #	R <sup>1a</sup>	A	B
1	CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
2	CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
3	CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
4	CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
5	CH <sub>3</sub>	phenyl	4-morpholino
6	CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
8	CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
9	CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
10	CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
11	CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
12	CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
14	CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
15	CH <sub>3</sub>	2-pyridyl	4-morpholino
16	CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
18	CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
19	CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
20	CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
22	CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
24	CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
25	CH <sub>3</sub>	3-pyridyl	4-morpholino
26	CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
28	CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
29	CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
30	CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
34	CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	CH <sub>3</sub>	2-pyrimidyl	4-morpholino
36	CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
38	CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
39	CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
40	CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl



41	CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	CH <sub>3</sub>	5-pyrimidyl	2-(methylaninosulfonyl)phenyl
43	CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
44	CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	CH <sub>3</sub>	5-pyrimidyl	4-morpholino
46	CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
47	CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
48	CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
49	CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
50	CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	CH <sub>3</sub>	2-Cl-phenyl	2-(methylaninosulfonyl)phenyl
53	CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
56	CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
58	CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
59	CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
60	CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
62	CH <sub>3</sub>	2-F-phenyl	2-(methylaninosulfonyl)phenyl
63	CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
64	CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
65	CH <sub>3</sub>	2-F-phenyl	4-morpholino
66	CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
68	CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
69	CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
70	CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaninosulfonyl)phenyl
73	CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
76	CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
78	CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
81	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
82	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaninosulfonyl)phenyl
83	CH <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
84	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
85	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
86	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
87	CH <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
88	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl

89	CH <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
90	CH <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
91	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
92	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
93	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
94	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
95	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
96	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
97	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
98	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
99	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
100	CH <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
101	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
102	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
103	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
104	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
105	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
106	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
107	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
108	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
109	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
110	CH <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
111	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
112	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
113	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
114	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
115	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
116	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
117	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
118	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
119	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
120	CH <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
121	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
122	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
123	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
124	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
125	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
126	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
127	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
128	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
129	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
130	CH <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
131	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
132	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
133	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
134	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
135	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
136	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl

137	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
138	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
139	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
140	CH <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
141	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
142	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
143	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
144	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
145	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
146	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
147	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
148	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
149	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
150	CH <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
151	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
152	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
153	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
154	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
155	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
156	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
157	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
158	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
159	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
160	CH <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
161	CF <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
162	CF <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
163	CF <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
164	CF <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
165	CF <sub>3</sub>	phenyl	4-morpholino
166	CF <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
167	CF <sub>3</sub>	phenyl	4-morpholinocarbonyl
168	CF <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
169	CF <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
170	CF <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
171	CF <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
172	CF <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
173	CF <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
174	CF <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
175	CF <sub>3</sub>	2-pyridyl	4-morpholino
176	CF <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
177	CF <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
178	CF <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
179	CF <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
180	CF <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
181	CF <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
182	CF <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
183	CF <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
184	CF <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl

185	CF <sub>3</sub>	3-pyridyl	4-morpholino
186	CF <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
187	CF <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
188	CF <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
189	CF <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
190	CF <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
191	CF <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
192	CF <sub>3</sub>	2-pyrimidyl	2-(methyaminosulfonyl)phenyl
193	CF <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
194	CF <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
195	CF <sub>3</sub>	2-pyrimidyl	4-morpholino
196	CF <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
197	CF <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
198	CF <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
199	CF <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
200	CF <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
201	CF <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
202	CF <sub>3</sub>	5-pyrimidyl	2-(methyaminosulfonyl)phenyl
203	CF <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
204	CF <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
205	CF <sub>3</sub>	5-pyrimidyl	4-morpholino
206	CF <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
207	CF <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
208	CF <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
209	CF <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
210	CF <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
211	CF <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
212	CF <sub>3</sub>	2-Cl-phenyl	2-(methyaminosulfonyl)phenyl
213	CF <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
214	CF <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
215	CF <sub>3</sub>	2-Cl-phenyl	4-morpholino
216	CF <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
217	CF <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
218	CF <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
219	CF <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
220	CF <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
221	CF <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
222	CF <sub>3</sub>	2-F-phenyl	2-(methyaminosulfonyl)phenyl
223	CF <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
224	CF <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
225	CF <sub>3</sub>	2-F-phenyl	4-morpholino
226	CF <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
227	CF <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
228	CF <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
229	CF <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
230	CF <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
231	CF <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
232	CF <sub>3</sub>	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl

233	CF <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
234	CF <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
235	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholino
236	CF <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
237	CF <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
238	CF <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
239	CF <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
240	CF <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
241	SCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
242	SCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
243	SCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
244	SCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
245	SCH <sub>3</sub>	phenyl	4-morpholino
246	SCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
247	SCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
248	SCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
249	SCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
250	SCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
251	SCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
252	SCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
253	SCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
254	SCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
255	SCH <sub>3</sub>	2-pyridyl	4-morpholino
256	SCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
257	SCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
258	SCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
259	SCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
260	SCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
261	SCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
262	SCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
263	SCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
264	SCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
265	SCH <sub>3</sub>	3-pyridyl	4-morpholino
266	SCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
267	SCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
268	SCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
269	SCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
270	SCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
271	SCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
272	SCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
273	SCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
274	SCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
275	SCH <sub>3</sub>	2-pyrimidyl	4-morpholino
276	SCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
277	SCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
278	SCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
279	SCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
280	SCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl

281	SCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
282	SCH <sub>3</sub>	5-pyrimidyl	2-(methyaminosulfonyl)phenyl
283	SCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
284	SCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
285	SCH <sub>3</sub>	5-pyrimidyl	4-morpholino
286	SCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
287	SCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
288	SCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
289	SCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
290	SCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
291	SCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
292	SCH <sub>3</sub>	2-Cl-phenyl	2-(methyaminosulfonyl)phenyl
293	SCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
294	SCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
295	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
296	SCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
297	SCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
298	SCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
299	SCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
300	SCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
301	SCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
302	SCH <sub>3</sub>	2-F-phenyl	2-(methyaminosulfonyl)phenyl
303	SCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
304	SCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
305	SCH <sub>3</sub>	2-F-phenyl	4-morpholino
306	SCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
307	SCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
308	SCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
309	SCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
310	SCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
311	SCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
312	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
313	SCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
314	SCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
315	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
316	SCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
317	SCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
318	SCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
319	SCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
320	SCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
321	SOCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
322	SOCH <sub>3</sub>	phenyl	2-(methyaminosulfonyl)phenyl
323	SOCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
324	SOCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
325	SOCH <sub>3</sub>	phenyl	4-morpholino
326	SOCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
327	SOCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
328	SOCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl

329	SOCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
330	SOCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
331	SOCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
332	SOCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
333	SOCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
334	SOCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
335	SOCH <sub>3</sub>	2-pyridyl	4-morpholino
336	SOCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
337	SOCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
338	SOCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
339	SOCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
340	SOCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
341	SOCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
342	SOCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
343	SOCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
344	SOCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
345	SOCH <sub>3</sub>	3-pyridyl	4-morpholino
346	SOCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
347	SOCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
348	SOCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
349	SOCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
350	SOCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
351	SOCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
352	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
353	SOCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
354	SOCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
355	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholino
356	SOCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
357	SOCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
358	SOCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
359	SOCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
360	SOCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
361	SOCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
362	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
363	SOCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
364	SOCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
365	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholino
366	SOCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
367	SOCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
368	SOCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
369	SOCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
370	SOCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
371	SOCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
372	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
373	SOCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
374	SOCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
375	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
376	SOCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl

377	SOCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
378	SOCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
379	SOCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
380	SOCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
381	SOCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
382	SOCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
383	SOCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
384	SOCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
385	SOCH <sub>3</sub>	2-F-phenyl	4-morpholino
386	SOCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
387	SOCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
388	SOCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
389	SOCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
390	SOCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
391	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
392	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
393	SOCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
394	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
395	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
396	SOCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
397	SOCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
398	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
399	SOCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
400	SOCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
401	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
402	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
403	SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
404	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
405	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
406	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
407	SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
408	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
409	SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
410	SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
411	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
412	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
413	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
414	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
415	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
416	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
417	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
418	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
419	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
420	SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
421	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
422	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
423	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
424	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl



425	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
426	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
427	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
428	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
429	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
430	SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
431	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
432	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
433	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
434	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
435	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
436	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
437	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
438	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
439	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
440	SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
441	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
442	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
443	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
444	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
445	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
446	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
447	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
448	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
449	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
450	SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
451	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
452	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
453	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
454	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
455	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
456	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
457	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
458	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
459	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
460	SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
461	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
462	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
463	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
464	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
465	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
466	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
467	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
468	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
469	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
470	SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
471	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
472	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl

473	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
474	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
475	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
476	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
477	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
478	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
479	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
480	SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
481	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
482	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
483	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
484	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
485	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
486	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
487	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
488	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
489	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
490	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
491	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
492	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
493	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
494	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
495	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
496	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
497	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
498	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
499	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
500	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl

501	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
502	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
503	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
504	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
505	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
506	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
507	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
508	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
509	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
510	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
511	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
512	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
513	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
514	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
515	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
516	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
517	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
518	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
519	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
520	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
521	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
522	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
523	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
524	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl

525	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
526	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
527	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
528	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
529	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
530	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
531	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
532	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
533	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
534	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
535	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
536	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
537	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
538	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
539	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
540	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
541	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
542	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
543	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
544	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
545	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
546	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
547	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
548	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl

549	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
550	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
551	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
552	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methyaminosulfonyl)phenyl
553	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
554	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
555	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
556	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
557	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
558	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
559	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
560	CH <sub>2</sub> NH-SO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
561	Cl	phenyl	2-(aminosulfonyl)phenyl
562	Cl	phenyl	2-(methyaminosulfonyl)phenyl
563	Cl	phenyl	1-pyrrolidinocarbonyl
564	Cl	phenyl	2-(methylsulfonyl)phenyl
565	Cl	phenyl	4-morpholino
566	Cl	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
567	Cl	phenyl	4-morpholinocarbonyl
568	Cl	phenyl	2-methyl-1-imidazolyl
569	Cl	phenyl	5-methyl-1-imidazolyl
570	Cl	phenyl	2-methylsulfonyl-1-imidazolyl
571	Cl	2-pyridyl	2-(aminosulfonyl)phenyl
572	Cl	2-pyridyl	2-(methyaminosulfonyl)phenyl
573	Cl	2-pyridyl	1-pyrrolidinocarbonyl
574	Cl	2-pyridyl	2-(methylsulfonyl)phenyl
575	Cl	2-pyridyl	4-morpholino
576	Cl	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
577	Cl	2-pyridyl	4-morpholinocarbonyl
578	Cl	2-pyridyl	2-methyl-1-imidazolyl
579	Cl	2-pyridyl	5-methyl-1-imidazolyl
580	Cl	2-pyridyl	2-methylsulfonyl-1-imidazolyl
581	Cl	3-pyridyl	2-(aminosulfonyl)phenyl
582	Cl	3-pyridyl	2-(methyaminosulfonyl)phenyl
583	Cl	3-pyridyl	1-pyrrolidinocarbonyl
584	Cl	3-pyridyl	2-(methylsulfonyl)phenyl
585	Cl	3-pyridyl	4-morpholino
586	Cl	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
587	Cl	3-pyridyl	4-morpholinocarbonyl
588	Cl	3-pyridyl	2-methyl-1-imidazolyl

589	Cl	3-pyridyl	5-methyl-1-imidazolyl
590	Cl	3-pyridyl	2-methylsulfonyl-1-imidazolyl
591	Cl	2-pyrimidyl	2-(aminosulfonyl)phenyl
592	Cl	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
593	Cl	2-pyrimidyl	1-pyrrolidinocarbonyl
594	Cl	2-pyrimidyl	2-(methylsulfonyl)phenyl
595	Cl	2-pyrimidyl	4-morpholino
596	Cl	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
597	Cl	2-pyrimidyl	4-morpholinocarbonyl
598	Cl	2-pyrimidyl	2-methyl-1-imidazolyl
599	Cl	2-pyrimidyl	5-methyl-1-imidazolyl
600	Cl	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
601	Cl	5-pyrimidyl	2-(aminosulfonyl)phenyl
602	Cl	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
603	Cl	5-pyrimidyl	1-pyrrolidinocarbonyl
604	Cl	5-pyrimidyl	2-(methylsulfonyl)phenyl
605	Cl	5-pyrimidyl	4-morpholino
606	Cl	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
607	Cl	5-pyrimidyl	4-morpholinocarbonyl
608	Cl	5-pyrimidyl	2-methyl-1-imidazolyl
609	Cl	5-pyrimidyl	5-methyl-1-imidazolyl
610	Cl	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
611	Cl	2-Cl-phenyl	2-(aminosulfonyl)phenyl
612	Cl	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
613	Cl	2-Cl-phenyl	1-pyrrolidinocarbonyl
614	Cl	2-Cl-phenyl	2-(methylsulfonyl)phenyl
615	Cl	2-Cl-phenyl	4-morpholino
616	Cl	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
617	Cl	2-Cl-phenyl	4-morpholinocarbonyl
618	Cl	2-Cl-phenyl	2-methyl-1-imidazolyl
619	Cl	2-Cl-phenyl	5-methyl-1-imidazolyl
620	Cl	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
621	Cl	2-F-phenyl	2-(aminosulfonyl)phenyl
622	Cl	2-F-phenyl	2-(methylaminosulfonyl)phenyl
623	Cl	2-F-phenyl	1-pyrrolidinocarbonyl
624	Cl	2-F-phenyl	2-(methylsulfonyl)phenyl
625	Cl	2-F-phenyl	4-morpholino
626	Cl	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
627	Cl	2-F-phenyl	4-morpholinocarbonyl
628	Cl	2-F-phenyl	2-methyl-1-imidazolyl
629	Cl	2-F-phenyl	5-methyl-1-imidazolyl
630	Cl	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
631	Cl	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
632	Cl	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
633	Cl	2,6-diF-phenyl	1-pyrrolidinocarbonyl
634	Cl	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
635	Cl	2,6-diF-phenyl	4-morpholino
636	Cl	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
637	Cl	2,6-diF-phenyl	4-morpholinocarbonyl
638	Cl	2,6-diF-phenyl	2-methyl-1-imidazolyl
639	Cl	2,6-diF-phenyl	5-methyl-1-imidazolyl
640	Cl	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
641	F	phenyl	2-(aminosulfonyl)phenyl
642	F	phenyl	2-(methylaminosulfonyl)phenyl
643	F	phenyl	1-pyrrolidinocarbonyl

644	F	phenyl	2-(methylsulfonyl)phenyl
645	F	phenyl	4-morpholino
646	F	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
647	F	phenyl	4-morpholinocarbonyl
648	F	phenyl	2-methyl-1-imidazolyl
649	F	phenyl	5-methyl-1-imidazolyl
650	F	phenyl	2-methylsulfonyl-1-imidazolyl
651	F	2-pyridyl	2-(aminosulfonyl)phenyl
652	F	2-pyridyl	2-(methylaminosulfonyl)phenyl
653	F	2-pyridyl	1-pyrrolidinocarbonyl
654	F	2-pyridyl	2-(methylsulfonyl)phenyl
655	F	2-pyridyl	4-morpholino
656	F	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
657	F	2-pyridyl	4-morpholinocarbonyl
658	F	2-pyridyl	2-methyl-1-imidazolyl
659	F	2-pyridyl	5-methyl-1-imidazolyl
660	F	2-pyridyl	2-methylsulfonyl-1-imidazolyl
661	F	3-pyridyl	2-(aminosulfonyl)phenyl
662	F	3-pyridyl	2-(methylaminosulfonyl)phenyl
663	F	3-pyridyl	1-pyrrolidinocarbonyl
664	F	3-pyridyl	2-(methylsulfonyl)phenyl
665	F	3-pyridyl	4-morpholino
666	F	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
667	F	3-pyridyl	4-morpholinocarbonyl
668	F	3-pyridyl	2-methyl-1-imidazolyl
669	F	3-pyridyl	5-methyl-1-imidazolyl
670	F	3-pyridyl	2-methylsulfonyl-1-imidazolyl
671	F	2-pyrimidyl	2-(aminosulfonyl)phenyl
672	F	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
673	F	2-pyrimidyl	1-pyrrolidinocarbonyl
674	F	2-pyrimidyl	2-(methylsulfonyl)phenyl
675	F	2-pyrimidyl	4-morpholino
676	F	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
677	F	2-pyrimidyl	4-morpholinocarbonyl
678	F	2-pyrimidyl	2-methyl-1-imidazolyl
679	F	2-pyrimidyl	5-methyl-1-imidazolyl
680	F	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
681	F	5-pyrimidyl	2-(aminosulfonyl)phenyl
682	F	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
683	F	5-pyrimidyl	1-pyrrolidinocarbonyl
684	F	5-pyrimidyl	2-(methylsulfonyl)phenyl
685	F	5-pyrimidyl	4-morpholino
686	F	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
687	F	5-pyrimidyl	4-morpholinocarbonyl
688	F	5-pyrimidyl	2-methyl-1-imidazolyl
689	F	5-pyrimidyl	5-methyl-1-imidazolyl
690	F	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
691	F	2-Cl-phenyl	2-(aminosulfonyl)phenyl
692	F	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
693	F	2-Cl-phenyl	1-pyrrolidinocarbonyl
694	F	2-Cl-phenyl	2-(methylsulfonyl)phenyl
695	F	2-Cl-phenyl	4-morpholino
696	F	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
697	F	2-Cl-phenyl	4-morpholinocarbonyl
698	F	2-Cl-phenyl	2-methyl-1-imidazolyl

699	F	2-Cl-phenyl	5-methyl-1-imidazolyl
700	F	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
701	F	2-F-phenyl	2-(aminosulfonyl)phenyl
702	F	2-F-phenyl	2-(methylaminosulfonyl)phenyl
703	F	2-F-phenyl	1-pyrrolidinocarbonyl
704	F	2-F-phenyl	2-(methylsulfonyl)phenyl
705	F	2-F-phenyl	4-morpholino
706	F	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
707	F	2-F-phenyl	4-morpholinocarbonyl
708	F	2-F-phenyl	2-methyl-1-imidazolyl
709	F	2-F-phenyl	5-methyl-1-imidazolyl
710	F	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
711	F	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
712	F	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
713	F	2,6-diF-phenyl	1-pyrrolidinocarbonyl
714	F	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
715	F	2,6-diF-phenyl	4-morpholino
716	F	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
717	F	2,6-diF-phenyl	4-morpholinocarbonyl
718	F	2,6-diF-phenyl	2-methyl-1-imidazolyl
719	F	2,6-diF-phenyl	5-methyl-1-imidazolyl
720	F	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
721	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
722	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
723	CO <sub>2</sub> CH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
724	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
725	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholino
726	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
727	CO <sub>2</sub> CH <sub>3</sub>	phenyl	4-morpholinocarbonyl
728	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
729	CO <sub>2</sub> CH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
730	CO <sub>2</sub> CH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
731	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
732	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
733	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
734	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
735	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholino
736	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
737	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
738	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
739	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
740	CO <sub>2</sub> CH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
741	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
742	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
743	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
744	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
745	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholino
746	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
747	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
748	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
749	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl



750	CO <sub>2</sub> CH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
751	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
752	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
753	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
754	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
755	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholino
756	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
757	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
758	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
759	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
760	CO <sub>2</sub> CH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
761	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
762	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
763	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
764	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
765	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholino
766	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
767	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
768	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
769	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
770	CO <sub>2</sub> CH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
771	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
772	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
773	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
774	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
775	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholino
776	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
777	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
778	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
779	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
780	CO <sub>2</sub> CH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
781	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
782	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
783	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
784	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
785	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholino
786	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
787	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
788	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
789	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
790	CO <sub>2</sub> CH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
791	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
792	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
793	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
794	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
795	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
796	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
797	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl


798	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
799	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
800	CO <sub>2</sub> CH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
801	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(aminosulfonyl)phenyl
802	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylaminosulfonyl)phenyl
803	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	1-pyrrolidinocarbonyl
804	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(methylsulfonyl)phenyl
805	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholino
806	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
807	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	4-morpholinocarbonyl
808	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methyl-1-imidazolyl
809	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	5-methyl-1-imidazolyl
810	CH <sub>2</sub> OCH <sub>3</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
811	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
812	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
813	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	1-pyrrolidinocarbonyl
814	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
815	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholino
816	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
817	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	4-morpholinocarbonyl
818	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methyl-1-imidazolyl
819	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	5-methyl-1-imidazolyl
820	CH <sub>2</sub> OCH <sub>3</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
821	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
822	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
823	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	1-pyrrolidinocarbonyl
824	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
825	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholino
826	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
827	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	4-morpholinocarbonyl
828	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methyl-1-imidazolyl
829	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	5-methyl-1-imidazolyl
830	CH <sub>2</sub> OCH <sub>3</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
831	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
832	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
833	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
834	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
835	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholino
836	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
837	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	4-morpholinocarbonyl
838	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
839	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
840	CH <sub>2</sub> OCH <sub>3</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
841	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
842	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
843	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
844	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
845	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholino

846	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
847	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	4-morpholinocarbonyl
848	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
849	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
850	CH <sub>2</sub> OCH <sub>3</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
851	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
852	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
853	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
854	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
855	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholino
856	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
857	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	4-morpholinocarbonyl
858	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
859	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
860	CH <sub>2</sub> OCH <sub>3</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
861	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl
862	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
863	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
864	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
865	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholino
866	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
867	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	4-morpholinocarbonyl
868	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methyl-1-imidazolyl
869	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	5-methyl-1-imidazolyl
870	CH <sub>2</sub> OCH <sub>3</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
871	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
872	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
873	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
874	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
875	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholino
876	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
877	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
878	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
879	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
880	CH <sub>2</sub> OCH <sub>3</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl
881	CONH <sub>2</sub>	phenyl	2-(aminosulfonyl)phenyl
882	CONH <sub>2</sub>	phenyl	2-(methylaminosulfonyl)phenyl
883	CONH <sub>2</sub>	phenyl	1-pyrrolidinocarbonyl
884	CONH <sub>2</sub>	phenyl	2-(methylsulfonyl)phenyl
885	CONH <sub>2</sub>	phenyl	4-morpholino
886	CONH <sub>2</sub>	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
887	CONH <sub>2</sub>	phenyl	4-morpholinocarbonyl
888	CONH <sub>2</sub>	phenyl	2-methyl-1-imidazolyl
889	CONH <sub>2</sub>	phenyl	5-methyl-1-imidazolyl
890	CONH <sub>2</sub>	phenyl	2-methylsulfonyl-1-imidazolyl
891	CONH <sub>2</sub>	2-pyridyl	2-(aminosulfonyl)phenyl
892	CONH <sub>2</sub>	2-pyridyl	2-(methylaminosulfonyl)phenyl
893	CONH <sub>2</sub>	2-pyridyl	1-pyrrolidinocarbonyl

894	CONH <sub>2</sub>	2-pyridyl	2-(methylsulfonyl)phenyl
895	CONH <sub>2</sub>	2-pyridyl	4-morpholino
896	CONH <sub>2</sub>	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
897	CONH <sub>2</sub>	2-pyridyl	4-morpholinocarbonyl
898	CONH <sub>2</sub>	2-pyridyl	2-methyl-1-imidazolyl
899	CONH <sub>2</sub>	2-pyridyl	5-methyl-1-imidazolyl
900	CONH <sub>2</sub>	2-pyridyl	2-methylsulfonyl-1-imidazolyl
901	CONH <sub>2</sub>	3-pyridyl	2-(aminosulfonyl)phenyl
902	CONH <sub>2</sub>	3-pyridyl	2-(methylaminosulfonyl)phenyl
903	CONH <sub>2</sub>	3-pyridyl	1-pyrrolidinocarbonyl
904	CONH <sub>2</sub>	3-pyridyl	2-(methylsulfonyl)phenyl
905	CONH <sub>2</sub>	3-pyridyl	4-morpholino
906	CONH <sub>2</sub>	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
907	CONH <sub>2</sub>	3-pyridyl	4-morpholinocarbonyl
908	CONH <sub>2</sub>	3-pyridyl	2-methyl-1-imidazolyl
909	CONH <sub>2</sub>	3-pyridyl	5-methyl-1-imidazolyl
910	CONH <sub>2</sub>	3-pyridyl	2-methylsulfonyl-1-imidazolyl
911	CONH <sub>2</sub>	2-pyrimidyl	2-(aminosulfonyl)phenyl
912	CONH <sub>2</sub>	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
913	CONH <sub>2</sub>	2-pyrimidyl	1-pyrrolidinocarbonyl
914	CONH <sub>2</sub>	2-pyrimidyl	2-(methylsulfonyl)phenyl
915	CONH <sub>2</sub>	2-pyrimidyl	4-morpholino
916	CONH <sub>2</sub>	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
917	CONH <sub>2</sub>	2-pyrimidyl	4-morpholinocarbonyl
918	CONH <sub>2</sub>	2-pyrimidyl	2-methyl-1-imidazolyl
919	CONH <sub>2</sub>	2-pyrimidyl	5-methyl-1-imidazolyl
920	CONH <sub>2</sub>	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
921	CONH <sub>2</sub>	5-pyrimidyl	2-(aminosulfonyl)phenyl
922	CONH <sub>2</sub>	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
923	CONH <sub>2</sub>	5-pyrimidyl	1-pyrrolidinocarbonyl
924	CONH <sub>2</sub>	5-pyrimidyl	2-(methylsulfonyl)phenyl
925	CONH <sub>2</sub>	5-pyrimidyl	4-morpholino
926	CONH <sub>2</sub>	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
927	CONH <sub>2</sub>	5-pyrimidyl	4-morpholinocarbonyl
928	CONH <sub>2</sub>	5-pyrimidyl	2-methyl-1-imidazolyl
929	CONH <sub>2</sub>	5-pyrimidyl	5-methyl-1-imidazolyl
930	CONH <sub>2</sub>	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
931	CONH <sub>2</sub>	2-Cl-phenyl	2-(aminosulfonyl)phenyl
932	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
933	CONH <sub>2</sub>	2-Cl-phenyl	1-pyrrolidinocarbonyl
934	CONH <sub>2</sub>	2-Cl-phenyl	2-(methylsulfonyl)phenyl
935	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholino
936	CONH <sub>2</sub>	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
937	CONH <sub>2</sub>	2-Cl-phenyl	4-morpholinocarbonyl
938	CONH <sub>2</sub>	2-Cl-phenyl	2-methyl-1-imidazolyl
939	CONH <sub>2</sub>	2-Cl-phenyl	5-methyl-1-imidazolyl
940	CONH <sub>2</sub>	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
941	CONH <sub>2</sub>	2-F-phenyl	2-(aminosulfonyl)phenyl

942	CONH <sub>2</sub>	2-F-phenyl	2-(methylaminosulfonyl)phenyl
943	CONH <sub>2</sub>	2-F-phenyl	1-pyrrolidinocarbonyl
944	CONH <sub>2</sub>	2-F-phenyl	2-(methylsulfonyl)phenyl
945	CONH <sub>2</sub>	2-F-phenyl	4-morpholino
946	CONH <sub>2</sub>	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
947	CONH <sub>2</sub>	2-F-phenyl	4-morpholinocarbonyl
948	CONH <sub>2</sub>	2-F-phenyl	2-methyl-1-imidazolyl
949	CONH <sub>2</sub>	2-F-phenyl	5-methyl-1-imidazolyl
950	CONH <sub>2</sub>	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
951	CONH <sub>2</sub>	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
952	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
953	CONH <sub>2</sub>	2,6-diF-phenyl	1-pyrrolidinocarbonyl
954	CONH <sub>2</sub>	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
955	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholino
956	CONH <sub>2</sub>	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
957	CONH <sub>2</sub>	2,6-diF-phenyl	4-morpholinocarbonyl
958	CONH <sub>2</sub>	2,6-diF-phenyl	2-methyl-1-imidazolyl
959	CONH <sub>2</sub>	2,6-diF-phenyl	5-methyl-1-imidazolyl
960	CONH <sub>2</sub>	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

Table 7



5

For each example, DE is:

10

- (A) pyridin-4-yl-CH<sub>2</sub>,  
 (B) 2-amino-pyrimidin-4-yl,  
 (C) 6-amino-pyridin-2-yl,  
 (D) 3-amidino-4-F-phenyl, or  
 (E) N-amidino-3-piperidinyl.

Ex #	A	B
1	phenyl	2-(aminosulfonyl)phenyl
2	phenyl	2-(methylaminosulfonyl)phenyl
3	phenyl	1-pyrrolidinocarbonyl
4	phenyl	2-(methylsulfonyl)phenyl
5	phenyl	4-morpholino
6	phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
7	phenyl	4-morpholinocarbonyl
8	phenyl	2-methyl-1-imidazolyl
9	phenyl	5-methyl-1-imidazolyl
10	phenyl	2-methylsulfonyl-1-imidazolyl
11	2-pyridyl	2-(aminosulfonyl)phenyl
12	2-pyridyl	2-(methylaminosulfonyl)phenyl
13	2-pyridyl	1-pyrrolidinocarbonyl
14	2-pyridyl	2-(methylsulfonyl)phenyl
15	2-pyridyl	4-morpholino
16	2-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
17	2-pyridyl	4-morpholinocarbonyl
18	2-pyridyl	2-methyl-1-imidazolyl
19	2-pyridyl	5-methyl-1-imidazolyl
20	2-pyridyl	2-methylsulfonyl-1-imidazolyl
21	3-pyridyl	2-(aminosulfonyl)phenyl
22	3-pyridyl	2-(methylaminosulfonyl)phenyl
23	3-pyridyl	1-pyrrolidinocarbonyl
24	3-pyridyl	2-(methylsulfonyl)phenyl
25	3-pyridyl	4-morpholino
26	3-pyridyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
27	3-pyridyl	4-morpholinocarbonyl
28	3-pyridyl	2-methyl-1-imidazolyl
29	3-pyridyl	5-methyl-1-imidazolyl
30	3-pyridyl	2-methylsulfonyl-1-imidazolyl
31	2-pyrimidyl	2-(aminosulfonyl)phenyl
32	2-pyrimidyl	2-(methylaminosulfonyl)phenyl
33	2-pyrimidyl	1-pyrrolidinocarbonyl
34	2-pyrimidyl	2-(methylsulfonyl)phenyl
35	2-pyrimidyl	4-morpholino

36	2-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
37	2-pyrimidyl	4-morpholinocarbonyl
38	2-pyrimidyl	2-methyl-1-imidazolyl
39	2-pyrimidyl	5-methyl-1-imidazolyl
40	2-pyrimidyl	2-methylsulfonyl-1-imidazolyl
41	5-pyrimidyl	2-(aminosulfonyl)phenyl
42	5-pyrimidyl	2-(methylaminosulfonyl)phenyl
43	5-pyrimidyl	1-pyrrolidinocarbonyl
44	5-pyrimidyl	2-(methylsulfonyl)phenyl
45	5-pyrimidyl	4-morpholino
46	5-pyrimidyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
47	5-pyrimidyl	4-morpholinocarbonyl
48	5-pyrimidyl	2-methyl-1-imidazolyl
49	5-pyrimidyl	5-methyl-1-imidazolyl
50	5-pyrimidyl	2-methylsulfonyl-1-imidazolyl
51	2-Cl-phenyl	2-(aminosulfonyl)phenyl
52	2-Cl-phenyl	2-(methylaminosulfonyl)phenyl
53	2-Cl-phenyl	1-pyrrolidinocarbonyl
54	2-Cl-phenyl	2-(methylsulfonyl)phenyl
55	2-Cl-phenyl	4-morpholino
56	2-Cl-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
57	2-Cl-phenyl	4-morpholinocarbonyl
58	2-Cl-phenyl	2-methyl-1-imidazolyl
59	2-Cl-phenyl	5-methyl-1-imidazolyl
60	2-Cl-phenyl	2-methylsulfonyl-1-imidazolyl
61	2-F-phenyl	2-(aminosulfonyl)phenyl
62	2-F-phenyl	2-(methylaminosulfonyl)phenyl
63	2-F-phenyl	1-pyrrolidinocarbonyl
64	2-F-phenyl	2-(methylsulfonyl)phenyl
65	2-F-phenyl	4-morpholino
66	2-F-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
67	2-F-phenyl	4-morpholinocarbonyl
68	2-F-phenyl	2-methyl-1-imidazolyl
69	2-F-phenyl	5-methyl-1-imidazolyl
70	2-F-phenyl	2-methylsulfonyl-1-imidazolyl
71	2,6-diF-phenyl	2-(aminosulfonyl)phenyl
72	2,6-diF-phenyl	2-(methylaminosulfonyl)phenyl
73	2,6-diF-phenyl	1-pyrrolidinocarbonyl
74	2,6-diF-phenyl	2-(methylsulfonyl)phenyl
75	2,6-diF-phenyl	4-morpholino
76	2,6-diF-phenyl	2-(1'-CF <sub>3</sub> -tetrazol-2-yl)phenyl
77	2,6-diF-phenyl	4-morpholinocarbonyl
78	2,6-diF-phenyl	2-methyl-1-imidazolyl
79	2,6-diF-phenyl	5-methyl-1-imidazolyl
80	2,6-diF-phenyl	2-methylsulfonyl-1-imidazolyl

### Utility

The compounds of this invention are useful as anticoagulants for the treatment or prevention of thromboembolic disorders in mammals. The term "thromboembolic disorders" as used herein includes arterial or venous cardiovascular or cerebrovascular thromboembolic disorders, including, for example, unstable angina, first or recurrent myocardial infarction, ischemic sudden death, transient ischemic attack, stroke, atherosclerosis, venous thrombosis, deep vein thrombosis, thrombophlebitis, arterial embolism, coronary and cerebral arterial thrombosis, cerebral embolism, kidney embolisms, and pulmonary embolisms. The anticoagulant effect of compounds of the present invention is believed to be due to inhibition of factor Xa or thrombin.

The effectiveness of compounds of the present invention as inhibitors of factor Xa was determined using purified human factor Xa and synthetic substrate. The rate of factor Xa hydrolysis of chromogenic substrate S2222 (Kabi Pharmacia, Franklin, OH) was measured both in the absence and presence of compounds of the present invention. Hydrolysis of the substrate resulted in the release of pNA, which was monitored spectrophotometrically by measuring the increase in absorbance at 405 nm. A decrease in the rate of absorbance change at 405 nm in the presence of inhibitor is indicative of enzyme inhibition. The results of this assay are expressed as inhibitory constant,  $K_i$ .

Factor Xa determinations were made in 0.10 M sodium phosphate buffer, pH 7.5, containing 0.20 M NaCl, and 0.5 % PEG 8000. The Michaelis constant,  $K_m$ , for substrate hydrolysis was determined at 25°C using the method of Lineweaver and Burk. Values of  $K_i$  were determined by allowing 0.2-0.5 nM human factor Xa (Enzyme Research Laboratories, South Bend, IN) to react with the substrate (0.20 mM-1 mM) in the presence of inhibitor. Reactions were allowed to go for 30 minutes and the velocities (rate of absorbance change vs time) were measured in the time frame of 25-30 minutes. The following relationship was used to calculate  $K_i$  values:

$$(v_o - v_s) / v_s = I / (K_i (1 + S / K_m))$$



where:

$v_0$  is the velocity of the control in the absence of inhibitor;

$v_s$  is the velocity in the presence of inhibitor;

5  $I$  is the concentration of inhibitor;

$K_i$  is the dissociation constant of the enzyme:inhibitor complex;

$S$  is the concentration of substrate;

$K_m$  is the Michaelis constant.

10 Using the methodology described above, a number of compounds of the present invention were found to exhibit a  $K_i$  of  $\leq 10 \mu M$ , thereby confirming the utility of the compounds of the present invention as effective Xa inhibitors.

The antithrombotic effect of compounds of the present invention can be demonstrated in a rabbit arterio-venous (AV) shunt thrombosis model. In this model, rabbits weighing 2-3 kg anesthetized with a mixture of xylazine (10 mg/kg i.m.) and ketamine (50 mg/kg i.m.) are used. A saline-filled AV shunt device is connected between the femoral arterial and the femoral venous cannulae. The AV shunt device consists of a piece of 6-cm tygon tubing which contains a piece of silk thread. Blood will flow from the femoral artery via the AV-shunt into the femoral vein. The exposure of flowing blood to a silk thread will induce the formation of a significant thrombus. After forty minutes, the shunt is disconnected and the silk thread covered with thrombus is weighed. Test agents or vehicle will be given (i.v., i.p., s.c., or orally) prior to the opening of the AV shunt. The percentage inhibition of thrombus formation is determined for each treatment group.

20 25 30 The ID50 values (dose which produces 50% inhibition of thrombus formation) are estimated by linear regression.

The compounds of formula (I) may also be useful as inhibitors of serine proteases, notably human thrombin, plasma kallikrein and plasmin. Because of their inhibitory action, these compounds are indicated for use in the prevention or treatment of physiological reactions, blood coagulation and inflammation, catalyzed by the aforesaid class of enzymes. Specifically, the compounds have utility as drugs for the

35

treatment of diseases arising from elevated thrombin activity such as myocardial infarction, and as reagents used as anticoagulants in the processing of blood to plasma for diagnostic and other commercial purposes.

5       Some compounds of the present invention were shown to be direct acting inhibitors of the serine protease thrombin by their ability to inhibit the cleavage of small molecule substrates by thrombin in a purified system. *In vitro* inhibition constants were determined by the method described  
10 by Kettner et al. in *J. Biol. Chem.* **265**, 18289-18297 (1990), herein incorporated by reference. In these assays, thrombin-mediated hydrolysis of the chromogenic substrate S2238 (Helena Laboratories, Beaumont, TX) was monitored spectrophotometrically. Addition of an inhibitor to the assay  
15 mixture results in decreased absorbance and is indicative of thrombin inhibition. Human thrombin (Enzyme Research Laboratories, Inc., South Bend, IN) at a concentration of 0.2 nM in 0.10 M sodium phosphate buffer, pH 7.5, 0.20 M NaCl, and 0.5% PEG 6000, was incubated with various substrate  
20 concentrations ranging from 0.20 to 0.02 mM. After 25 to 30 minutes of incubation, thrombin activity was assayed by monitoring the rate of increase in absorbance at 405 nm which arises owing to substrate hydrolysis. Inhibition constants were derived from reciprocal plots of the reaction velocity as  
25 a function of substrate concentration using the standard method of Lineweaver and Burk. Using the methodology described above, some compounds of this invention were evaluated and found to exhibit a  $K_i$  of less than 10  $\mu\text{M}$ , thereby confirming the utility of the compounds of the present  
30 invention as effective thrombin inhibitors.

      The compounds of the present invention can be administered alone or in combination with one or more additional therapeutic agents. These include other anti-coagulant or coagulation inhibitory agents, anti-platelet or  
35 platelet inhibitory agents, thrombin inhibitors, or thrombolytic or fibrinolytic agents.

      The compounds are administered to a mammal in a therapeutically effective amount. By "therapeutically

effective amount" it is meant an amount of a compound of Formula I that, when administered alone or in combination with an additional therapeutic agent to a mammal, is effective to prevent or ameliorate the thromboembolic disease condition or the progression of the disease.

By "administered in combination" or "combination therapy" it is meant that the compound of Formula I and one or more additional therapeutic agents are administered concurrently to the mammal being treated. When administered in combination each component may be administered at the same time or sequentially in any order at different points in time. Thus, each component may be administered separately but sufficiently closely in time so as to provide the desired therapeutic effect. Other anticoagulant agents (or coagulation inhibitory agents) that may be used in combination with the compounds of this invention include warfarin and heparin, as well as other factor Xa inhibitors such as those described in the publications identified above under Background of the Invention.

The term anti-platelet agents (or platelet inhibitory agents), as used herein, denotes agents that inhibit platelet function such as by inhibiting the aggregation, adhesion or granular secretion of platelets. Such agents include, but are not limited to, the various known non-steroidal anti-inflammatory drugs (NSAIDS) such as aspirin, ibuprofen, naproxen, sulindac, indomethacin, mefenamate, droxicam, diclofenac, sulfinpyrazone, and piroxicam, including pharmaceutically acceptable salts or prodrugs thereof. Of the NSAIDS, aspirin (acetylsalicyclic acid or ASA), and piroxicam are preferred. Other suitable anti-platelet agents include ticlopidine, including pharmaceutically acceptable salts or prodrugs thereof. Ticlopidine is also a preferred compound since it is known to be gentle on the gastro-intestinal tract in use. Still other suitable platelet inhibitory agents include IIb/IIIa antagonists, thromboxane-A<sub>2</sub>-receptor antagonists and thromboxane-A<sub>2</sub>-synthetase inhibitors, as well as pharmaceutically acceptable salts or prodrugs thereof.

The term thrombin inhibitors (or anti-thrombin agents), as used herein, denotes inhibitors of the serine protease thrombin. By inhibiting thrombin, various thrombin-mediated processes, such as thrombin-mediated platelet activation (that is, for example, the aggregation of platelets, and/or the granular secretion of plasminogen activator inhibitor-1 and/or serotonin) and/or fibrin formation are disrupted. A number of thrombin inhibitors are known to one of skill in the art and these inhibitors are contemplated to be used in combination with the present compounds. Such inhibitors include, but are not limited to, boroarginine derivatives, boroptides, heparins, hirudin and argatroban, including pharmaceutically acceptable salts and prodrugs thereof. Boroarginine derivatives and boroptides include N-acetyl and peptide derivatives of boronic acid, such as C-terminal  $\alpha$ -aminoboronic acid derivatives of lysine, ornithine, arginine, homoarginine and corresponding isothiuronium analogs thereof. The term hirudin, as used herein, includes suitable derivatives or analogs of hirudin, referred to herein as hirulogs, such as disulfatohirudin. Boroptide thrombin inhibitors include compounds described in Kettner et al., U.S. Patent No. 5,187,157 and European Patent Application Publication Number 293 881 A2, the disclosures of which are hereby incorporated herein by reference. Other suitable boroarginine derivatives and boroptide thrombin inhibitors include those disclosed in PCT Application Publication Number 92/07869 and European Patent Application Publication Number 471,651 A2, the disclosures of which are hereby incorporated herein by reference.

The term thrombolytics (or fibrinolytic) agents (or thrombolytics or fibrinolytics), as used herein, denotes agents that lyse blood clots (thrombi). Such agents include tissue plasminogen activator, anistreplase, urokinase or streptokinase, including pharmaceutically acceptable salts or prodrugs thereof. The term anistreplase, as used herein, refers to anisoylated plasminogen streptokinase activator complex, as described, for example, in European Patent Application No. 028,489, the disclosure of which is hereby

incorporated herein by reference herein. The term urokinase, as used herein, is intended to denote both dual and single chain urokinase, the latter also being referred to herein as prourokinase.

5 Administration of the compounds of Formula I of the invention in combination with such additional therapeutic agent, may afford an efficacy advantage over the compounds and agents alone, and may do so while permitting the use of lower doses of each. A lower dosage minimizes the potential of side effects, thereby providing an increased margin of safety.

10 The compounds of the present invention are also useful as standard or reference compounds, for example as a quality standard or control, in tests or assays involving the inhibition of factor Xa. Such compounds may be provided in a commercial kit, for example, for use in pharmaceutical research involving factor Xa. For example, a compound of the present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference compound. When developing new assays or protocols, compounds according to the present invention could be used to test their effectiveness.

20 The compounds of the present invention may also be used in diagnostic assays involving factor Xa. For example, the presence of factor Xa in an unknown sample could be determined by addition of chromogenic substrate S2222 to a series of solutions containing test sample and optionally one of the compounds of the present invention. If production of pNA is observed in the solutions containing test sample, but not in the presence of a compound of the present invention, then one would conclude factor Xa was present.

35

#### Dosage and Formulation

The compounds of this invention can be administered in such oral dosage forms as tablets, capsules (each of which includes sustained release or timed release formulations),

pills, powders, granules, elixirs, tinctures, suspensions, syrups, and emulsions. They may also be administered in intravenous (bolus or infusion), intraperitoneal, subcutaneous, or intramuscular form, all using dosage forms well known to those of ordinary skill in the pharmaceutical arts. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired. A physician or veterinarian can determine and prescribe the effective amount of the drug required to prevent, counter, or arrest the progress of the thromboembolic disorder.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.001 to 1000 mg/kg of body weight, preferably between about 0.01 to 100 mg/kg of body weight per day, and most preferably between about 1.0 to 20 mg/kg/day. Intravenously, the most preferred doses will range from about 1 to about 10 mg/kg/minute during a constant rate infusion. Compounds of this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

Compounds of this invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using transdermal skin patches. When administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

The compounds are typically administered in admixture with suitable pharmaceutical diluents, excipients, or carriers (collectively referred to herein as pharmaceutical carriers) suitably selected with respect to the intended form of administration, that is, oral tablets, capsules, elixirs, syrups and the like, and consistent with conventional pharmaceutical practices.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic, pharmaceutically acceptable, inert carrier such as lactose, starch, sucrose, glucose, methyl cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, mannitol, sorbitol and the like; for oral administration in liquid form, the oral drug components can be combined with any oral, non-toxic, pharmaceutically acceptable inert carrier such as ethanol, glycerol, water, and the like. Moreover, when desired or necessary, suitable binders, lubricants, disintegrating agents, and coloring agents can also be incorporated into the mixture. Suitable binders include starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth, or sodium alginate, carboxymethylcellulose, polyethylene glycol, waxes, and the like. Lubricants used in these dosage forms include sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum, and the like.

The compounds of the present invention can also be administered in the form of liposome delivery systems, such as small unilamellar vesicles, large unilamellar vesicles, and multilamellar vesicles. Liposomes can be formed from a variety of phospholipids, such as cholesterol, stearylamine, or phosphatidylcholines.

Compounds of the present invention may also be coupled with soluble polymers as targetable drug carriers. Such polymers can include polyvinylpyrrolidone, pyran copolymer, polyhydroxypropylmethacrylamide-phenol,

polyhydroxyethylaspartamidephenol, or polyethyleneoxide-polylysine substituted with palmitoyl residues. Furthermore, the compounds of the present invention may be coupled to a class of biodegradable polymers useful in achieving controlled  
5 release of a drug, for example, polylactic acid, polyglycolic acid, copolymers of polylactic and polyglycolic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters, polyacetals, polydihdropyrans, polycyanoacylates, and crosslinked or amphipathic block  
10 copolymers of hydrogels.

Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 100 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will  
15 ordinarily be present in an amount of about 0.5-95% by weight based on the total weight of the composition.

Gelatin capsules may contain the active ingredient and powdered carriers, such as lactose, starch, cellulose derivatives, magnesium stearate, stearic acid, and the like.  
20 Similar diluents can be used to make compressed tablets. Both tablets and capsules can be manufactured as sustained release products to provide for continuous release of medication over a period of hours. Compressed tablets can be sugar coated or film coated to mask any unpleasant taste and protect the  
25 tablet from the atmosphere, or enteric coated for selective disintegration in the gastrointestinal tract.

Liquid dosage forms for oral administration can contain coloring and flavoring to increase patient acceptance.

In general, water, a suitable oil, saline, aqueous  
30 dextrose (glucose), and related sugar solutions and glycols such as propylene glycol or polyethylene glycols are suitable carriers for parenteral solutions. Solutions for parenteral administration preferably contain a water soluble salt of the active ingredient, suitable stabilizing agents, and if  
35 necessary, buffer substances. Antioxidizing agents such as sodium bisulfite, sodium sulfite, or ascorbic acid, either alone or combined, are suitable stabilizing agents. Also used are citric acid and its salts and sodium EDTA. In addition,



parenteral solutions can contain preservatives, such as benzalkonium chloride, methyl- or propyl-paraben, and chlorobutanol.

Suitable pharmaceutical carriers are described in  
5 Remington's Pharmaceutical Sciences, Mack Publishing Company,  
a standard reference text in this field.

Representative useful pharmaceutical dosage-forms for  
administration of the compounds of this invention can be  
illustrated as follows:

10       Capsules

A large number of unit capsules can be prepared by  
filling standard two-piece hard gelatin capsules each with 100  
milligrams of powdered active ingredient, 150 milligrams of  
lactose, 50 milligrams of cellulose, and 6 milligrams  
15       magnesium stearate.

Soft Gelatin Capsules

A mixture of active ingredient in a digestable oil such  
as soybean oil, cottonseed oil or olive oil may be prepared  
and injected by means of a positive displacement pump into  
20       gelatin to form soft gelatin capsules containing 100  
milligrams of the active ingredient. The capsules should be  
washed and dried.

Tablets

Tablets may be prepared by conventional procedures so  
25       that the dosage unit is 100 milligrams of active ingredient,  
0.2 milligrams of colloidal silicon dioxide, 5 milligrams of  
magnesium stearate, 275 milligrams of microcrystalline  
cellulose, 11 milligrams of starch and 98.8 milligrams of  
lactose. Appropriate coatings may be applied to increase  
30       palatability or delay absorption.

Injectable

A parenteral composition suitable for administration by  
injection may be prepared by stirring 1.5% by weight of active  
ingredient in 10% by volume propylene glycol and water. The  
35       solution should be made isotonic with sodium chloride and  
sterilized.

### Suspension

An aqueous suspension can be prepared for oral administration so that each 5 mL contain 100 mg of finely divided active ingredient, 200 mg of sodium carboxymethyl cellulose, 5 mg of sodium benzoate, 1.0 g of sorbitol solution, U.S.P., and 0.025 mL of vanillin.

Where the compounds of this invention are combined with other anticoagulant agents, for example, a daily dosage may be about 0.1 to 100 milligrams of the compound of Formula I and about 1 to 7.5 milligrams of the second anticoagulant, per kilogram of patient body weight. For a tablet dosage form, the compounds of this invention generally may be present in an amount of about 5 to 10 milligrams per dosage unit, and the second anti-coagulant in an amount of about 1 to 5 milligrams per dosage unit.

Where the compounds of Formula I are administered in combination with an anti-platelet agent, by way of general guidance, typically a daily dosage may be about 0.01 to 25 milligrams of the compound of Formula I and about 50 to 150 milligrams of the anti-platelet agent, preferably about 0.1 to 1 milligrams of the compound of Formula I and about 1 to 3 milligrams of antiplatelet agents, per kilogram of patient body weight.

Where the compounds of Formula I are administered in combination with thrombolytic agent, typically a daily dosage may be about 0.1 to 1 milligrams of the compound of Formula I, per kilogram of patient body weight and, in the case of the thrombolytic agents, the usual dosage of the thrombolytic agent when administered alone may be reduced by about 70-80% when administered with a compound of Formula I.

Where two or more of the foregoing second therapeutic agents are administered with the compound of Formula I, generally the amount of each component in a typical daily dosage and typical dosage form may be reduced relative to the usual dosage of the agent when administered alone, in view of the additive or synergistic effect of the therapeutic agents when administered in combination.

Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of Formula I and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a material which effects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a lowviscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.


These as well as other ways of minimizing contact between the components of combination products of the present invention, whether administered in a single dosage form or administered in separate forms but at the same time by the same manner, will be readily apparent to those skilled in the art, once armed with the present disclosure.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the

scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

WHAT IS CLAIMED AS NEW AND DESIRED TO BE SECURED BY  
LETTER PATENT OF UNITED STATES IS:

1. A compound of formula I:



I

or a stereoisomer or pharmaceutically acceptable salt thereof,  
wherein;

ring M contains, in addition to J, 0-3 N atoms, provided that  
if M contains 2 N atoms then R<sup>1b</sup> is not present and if M  
contains 3 N atoms then R<sup>1a</sup> and R<sup>1b</sup> are not present;

J is N or NH;

D is selected from CN, C(=NR<sup>8</sup>)NR<sup>7</sup>R<sup>9</sup>, NHC(=NR<sup>8</sup>)NR<sup>7</sup>R<sup>9</sup>,  
NR<sup>8</sup>CH(=NR<sup>7</sup>), C(O)NR<sup>7</sup>R<sup>8</sup>, and (CR<sup>8</sup>R<sup>9</sup>)<sub>t</sub>NR<sup>7</sup>R<sup>8</sup>, provided that D  
is substituted meta or para to G on E;

E is selected from phenyl, pyridyl, pyrimidyl, pyrazinyl,  
pyridazinyl, and piperidinyl substituted with 1 R;

alternatively, D-E-G together represent pyridyl substituted  
with 1 R;

R is selected from H, halogen, (CH<sub>2</sub>)<sub>t</sub>OR<sup>3</sup>, C<sub>1-4</sub> alkyl, OCF<sub>3</sub>, and  
CF<sub>3</sub>;

G is absent or is selected from NHCH<sub>2</sub>, OCH<sub>2</sub>, and SCH<sub>2</sub>, provided  
that when s is 0, then G is attached to a carbon atom on  
ring M;

Z is selected from a C<sub>1-4</sub> alkylene, (CH<sub>2</sub>)<sub>r</sub>O(CH<sub>2</sub>)<sub>r</sub>,  
(CH<sub>2</sub>)<sub>r</sub>NR<sup>3</sup>(CH<sub>2</sub>)<sub>r</sub>, (CH<sub>2</sub>)<sub>r</sub>C(O)(CH<sub>2</sub>)<sub>r</sub>, (CH<sub>2</sub>)<sub>r</sub>C(O)O(CH<sub>2</sub>)<sub>r</sub>,

- $(\text{CH}_2)_r\text{OC}(\text{O})(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  
 $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{OC}(\text{O})\text{O}(\text{CH}_2)_r$ ,  
 $(\text{CH}_2)_r\text{OC}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})\text{O}(\text{CH}_2)_r$ ,  
 $(\text{CH}_2)_r\text{NR}^3\text{C}(\text{O})\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{S}(\text{O})_p(\text{CH}_2)_r$ ,  
5  $(\text{CH}_2)_r\text{SO}_2\text{NR}^3(\text{CH}_2)_r$ ,  $(\text{CH}_2)_r\text{NR}^3\text{SO}_2(\text{CH}_2)_r$ , and  
 $(\text{CH}_2)_r\text{NR}^3\text{SO}_2\text{NR}^3(\text{CH}_2)_r$ , provided that Z does not form a N-  
N, N-O, N-S, NCH<sub>2</sub>N, NCH<sub>2</sub>O, or NCH<sub>2</sub>S bond with ring M or  
group A;
- 10  $\text{R}^{1a}$  and  $\text{R}^{1b}$  are independently absent or selected from  
 $-(\text{CH}_2)_r\text{R}^{1'}$ ,  $\text{NCH}_2\text{R}^{1''}$ ,  $\text{OCH}_2\text{R}^{1''}$ ,  $\text{SCH}_2\text{R}^{1''}$ ,  $\text{N}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ ,  
 $\text{O}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ , and  $\text{S}(\text{CH}_2)_2(\text{CH}_2)_t\text{R}^{1'}$ , or combined to form  
a 5-8 membered saturated, partially saturated or  
unsaturated ring substituted with 0-2  $\text{R}^4$  and which  
15 contains from 0-2 heteroatoms selected from the group  
consisting of N, O, and S;
- $\text{R}^{1'}$  is selected from H, C<sub>1-3</sub> alkyl, halo,  $(\text{CF}_2)_r\text{CF}_3$ ,  $\text{OR}^2$ ,  
 $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(\text{O})\text{R}^{2c}$ ,  $\text{OC}(\text{O})\text{R}^2$ ,  $(\text{CF}_2)_r\text{CO}_2\text{R}^{2c}$ ,  $\text{S}(\text{O})_p\text{R}^{2b}$ ,  
20  $\text{NR}^2(\text{CH}_2)_r\text{OR}^2$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})\text{NHR}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2a}$ ,  
 $\text{OC}(\text{O})\text{NR}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ,  $\text{NR}^2\text{SO}_2\text{R}^{2b}$ , C<sub>3-6</sub>  
carbocyclic residue substituted with 0-2  $\text{R}^4$ , and 5-10  
membered heterocyclic system containing from 1-4  
heteroatoms selected from the group consisting of N, O,  
25 and S substituted with 0-2  $\text{R}^4$ ;
- $\text{R}^{1''}$  is selected from H,  $\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  $\text{S}(\text{O})\text{R}^{2b}$ ,  $\text{S}(\text{O})_2\text{R}^{2b}$ ,  
and  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ;
- 30  $\text{R}^2$ , at each occurrence, is selected from H,  $\text{CF}_3$ , C<sub>1-6</sub> alkyl,  
benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2  $\text{R}^{4b}$ ,  
and 5-6 membered heterocyclic system containing from 1-4  
heteroatoms selected from the group consisting of N, O,  
and S substituted with 0-2  $\text{R}^{4b}$ ;
- 35  $\text{R}^{2a}$ , at each occurrence, is selected from H,  $\text{CF}_3$ , C<sub>1-6</sub> alkyl,  
benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2  $\text{R}^{4b}$ ,  
and 5-6 membered heterocyclic system containing from 1-4

heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

5 R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkyl, benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2 R<sup>4b</sup>, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

10 R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, C<sub>1-4</sub> alkoxy, C<sub>1-6</sub> alkyl, benzyl, C<sub>3-6</sub> carbocyclic residue substituted with 0-2 R<sup>4b</sup>, and 5-6 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4b</sup>;

15 alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a 5 or 6 membered saturated, partially saturated or unsaturated ring substituted with 0-2 R<sup>4b</sup> which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

20

R<sup>3</sup>, at each occurrence, is selected from H, C<sub>1-4</sub> alkyl, and phenyl;

25 R<sup>3a</sup>, at each occurrence, is selected from H, C<sub>1-4</sub> alkyl, and phenyl;

A is selected from:

30 C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4</sup>, and 5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4</sup>;

B is selected from:

35 X-Y, NR<sup>2</sup>R<sup>2a</sup>, C(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>C(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4a</sup>, and

5-10 membered heterocyclic system containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4a</sup>;

- 5 X is selected from C<sub>1-4</sub> alkylene, -CR<sup>2</sup>(CR<sup>2</sup>R<sup>2b</sup>)(CH<sub>2</sub>)<sub>t</sub>-, -C(O)-, -C(=NR)-, -CR<sup>2</sup>(NR<sup>1</sup>\*R<sup>2</sup>)-, -CR<sup>2</sup>(OR<sup>2</sup>)-, -CR<sup>2</sup>(SR<sup>2</sup>)-, -C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)-, -S(O)<sub>p</sub>-, -S(O)<sub>p</sub>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>S(O)<sub>p</sub>-, -S(O)<sub>2</sub>NR<sup>2</sup>-, -NR<sup>2</sup>S(O)<sub>2</sub>-, -NR<sup>2</sup>S(O)<sub>2</sub>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>S(O)<sub>2</sub>NR<sup>2</sup>-, -NR<sup>2</sup>S(O)<sub>2</sub>NR<sup>2</sup>-, -C(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)-, -C(O)NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -NR<sup>2</sup>C(O)CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>C(O)NR<sup>2</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>C(O)-, -NR<sup>2</sup>C(O)O-, -OC(O)NR<sup>2</sup>-, -NR<sup>2</sup>C(O)NR<sup>2</sup>-, -NR<sup>2</sup>-, -NR<sup>2</sup>CR<sup>2</sup>R<sup>2a</sup>-, -CR<sup>2</sup>R<sup>2a</sup>NR<sup>2</sup>-, O, -CR<sup>2</sup>R<sup>2a</sup>O-, and -OCR<sup>2</sup>R<sup>2a</sup>-;
- 10
- 15 Y is selected from:  
 (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a N-N, O-N, or S-N bond,  
 C<sub>3-10</sub> carbocyclic residue substituted with 0-2 R<sup>4a</sup>, and  
 5-10 membered heterocyclic system containing from 1-4  
 20 heteroatoms selected from the group consisting of N, O, and S substituted with 0-2 R<sup>4a</sup>;
- R<sup>4</sup>, at each occurrence, is selected from =O, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, -CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>2b</sup>,  
 25 NR<sup>2</sup>C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>C(O)NR<sup>2</sup>R<sup>2a</sup>, CH(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, NHC(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>-C<sub>1-4</sub> alkyl, NR<sup>2</sup>SO<sub>2</sub>R<sup>5</sup>, S(O)<sub>p</sub>R<sup>5</sup>, (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>, NCH<sub>2</sub>R<sup>1</sup>\*, OCH<sub>2</sub>R<sup>1</sup>\*, SCH<sub>2</sub>R<sup>1</sup>\*, N(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>\*, O(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>\*, and S(CH<sub>2</sub>)<sub>2</sub>(CH<sub>2</sub>)<sub>t</sub>R<sup>1</sup>\*,
- 30 alternatively, one R<sup>4</sup> is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S;
- R<sup>4a</sup>, at each occurrence, is selected from =O, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, -CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>2b</sup>,  
 35 NR<sup>2</sup>C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>C(O)NR<sup>2</sup>R<sup>2a</sup>, CH(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, NHC(=NR<sup>2</sup>)NR<sup>2</sup>R<sup>2a</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>-C<sub>1-4</sub> alkyl, NR<sup>2</sup>SO<sub>2</sub>R<sup>5</sup>, S(O)<sub>p</sub>R<sup>5</sup>, and (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>;



alternatively, one R<sup>4a</sup> is a 5-6 membered aromatic heterocycle containing from 1-4 heteroatoms selected from the group consisting of N, O, and S substituted with 0-1 R<sup>5</sup>;

5 R<sup>4b</sup>, at each occurrence, is selected from =O, (CH<sub>2</sub>)<sub>r</sub>OR<sup>3</sup>, halo, C<sub>1-4</sub> alkyl, -CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>3</sup>R<sup>3a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>3</sup>, NR<sup>3</sup>C(O)R<sup>3a</sup>, C(O)NR<sup>3</sup>R<sup>3a</sup>, NR<sup>3</sup>C(O)NR<sup>3</sup>R<sup>3a</sup>, CH(=NR<sup>3</sup>)NR<sup>3</sup>R<sup>3a</sup>, NH<sup>3</sup>C(=NR<sup>3</sup>)NR<sup>3</sup>R<sup>3a</sup>, SO<sub>2</sub>NR<sup>3</sup>R<sup>3a</sup>, NR<sup>3</sup>SO<sub>2</sub>NR<sup>3</sup>R<sup>3a</sup>, NR<sup>3</sup>SO<sub>2</sub>-C<sub>1-4</sub> alkyl, NR<sup>3</sup>SO<sub>2</sub>CF<sub>3</sub>, NR<sup>3</sup>SO<sub>2</sub>-phenyl, S(O)<sub>p</sub>CF<sub>3</sub>, S(O)<sub>p</sub>-C<sub>1-4</sub> alkyl, S(O)<sub>p</sub>-phenyl, and (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>;

15 R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl, phenyl substituted with 0-2 R<sup>6</sup>, and benzyl substituted with 0-2 R<sup>6</sup>;

20 R<sup>6</sup>, at each occurrence, is selected from H, OH, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, CN, NO<sub>2</sub>, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, (CH<sub>2</sub>)<sub>r</sub>C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)NR<sup>2</sup>R<sup>2a</sup>, CH(=NH)NH<sub>2</sub>, NHC(=NH)NH<sub>2</sub>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, NR<sup>2</sup>SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and NR<sup>2</sup>SO<sub>2</sub>C<sub>1-4</sub> alkyl;

25 R<sup>7</sup>, at each occurrence, is selected from H, OH, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkylcarbonyl, C<sub>1-6</sub> alkoxy, C<sub>1-4</sub> alkoxycarbonyl, (CH<sub>2</sub>)<sub>n</sub>-phenyl, C<sub>6-10</sub> aryloxy, C<sub>6-10</sub> aryloxy carbonyl, C<sub>6-10</sub> arylmethylcarbonyl, C<sub>1-4</sub> alkylcarbonyloxy C<sub>1-4</sub> alkoxycarbonyl, C<sub>6-10</sub> arylcarbonyloxy C<sub>1-4</sub> alkoxycarbonyl, C<sub>1-6</sub> alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C<sub>1-4</sub> alkoxycarbonyl;

30 R<sup>8</sup>, at each occurrence, is selected from H, C<sub>1-6</sub> alkyl and (CH<sub>2</sub>)<sub>n</sub>-phenyl;

35 alternatively, R<sup>7</sup> and R<sup>8</sup> combine to form a 5 or 6 membered saturated, ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

$R^9$ , at each occurrence, is selected from H,  $C_{1-6}$  alkyl and  $(CH_2)_n$ -phenyl;

n, at each occurrence, is selected from 0, 1, 2, and 3;

m, at each occurrence, is selected from 0, 1, and 2;

p, at each occurrence, is selected from 0, 1, and 2;


r, at each occurrence, is selected from 0, 1, 2, and 3;

s, at each occurrence, is selected from 0, 1, and 2; and,

t, at each occurrence, is selected from 0 and 1;

provided that D-E-G- $(CH_2)_s$ - and -Z-A-B are not both benzamidines.

2. A compound according to Claim 1, wherein the compound is of formulae Ia-Ih:



wherein, groups D-E- and -Z-A-B are attached to adjacent atoms on the ring;

Z is selected from a  $\text{CH}_2\text{O}$ ,  $\text{OCH}_2$ ,  $\text{CH}_2\text{NH}$ ,  $\text{NHCH}_2$ ,  $\text{C}(\text{O})$ ,  $\text{CH}_2\text{C}(\text{O})$ ,  $\text{C}(\text{O})\text{CH}_2$ ,  $\text{NHC}(\text{O})$ ,  $\text{C}(\text{O})\text{NH}$ ,  $\text{CH}_2\text{S}(\text{O})_2$ ,  $\text{S}(\text{O})_2(\text{CH}_2)$ ,  $\text{SO}_2\text{NH}$ , and  $\text{NHSO}_2$ , provided that Z does not form a N-N, N-O,  $\text{NCH}_2\text{N}$ , or  $\text{NCH}_2\text{O}$  bond with ring M or group A;

5  
A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2  $\text{R}^4$ ;  
phenyl, piperidinyl, piperazinyl, pyridyl,  
pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,  
10 pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,  
isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,  
thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,  
1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,  
1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,  
15 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,  
1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl,  
benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl,  
benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl,  
benziso-thiazolyl, and isoindazolyl;

20  
B is selected from: Y, X-Y,  $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ , and  $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ ;

25  
X is selected from  $\text{C}_{1-4}$  alkylene,  $-\text{C}(\text{O})-$ ,  $-\text{C}(=\text{NR})-$ ,  
 $-\text{CR}^2(\text{NR}^2\text{R}^{2a})-$ ,  $-\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$ ,  $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})-$ ,  $-\text{C}(\text{O})\text{NR}^2-$ ,  
 $-\text{NR}^2\text{C}(\text{O})-$ ,  $-\text{C}(\text{O})\text{NR}^2\text{CR}^2\text{R}^{2a}-$ ,  $-\text{NR}^2\text{C}(\text{O})\text{CR}^2\text{R}^{2a}-$ ,  
 $-\text{CR}^2\text{R}^{2a}\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{CR}^2\text{R}^{2a}\text{NR}^2\text{C}(\text{O})-$ ,  $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{NR}^2-$ ,  
 $-\text{NR}^2\text{CR}^2\text{R}^{2a}-$ ,  $-\text{CR}^2\text{R}^{2a}\text{NR}^2-$ , O,  $-\text{CR}^2\text{R}^{2a}\text{O}-$ , and  $-\text{OCR}^2\text{R}^{2a}-$ ;


30 Y is  $\text{NR}^2\text{R}^{2a}$ , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2  $\text{R}^{4a}$ ;  
35 cyclopropyl, cyclopentyl, cyclohexyl, phenyl,  
piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl,  
morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl,  
oxazolyl, isoxazolyl, isoxazolinyl, thiazolyl,

isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl,  
 thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,  
 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,  
 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,  
 5 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,  
 1,2,4-triazolyl, 1,2,5-triazolyl, 1,3,4-triazolyl,  
 benzofuranyl, benzothiofuranyl, indolyl, benzimidazolyl,  
 benzoxazolyl, benzthiazolyl, indazolyl, benzisoxazolyl,  
 benzisothiazolyl, and isoindazolyl;

10


alternatively, Y is selected from the following bicyclic  
 heteroaryl ring systems:



15 K is selected from O, S, NH, and N.

3. A compound according to Claim 2, the compound is of  
 formulae IIa-IIf:

20



wherein;

5 Z is selected from a C(O), CH<sub>2</sub>C(O), C(O)CH<sub>2</sub>, NHC(O), C(O)NH, C(O)N(CH<sub>3</sub>), CH<sub>2</sub>S(O)<sub>2</sub>, S(O)<sub>2</sub>(CH<sub>2</sub>), SO<sub>2</sub>NH, and NHSO<sub>2</sub>, provided that Z does not form a N-N or NCH<sub>2</sub>N bond with ring M or group A.

4. A compound according to Claim 3, wherein;

10 E is phenyl substituted with R or 2-pyridyl substituted with R;

D is selected from NH<sub>2</sub>, C(O)NH<sub>2</sub>, C(=NH)NH<sub>2</sub>, CH<sub>2</sub>NH<sub>2</sub>, CH<sub>2</sub>NHCH<sub>3</sub>, CH(CH<sub>3</sub>)NH<sub>2</sub>, and C(CH<sub>3</sub>)<sub>2</sub>NH<sub>2</sub>, provided that D is substituted  
15 meta or para to ring M on E; and,

R is selected from H, OCH<sub>3</sub>, Cl, and F.

20 5. A compound according to Claim 4, wherein;

D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, and 6-(2-amino-2-propyl)pyrid-2-yl.  
25  
30  
35

6. A compound according to Claim 3, wherein;

Z is C(O)CH<sub>2</sub> and CONH, provided that Z does not form a N-N bond with group A;

5 A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2 R<sup>4</sup>; and,

B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1 R<sup>4a</sup>;

10 R<sup>4</sup>, at each occurrence, is selected from OH, (CH<sub>2</sub>)<sub>r</sub>OR<sup>2</sup>, halo, C<sub>1-4</sub> alkyl, (CH<sub>2</sub>)<sub>r</sub>NR<sup>2</sup>R<sup>2a</sup>, and (CF<sub>2</sub>)<sub>r</sub>CF<sub>3</sub>;

15 R<sup>4a</sup> is selected from C<sub>1-4</sub> alkyl, CF<sub>3</sub>, S(O)<sub>p</sub>R<sup>5</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and 1-CF<sub>3</sub>-tetrazol-2-yl;

R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl, phenyl, and benzyl;

20 X is CH<sub>2</sub> or C(O); and,

Y is selected from pyrrolidino and morpholino.

25 7. A compound according to Claim 6, wherein;

A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,

30 B is selected from the group: 2-CF<sub>3</sub>-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF<sub>3</sub>-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

8. A compound according to Claim 3, wherein;
- 5 E is phenyl substituted with R or 2-pyridyl substituted with R;
- D is selected from  $\text{NH}_2$ ,  $\text{C}(\text{O})\text{NH}_2$ ,  $\text{C}(=\text{NH})\text{NH}_2$ ,  $\text{CH}_2\text{NH}_2$ ,  $\text{CH}_2\text{NHCH}_3$ ,  $\text{CH}(\text{CH}_3)\text{NH}_2$ , and  $\text{C}(\text{CH}_3)_2\text{NH}_2$ , provided that D is substituted  
10 meta or para to ring M on E; and,
- R is selected from H,  $\text{OCH}_3$ , Cl, and F;
- Z is  $\text{C}(\text{O})\text{CH}_2$  and  $\text{CONH}$ , provided that Z does not form a N-N bond  
15 with group A;
- A is selected from phenyl, pyridyl, and pyrimidyl, and is substituted with 0-2  $\text{R}^4$ ; and,
- 20 B is selected from X-Y, phenyl, pyrrolidino, morpholino, 1,2,3-triazolyl, and imidazolyl, and is substituted with 0-1  $\text{R}^{4a}$ ;
- $\text{R}^4$ , at each occurrence, is selected from OH,  $(\text{CH}_2)_r\text{OR}^2$ , halo,   
25  $\text{C}_{1-4}$  alkyl,  $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$ , and  $(\text{CF}_2)_r\text{CF}_3$ ;
- $\text{R}^{4a}$  is selected from  $\text{C}_{1-4}$  alkyl,  $\text{CF}_3$ ,  $\text{S}(\text{O})_p\text{R}^5$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ , and 1- $\text{CF}_3$ -tetrazol-2-yl;
- 30  $\text{R}^5$ , at each occurrence, is selected from  $\text{CF}_3$ ,  $\text{C}_{1-6}$  alkyl, phenyl, and benzyl;
- X is  $\text{CH}_2$  or  $\text{C}(\text{O})$ ; and,
- 35 Y is selected from pyrrolidino and morpholino.

9. A compound according to Claim 8, wherein;

- D-E is selected from 3-aminophenyl, 3-amidinophenyl, 3-aminomethylphenyl, 3-aminocarbonylphenyl, 3-(methylaminomethyl)phenyl, 3-(1-aminoethyl)phenyl, 3-(2-amino-2-propyl)phenyl, 4-chloro-3-aminophenyl, 4-chloro-3-amidinophenyl, 4-chloro-3-aminomethylphenyl, 4-chloro-3-(methylaminomethyl)phenyl, 4-fluoro-3-aminophenyl, 4-fluoro-3-amidinophenyl, 4-fluoro-3-aminomethylphenyl, 4-fluoro-3-(methylaminomethyl)phenyl, 6-aminopyrid-2-yl, 6-amidinopyrid-2-yl, 6-aminomethylpyrid-2-yl, 6-aminocarbonylpyrid-2-yl, 6-(methylaminomethyl)pyrid-2-yl, 6-(1-aminoethyl)pyrid-2-yl, 6-(2-amino-2-propyl)pyrid-2-yl;
- 15 A is selected from the group: phenyl, 2-pyridyl, 3-pyridyl, 2-pyrimidyl, 2-Cl-phenyl, 3-Cl-phenyl, 2-F-phenyl, 3-F-phenyl, 2-methylphenyl, 2-aminophenyl, and 2-methoxyphenyl; and,
- 20 B is selected from the group: 2-CF<sub>3</sub>-phenyl, 2-(aminosulfonyl)phenyl, 2-(methylaminosulfonyl)phenyl, 2-(dimethylaminosulfonyl)phenyl, 1-pyrrolidinocarbonyl, 2-(methylsulfonyl)phenyl, 4-morpholino, 2-(1'-CF<sub>3</sub>-tetrazol-2-yl)phenyl, 4-morpholinocarbonyl, 2-methyl-1-imidazolyl, 5-methyl-1-imidazolyl, 2-methylsulfonyl-1-imidazolyl and, 5-methyl-1,2,3-triazolyl.

10. A compound according to Claim 9, wherein the compound is of formula IIa.

11. A compound according to Claim 9, wherein the compound is of formula IIb.

35

12. A compound according to Claim 9, wherein the compound is of formula IIc.



13. A compound according to Claim 9, wherein the compound is of formula IIId.

5

14. A compound according to Claim 9, wherein the compound is of formula IIe.

10

15. A compound according to Claim 9, wherein the compound is of formula IIIf.

15

16. A compound according to Claim 3, wherein;

D is selected from  $C(=NR^8)NR^7R^9$ ,  $C(O)NR^7R^8$ ,  $NR^7R^8$ , and  $CH_2NR^7R^8$ , provided that D is substituted meta or para to ring M on E;

20

E is phenyl substituted with R or pyridyl substituted with R;

R is selected from H, Cl, F,  $OR^3$ ,  $CH_3$ ,  $CH_2CH_3$ ,  $OCF_3$ , and  $CF_3$ ;

25 Z is selected from  $C(O)$ ,  $CH_2C(O)$ ,  $C(O)CH_2$ ,  $NHC(O)$ , and  $C(O)NH$ , provided that Z does not form a N-N bond with ring M or group A;

30  $R^{1a}$  and  $R^{1b}$  are independently absent or selected from  $-(CH_2)_r-R^{1'}$ ,  $NCH_2R^{1'}$ ,  $OCH_2R^{1'}$ ,  $SCH_2R^{1'}$ ,  $N(CH_2)_2(CH_2)_tR^{1'}$ ,  $O(CH_2)_2(CH_2)_tR^{1'}$ , and  $S(CH_2)_2(CH_2)_tR^{1'}$ , or combined to form a 5-8 membered saturated, partially saturated or unsaturated ring substituted with 0-2  $R^4$  and which contains from 0-2 heteroatoms selected from the group  
35 consisting of N, O, and S;

$R^{1'}$ , at each occurrence, is selected from H,  $C_{1-3}$  alkyl, halo,  $(CF_2)_rCF_3$ ,  $OR^2$ ,  $NR^2R^{2a}$ ,  $C(O)R^{2c}$ ,  $(CF_2)_rCO_2R^{2c}$ ,  $S(O)_pR^{2b}$ ,

$\text{NR}^2(\text{CH}_2)_r\text{OR}^2$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})_2\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  
 $\text{SO}_2\text{NR}^2\text{R}^{2a}$ , and  $\text{NR}^2\text{SO}_2\text{R}^{2b}$ ;

5 A is selected from one of the following carbocyclic and  
 heterocyclic systems which are substituted with 0-2  $\text{R}^4$ ;  
 phenyl, piperidinyl, piperazinyl, pyridyl,  
 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,  
 pyrrolidinyl, oxazolyl, isoxazolyl, thiazolyl,  
 isothiazolyl, pyrazolyl, and imidazolyl;

10 B is selected from: Y, X-Y,  $\text{NR}^2\text{R}^{2a}$ ,  $\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ , and  
 $\text{NR}^2\text{C}(=\text{NR}^2)\text{NR}^2\text{R}^{2a}$ ;

15 X is selected from  $\text{CH}_2$ ,  $-\text{CR}^2(\text{CR}^2\text{R}^{2b})(\text{CH}_2)_t-$ ,  $-\text{C}(\text{O})-$ ,  $-\text{C}(=\text{NR})-$ ,  
 $-\text{CH}(\text{NR}^2\text{R}^{2a})-$ ,  $-\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{NR}^2\text{C}(\text{O})-$ ,  $-\text{NR}^2\text{C}(\text{O})\text{NR}^2-$ ,  $-\text{NR}^2-$ ,  
 and O;

Y is  $\text{NR}^2\text{R}^{2a}$ , provided that X-Y do not form a N-N or O-N bond;

20 alternatively, Y is selected from one of the following  
 carbocyclic and heterocyclic systems which are  
 substituted with 0-2  $\text{R}^{4a}$ ;  
 phenyl, piperidinyl, piperazinyl, pyridyl,  
 pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl,  
 25 pyrrolidinyl, oxazolyl, isoxazolyl, isoxazoliny,   
 thiazolyl, isothiazolyl, pyrazolyl, imidazolyl,  
 oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl,  
 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl,  
 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl,  
 30 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl,  
 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

35  $\text{R}^4$ , at each occurrence, is selected from =O, OH, Cl, F,  $\text{C}_{1-4}$   
 alkyl,  $(\text{CH}_2)_r\text{NR}^2\text{R}^{2a}$ ,  $(\text{CH}_2)_r\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{NR}^2\text{C}(\text{O})\text{R}^{2b}$ ,  $\text{C}(\text{O})\text{NR}^2\text{R}^{2a}$ ,  
 $\text{CH}(=\text{NH})\text{NH}_2$ ,  $\text{NHC}(=\text{NH})\text{NH}_2$ ,  $\text{SO}_2\text{NR}^2\text{R}^{2a}$ ,  $\text{NR}^2\text{SO}_2\text{-C}_{1-4}$  alkyl,  
 $\text{NR}^2\text{SO}_2\text{R}^5$ ,  $\text{S}(\text{O})\text{R}^5$ , and  $(\text{CF}_2)_r\text{CF}_3$ ;

- 5  $R^{4a}$ , at each occurrence, is selected from =O, OH, Cl, F,  $C_{1-4}$  alkyl,  $(CH_2)_rNR^2R^{2a}$ ,  $(CH_2)_rC(O)R^{2b}$ ,  $NR^2C(O)R^{2b}$ ,  $C(O)NR^2R^{2a}$ ,  $CH(=NH)NH_2$ ,  $NHC(=NH)NH_2$ ,  $SO_2NR^2R^{2a}$ ,  $NR^2SO_2-C_{1-4}$  alkyl,  $NR^2SO_2R^5$ ,  $S(O)_pR^5$ ,  $(CF_2)_rCF_3$ , and 1- $CF_3$ -tetrazol-2-yl;
- 10  $R^5$ , at each occurrence, is selected from  $CF_3$ ,  $C_{1-6}$  alkyl, phenyl substituted with 0-2  $R^6$ , and benzyl substituted with 0-2  $R^6$ ;
- 15  $R^6$ , at each occurrence, is selected from H, =O, OH,  $OR^2$ , Cl, F,  $CH_3$ , CN,  $NO_2$ ,  $(CH_2)_rNR^2R^{2a}$ ,  $(CH_2)_rC(O)R^{2b}$ ,  $NR^2C(O)R^{2b}$ ,  $CH(=NH)NH_2$ ,  $NHC(=NH)NH_2$ , and  $SO_2NR^2R^{2a}$ ;
- 20  $R^7$ , at each occurrence, is selected from H, OH,  $C_{1-6}$  alkyl,  $C_{1-6}$  alkylcarbonyl,  $C_{1-6}$  alkoxy,  $C_{1-4}$  alkoxycarbonyl, benzyl,  $C_{6-10}$  aryloxy,  $C_{6-10}$  aryloxycarbonyl,  $C_{6-10}$  arylmethylcarbonyl,  $C_{1-4}$  alkylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl,  $C_{6-10}$  arylcarbonyloxy  $C_{1-4}$  alkoxycarbonyl,  $C_{1-6}$  alkylaminocarbonyl, phenylaminocarbonyl, and phenyl  $C_{1-4}$  alkoxycarbonyl;
- 25  $R^8$ , at each occurrence, is selected from H,  $C_{1-6}$  alkyl and benzyl; and
- alternatively,  $R^7$  and  $R^8$  combine to form a morpholino group; and,
- 30  $R^9$ , at each occurrence, is selected from H,  $C_{1-6}$  alkyl and benzyl.
- 35 17. A compound according to Claim 16, wherein;
- E is phenyl substituted with R or 2-pyridyl substituted with R;
- R is selected from H, Cl, F,  $OCH_3$ ,  $CH_3$ ,  $OCF_3$ , and  $CF_3$ ;

Z is selected from a  $C(O)CH_2$  and  $C(O)NH$ , provided that Z does not form a N-N bond with group A;

5  $R^{1a}$  is selected from H,  $CH_3$ ,  $CH_2CH_3$ , Cl, F,  $CF_3$ ,  $OCH_3$ ,  $NR^2R^{2a}$ ,  $S(O)_pR^{2b}$ ,  $CH_2S(O)_pR^{2b}$ ,  $CH_2NR^2S(O)_pR^{2b}$ ,  $C(O)R^{2c}$ ,  $CH_2C(O)R^{2c}$ ,  $C(O)NR^2R^{2a}$ , and  $SO_2NR^2R^{2a}$ ;

10  $R^{1b}$  is selected from H,  $CH_3$ ,  $CH_2CH_3$ , Cl, F,  $CF_3$ ,  $OCH_3$ ,  $NR^2R^{2a}$ ,  $S(O)_pR^{2b}$ ,  $CH_2S(O)_pR^{2b}$ ,  $CH_2NR^2S(O)_pR^{2b}$ ,  $C(O)R^{2c}$ ,  $CH_2C(O)R^{2c}$ ,  $C(O)NR^2R^{2a}$ , and  $SO_2NR^2R^{2a}$ ;

15 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2  $R^4$ : phenyl, pyridyl, pyrimidyl, furanyl, thiophenyl, pyrrolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, and imidazolyl;

B is selected from: Y and X-Y;

20 X is selected from  $CH_2$ ,  $-CR^2(CR^2R^{2b})-$ ,  $-C(O)-$ ,  $-C(=NR)-$ ,  $-CH(NR^2R^{2a})-$ ,  $-C(O)NR^2-$ ,  $-NR^2C(O)-$ ,  $-NR^2C(O)NR^2-$ ,  $-NR^2-$ , and O;

25 Y is  $NR^2R^{2a}$ , provided that X-Y do not form a N-N or O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2  $R^{4a}$ ;

30 phenyl, piperidinyl, piperazinyl, pyridyl, pyrimidyl, furanyl, morpholinyl, thiophenyl, pyrrolyl, pyrrolidinyl, oxazolyl, isoxazolyl, isoxazoliny, thiazolyl, isothiazolyl, pyrazolyl, imidazolyl, oxadiazolyl, thiadiazolyl, triazolyl, 1,2,3-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,5-oxadiazolyl, 1,3,4-oxadiazolyl, 35 1,2,3-thiadiazolyl, 1,2,4-thiadiazolyl, 1,2,5-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, 1,2,5-triazolyl, and 1,3,4-triazolyl;

R<sup>2</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

5 R<sup>2a</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, OCH<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

10 R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, OCH<sub>3</sub>, CH<sub>3</sub>, benzyl, and phenyl;

15 alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a 5 or 6 membered saturated, partially unsaturated, or unsaturated ring which contains from 0-1 additional heteroatoms selected from the group consisting of N, O, and S;

R<sup>3</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, and phenyl;

20 R<sup>3a</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, and phenyl;

25 R<sup>4</sup>, at each occurrence, is selected from OH, Cl, F, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, C(O)R<sup>2b</sup>, NR<sup>2</sup>C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, and CF<sub>3</sub>;

30 R<sup>4a</sup>, at each occurrence, is selected from OH, Cl, F, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, C(O)R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>5</sup>, CF<sub>3</sub>, and 1-CF<sub>3</sub>-tetrazol-2-yl;

R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub>, C<sub>1-6</sub> alkyl, phenyl substituted with 0-2 R<sup>6</sup>, and benzyl substituted with 1 R<sup>6</sup>;

35 R<sup>6</sup>, at each occurrence, is selected from H, OH, OCH<sub>3</sub>, Cl, F, CH<sub>3</sub>, CN, NO<sub>2</sub>, NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

5 R<sup>7</sup>, at each occurrence, is selected from H, OH, C<sub>1-3</sub> alkyl, C<sub>1-3</sub> alkylcarbonyl, C<sub>1-3</sub> alkoxy, C<sub>1-4</sub> alkoxy carbonyl, benzyl, phenoxy, phenoxycarbonyl, benzylcarbonyl, C<sub>1-4</sub> alkylcarbonyloxy C<sub>1-4</sub> alkoxy carbonyl, phenylcarbonyloxy C<sub>1-4</sub> alkoxy carbonyl, C<sub>1-6</sub> alkylaminocarbonyl, phenylaminocarbonyl, and phenyl C<sub>1-4</sub> alkoxy carbonyl;

10 R<sup>8</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, and benzyl; and,

alternatively, R<sup>7</sup> and R<sup>8</sup> combine to form a morpholino group;

15 R<sup>9</sup>, at each occurrence, is selected from H, CH<sub>3</sub>, and benzyl.

18. A compound according to Claim 17, wherein;

20 R<sup>1a</sup> is absent or is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>, OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2c</sup>, CH<sub>2</sub>C(O)R<sup>2c</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

25 R<sup>1b</sup> is absent or is selected from H, CH<sub>3</sub>, CH<sub>2</sub>CH<sub>3</sub>, Cl, F, CF<sub>3</sub>, OCH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>2b</sup>, C(O)NR<sup>2</sup>R<sup>2a</sup>, CH<sub>2</sub>S(O)<sub>p</sub>R<sup>2b</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>p</sub>R<sup>2b</sup>, C(O)R<sup>2b</sup>, CH<sub>2</sub>C(O)R<sup>2b</sup>, and SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>;

30 A is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R<sup>4</sup>; phenyl, pyridyl, and pyrimidyl;

35 B is selected from: Y and X-Y;

X is selected from -C(O)- and O;

Y is NR<sup>2</sup>R<sup>2a</sup>, provided that X-Y do not form a O-N bond;

alternatively, Y is selected from one of the following carbocyclic and heterocyclic systems which are substituted with 0-2 R<sup>4a</sup>;

phenyl, piperazinyl, pyridyl, pyrimidyl,  
morpholinyl, pyrrolidinyl, imidazolyl, and 1,2,3-  
triazolyl;

5 R<sup>2</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl,  
and phenyl;

R<sup>2a</sup>, at each occurrence, is selected from H, CF<sub>3</sub>, CH<sub>3</sub>, benzyl,  
and phenyl;

10 R<sup>2b</sup>, at each occurrence, is selected from CF<sub>3</sub>, OCH<sub>3</sub>, CH<sub>3</sub>,  
benzyl, and phenyl;

15 R<sup>2c</sup>, at each occurrence, is selected from CF<sub>3</sub>, OH, OCH<sub>3</sub>, CH<sub>3</sub>,  
benzyl, and phenyl;

alternatively, R<sup>2</sup> and R<sup>2a</sup> combine to form a ring system  
selected from pyrrolidinyl, piperazinyl and morpholino;

20 R<sup>4</sup>, at each occurrence, is selected from Cl, F, CH<sub>3</sub>, NR<sup>2</sup>R<sup>2a</sup>,  
and CF<sub>3</sub>;

R<sup>4a</sup>, at each occurrence, is selected from Cl, F, CH<sub>3</sub>,  
SO<sub>2</sub>NR<sup>2</sup>R<sup>2a</sup>, S(O)<sub>p</sub>R<sup>5</sup>, and CF<sub>3</sub>; and,

25 R<sup>5</sup>, at each occurrence, is selected from CF<sub>3</sub> and CH<sub>3</sub>.

19. A compound according to Claim 1, wherein the  
30 compound is selected from the group:

1-(3-amidinophenyl)-2-[[ (2'-aminosulfonyl-[1,1']-biphen-4-yl)-  
aminocarbonyl]pyrrole;

35 1-(3-amidinophenyl)-2-[[ (2'-tert-butylaminosulfonyl-[1,1']-  
biphen-4-yl)-aminocarbonyl]pyrrole;

1-(3-amidinophenyl)-2-[[ (2'-aminosulfonyl-[1,1']-biphen-4-yl)-  
aminocarbonyl]-4-bromopyrrole;

40

- 1-(3-amidinophenyl)-2-[[5-(2'-aminosulfonylphen-1-yl) pyridin-2-yl]-aminocarbonyl]pyrrole;
- 5 1-benzyl-3-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole;
- 1-benzyl-3-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-(3-amidinophenyl)pyrrole;
- 10 1-(3-amidinophenyl)-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 1-(3-amidinophenyl)-4-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 15 1-(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylamino]pyrazole;
- 25 1-(3-amidinophenyl)-3-methyl-5-(2'-(5''-CF<sub>3</sub>-tetrazolyl)-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-4-chloro-3-methyl-pyrazole;
- 30 1-(3-amidinophenyl)-5-((2'-t-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-4-methoxy-5-((2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)-3-trifluoromethyl-pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-(4'-(imidazol-1-yl-phenyl)aminocarbonyl)pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[(4'-(2''-sulfonylmethyl)phenoxyphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methylcarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 1-(3-amidinophenyl)-5-((2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl)tetrazole;
- 50 1-(3-amidinophenyl)-5-((2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylthio)tetrazole;
- 55 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylsulfoxide]tetrazole;



- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)methylsulfonyl]tetrazole;
- 5 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-aminosulfonylphenyl-1-yl)pyrimidin-2-yl]-aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-2-chloro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-2-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-4'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-chloro-2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 25 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-methyl-2-[[5-(2'-trifluoromethylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(3-chloro-2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)(N'-methyl)aminocarbonyl]pyrazole;
- 45 1-(3-amidinophenyl)-3-n-butyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-n-butyl-5-[[5-(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 50 1-(3-amidinophenyl)-3-n-butyl-5-[[5-(2'-trifluoromethylphenyl-1-yl)pyridin-2-yl]-aminocarbonyl]pyrazole;
- 55

- 1-(3-amidinophenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 5 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-4-methoxy-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-[(4-trifluoromethylphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-4-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-imidazole;
- 15 1-(3-amidinophenyl)-5-[(2'-aminosulfonylphenyl-1-yl)pyridin-2-yl)-aminocarbonyl]-1,2,3-triazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 20 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-1,2,4-triazole;
- 25 3-methyl-1-(3-amidinophenyl)-5-(4'-(4''-chlorophenyl)thiazol-2'-yl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfide-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfoxide-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(2'-trifluoromethylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-(3-amidino)phenyl-3-methyl-5-[4'-(carboxymethyl)phenylaminocarbonyl]pyrazole;
- 40 1-(3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminocarbonyl)phenylaminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[4'-(N,N-dimethylaminosulfonyl)phenylaminocarbonyl]pyrazole;
- 45 1-(3-amidino)phenyl-3-methyl-5-[(4'-tert-butylaminosulfonylphenyl)aminocarbonyl]pyrazole;
- 1-(3-amidino)phenyl-3-methyl-5-[(4'-aminosulfonylphenyl)aminocarbonyl]pyrazole;
- 50 1-(3-amidino)phenyl-3-methyl-5-[(4'-trifluoromethylphenyl)-aminocarbonyl]pyrazole;
- 55 1-(3-amidino)phenyl-3-methyl-5-[(4'-benzylsulfonylpiperidyl)-aminocarbonyl]pyrazole;

- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)-N-methylaminocarbonyl]-3-methyl-pyrazole;
- 5 1-(3-amidinophenyl)-5-[(4'-fluoro-[1,1']-biphen-4-yl)-aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 10 1-(3-cyanophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 15 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole; and,
- 20 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl)-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- and a pharmaceutically acceptable salt.

25

20. A compound according to Claim 1, wherein the compound is selected from the group:

- 30 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl)-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methylpyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-n-butylpyrazole;
- 35 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-n-butylpyrazole;
- 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-n-butylpyrazole;
- 40 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-4-methoxypyrazole;
- 1-(3-amidinophenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 45 1-(3-amidinophenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-3-bromo-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 55

- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl)-[1,1']-biphen-4-yl)methylcarbonyl]pyrazole;
- 5 1-(3-aminocarbonylphenyl)-5-[5-[(2'-aminosulfonylphen-1-yl)pyridin-2-yl]aminocarbonyl]-3-methyl-pyrazole;
- 10 1-(3-amidinophenyl)-5-[[5-(2'-t-butylaminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 15 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyrimidin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 20 1-(3-cyanophenyl)-5-[(4'-(imidazol-1-yl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 25 1-(3-amidinophenyl)-5-[(4'-(morpholin-1-yl)phenyl)-aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-aminocarbonylphenyl)-5-[(4'-(morpholin-1-yl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 30 1-(3-amidinophenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 35 1-(3-aminocarbonylphenyl)-5-[[5-(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-[(4'-(3-methyltetrazol-1-yl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-5-(2'-naphthylaminosulfonyl)-3-methyl-pyrazole;
- 40 1-(3-amidinophenyl)-5-[(4-bromophenyl)aminosulfonyl]-3-methyl-pyrazole;
- 45 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 50 1-(3-amidinophenyl)-3-methyl-5-[(2'-trifluoromethylphenyl)pyrid-2-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(2'-aminosulfonyl-1-yl)pyrimid-5-yl]aminocarbonyl]pyrazole;
- 55 1-(3-amidinophenyl)-3-methyl-5-[(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;

- 1-(3-amidinophenyl)-3-methyl-5-[3-chloro-(2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-2'-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-[5-(2'-fluorophen-1-yl)pyrid-2-yl]aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-tertbutylaminosulfonylphenyl)pyrimid-2-yl]aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[[5-(2'-aminosulfonylphenyl)-[1,6]-dihydropyrimid-2-yl]aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(4-(pyrid-3'-yl)phen-1-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[[2-(2'-pyridyl)ethyl]aminocarbonyl]pyrazole;
- 25 1-(3-amidinophenyl)-3-methyl-5-[(3-phenylpropyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[4-(pyrid-2'-yl)phen-1-ylaminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-methyl-5-[(4-(isopropoxy)phenyl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(5-(2'-trifluoromethylphenyl)-pyrimidin-2-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidinosulfonyl)phenyl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-3-methyl-5-[(4-(piperidinocarbonyl)phenyl)aminocarbonyl]pyrazole;
- 1-(3-amidino-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 45 1-(3-aminocarbonyl-4-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 50 1-methyl-3-(3-amidino)phenyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole; and,
- 1-(3-amidinophenyl)-3-methyl-5-[[4-(pyrazol-4'-yl)phen-1-yl]aminocarbonyl]pyrazole;
- 55

and a pharmaceutically acceptable salt.

21. A compound according to Claim 1, wherein the  
5 compound is selected from the group:

- 1-(3-amidinophenyl)-3-methyl-5-([5-(2'-  
methysulfonylphenyl)pyrid-2-yl]aminocarbonyl)pyrazole;
- 10 1-(3-amidinophenyl)-3-methyl-5-([5-(2'-  
methysulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole;
- 1-(3-cyanophenyl)-3-methyl-5-([5-(2'-  
methysulfonylphenyl)pyrimid-2-  
15 yl]aminocarbonyl)pyrazole,;
- 1-(3-aminocarbonylphenyl)-3-methyl-5-([5-(2'-  
methysulfonylphenyl)pyrimid-2-yl]aminocarbonyl)pyrazole;
- 20 1-(3-(N-aminoamidino)phenyl)-3-methyl-5-[(2'-tert-  
butylaminosulfonyl-[1,1']-biphen-4-  
yl)aminocarbonyl]pyrazole;
- 1-(3-(N-aminoamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-  
25 [1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-(N-methyl-N-hydroxyamidino)phenyl)-3-methyl-5-[(4'-t-  
butylaminosulfonyl-[1,1']-biphen-4-  
yl)aminocarbonyl]pyrazole;
- 30 1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-tert-  
butylaminosulfonyl-[1,1']-biphen-4-  
yl)aminocarbonyl]pyrazole;
- 35 1-(3-(N-methylamidino)phenyl)-3-methyl-5-[(2'-aminosulfonyl-  
[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-  
yl]aminocarbonyl]tetrazole;
- 40 1-(3-aminocarbonylphenyl)-5-([5-(2'-  
aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl)tetrazole;
- 1-(3-amidinophenyl)-5-([5-(2'-trifluoromethylphen-1-  
45 yl)pyridin-2-yl]aminocarbonyl)tetrazole;
- 1-(3-amidinophenyl)-5-[(4'-bromophen-1-yl)  
aminocarbonyl]tetrazole;
- 50 1-(3-aminocarbonylphenyl)-5-([5-(2'-trifluoromethylphen-1-  
yl)pyridin-2-yl]aminocarbonyl)tetrazole;
- 5-(3-amidinophenyl)-1-[(2'-trifluoromethyl-[1,1']-biphen-4-  
55 yl)methyl]tetrazole;

- 1-[(3-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 5 1-[(4-amidinophenyl)methyl]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole
- 1-[(3-amidinophenyl)-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 10 1-[(3-amidinophenyl)-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 1-[(3-amidinophenyl)-5-chloro-4-methyl-2-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 15 5-[(3-amidinophenyl)-2-methyl-4-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 20 1-[(3-amidinophenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole;
- 1-[(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(benzimidazol-1-yl)phen-1-yl)aminocarbonyl]pyrazole;
- 25 1-[(3-amidinophenyl)-3-methyl-5-[(4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 1-[(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 30 1-[(3-amidinophenyl)-3-methyl-5-[[4'-(1,2,4-triazol-2-yl)-phenyl]aminocarbonyl]pyrazole;
- 1-[(3-amidinophenyl)-3-methyl-5-[(4'-cyclohexylphenyl)aminocarbonyl]pyrazole;
- 35 1-[(3-amidinophenyl)-3-methyl-5-[[1,1']-biphen-4-ylaminocarbonyl]pyrazole;
- 40 1-[(3-amidinophenyl)-3-methyl-5-[(4'-morpholinophenyl)aminocarbonyl]pyrazole;
- 1-[(3-amidinophenyl)-3-methyl-5-[(4'-(2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 45 1-[(3-aminomethylphenyl)-3-methyl-5-[(4'-(2-trifluoromethyl)tetrazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 50 1-[(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-dimethylamino)carbonylamino)phen-1'-yl)aminocarbonyl]pyrazole;
- 55 1-[(3-amidinophenyl)-3-methyl-5-[(4'-(N,N-diethylamino)phenyl)aminocarbonyl]pyrazole;

- 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-N,N-diethylamino)phenyl]aminocarbonyl]pyrazole;
- 5 1-(3-amidinophenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole;
- 10 1-(3-aminocarbonylphenyl)-3-methyl-5-[(4'-(1-tetrazolyl)phenyl)aminocarbonyl]pyrazole;
- 15 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-acetylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole;
- 20 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-tert-butyloxycarbonylpiperizin-1-yl)phenyl)aminocarbonyl]pyrazole,;
- 25 1-(3-amidinophenyl)-3-methyl-5-[(4'-piperizin-1-yl-phenyl)aminocarbonyl]pyrazole;
- 30 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-cyclohexylphenyl)aminocarbonyl]pyrazole;
- 35 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholino)-3'-chlorophenyl)aminocarbonyl]pyrazole;
- 40 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 45 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfinyl)pyrazole;
- 50 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole;
- 55 1-(3-aminocarbonylphenyl)-5-[(2'-trifluoromethyl-[1,1']-biphen-4-yl)methyl]tetrazole;
- 1-(3-aminocarbonylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)methyl]tetrazole;
- 1-(3-amidinophenyl)-5-[(4'-cyclopentyloxyphenyl)aminocarbonyl]-3-methyl-pyrazole;
- 45 1-(3-amidinophenyl)-5-[(3-((pyrid-2-yl)methylamino)phenyl)aminocarbonyl]-3-methyl-pyrazole;
- 50 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]pyrazole;
- 55 1-(3-amidinophenyl)-3-trifluoromethyl-5-[(4'-(N-morpholino)-3-chlorophenyl)aminocarbonyl]pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-pyrrolidinocarbonyl)-3'-chlorophenyl)aminocarbonyl]pyrazole;



- 1-(3-amidinophenyl)-3-methyl-5-[(4'-(N-morpholinocarbonyl)-3-chlorophenyl)aminocarbonyl]pyrazole;
- 5 1-(3-cyanophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 10 1-(3-amidinophenyl)-5-[(4'-(N-imidazolyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 15 1-(3-amidinophenyl)-5-[(4'-(N-methyltetrazolon-1-yl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole; and,
- 1-(3'-aminocarbonylphenyl)-5-[(2'-aminosulfonylphenyl-[1,1']-biphen-4-yl)methylcarbonyl]-3-methyl-pyrazole;
- and a pharmaceutically acceptable salt.

20 22. A compound according to Claim 1, wherein the compound is selected from the group:

- 1-(3-amidinophenyl)-5-[4'-(pyrrolidinomethyl)phenyl)aminocarbonyl]-3-methyl-pyrazole;
- 25 1-(3-aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-(2'-aminophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amino-4'-chlorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-(3-amino-4'-fluorophenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amino-4'-methoxyphenyl)-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 40 1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 45 1-(3-amino-4'-chlorophenyl)-5-[(2'-aminosulfonylphenyl)pyridin-2-yl]aminocarbonyl]tetrazole;
- 1-(3-amino-4'-methoxyphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 50 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonylphenyl)pyrid-2-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethyl-4'-methylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 55

- 1-(3-aminomethyl-4'-fluorophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 5 1-(3-aminomethylphenyl)-5-[(4'-(N-pyrrolidino-carbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 10 1-(3-ethylcarboxyamidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-(1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 15 1-(3-(1'-imino-1'-(N-morpholino)methyl)phenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 20 1-[3-[N-((5-methyl-2-oxo-1,3-dioxol-4-yl)methoxycarbonyl)amidino]phenyl]-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(pyrid-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 25 1-(6-bromopyridin-2-yl)-3-methyl-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-(3-amino-4-chlorophenyl)-5-[(2'-aminosulfonyl-3-chloro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 30 1-(3-amino-4-chlorophenyl)-5-[(4'-(1-pyrrolidinocarbonyl)phenyl)aminocarbonyl]tetrazole;
- 35 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]tetrazole;
- 40 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 45 1-(3-aminomethylphenyl)-5-[(2'-methylsulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 1-(3-amidinophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]imidazole;
- 50 1-[3-(methylaminomethyl)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 55 1-[3-(methylaminomethyl)phenyl]-5-[(2'-methylsulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;

1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

5 1-(3-aminomethylphenyl)-5-[(2-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

10 1-(3-aminomethylphenyl)-5-[(3-fluoro-4-(N-pyrrolidinocarbonyl)phenyl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

1-(3-aminomethylphenyl)-5-[(2'-sulfonylmethyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

15 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

20 1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)-[1,6-dihydro]pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

1-(3-aminomethylphenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

25 1-[3-(2'-ethylaminophenyl)-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

30 1-[3-(1-(N-morpholino)imino)phenyl]-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

1-(3-aminomethylphenyl)-5-[2-(2'-aminosulfonyl-[1,1']-biphen-4-yl)-1-hydroxyethyl]-3-trifluoromethyl-pyrazole;

35 1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

40 1-(3-aminomethylphenyl)-5-[(5-(2'-methylsulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

1-[3-amidinophenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;  
and,

45 1-[3-amidinophenyl]-5-[(3-fluoro-2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;

and a pharmaceutically acceptable salt.

50

23. A compound according to Claim 1, wherein the compound is selected from the group:

55 1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)carbonylmethyl]-3-trifluoromethyl-pyrazole;

- 1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonylmethyl)pyrazole;
- 5 1-(3-amidino)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylaminosulfonylmethyl)pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-
- 10 (methylaminosulfonylmethyl)pyrazole;
- 1-(3-(N-carboxymethyl)amidinophenyl)-5-[(5-(2'-aminosulfonylphenyl)pyrimid-2-yl)aminocarbonyl]-3-methyl-
- 15 pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethylphenyl)-5-[(2'-aminosulfonyl-3-methyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 20 1-(3-aminomethylphenyl)-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-1,2,3-triazole;
- 25 1-(3-aminomethyl-4-methyl)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 30 1-(3-aminomethyl-4-chloro)phenyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 1-(3-aminomethyl-4-fluoro)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-
- 35 pyrazole;
- 1-(3-aminomethyl)phenyl-5-[(2'-aminosulfonyl-3-fluoro-[1,1']-biphen-4-yl)aminocarbonyl]-3-methyl-pyrazole;
- 40 1-(3-aminomethyl)phenyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 1-(3-amidinophenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 45 1-(3-aminomethylphenyl)-3-methyl-5-[(3-fluoro-4-(N-morpholino)phenyl)aminocarbonyl]pyrazole;
- 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-[(3-fluoro-4-(2-methylimidazol-1-yl)phenyl)aminocarbonyl]pyrazole;
- 50 1-(3-cyanophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethylpyrazole;
- 55

- 1-(3-amidinophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl]pyrazole;
- 5 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-([1,1']-biphen-4-yl)oxymethyl]pyrazole;
- 1-(3-amidinophenyl)-3-trifluoromethyl-5-((2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole;
- 10 1-(3-carboxamidophenyl)-3-trifluoromethyl-5-((2-fluoro-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole;
- 15 1-(3-aminomethylphenyl)-3-trifluoromethyl-5-((3-trifluoromethyl-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole;
- 1-(3-aminomethylphenyl)-3-ethyl-5-((3-fluoro-2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole;
- 20 1-(3-aminomethylphenyl)-3-ethyl-5-((3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl)pyrazole;
- 25 1-(3-aminomethylphenyl)-3-ethyl-5-((2-fluoro-4-(2-methylsulfonylimidazol-1-yl)phenyl)aminocarbonyl)pyrazole;
- 1-[(6-(aminomethyl)pyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 30 1-[(6-(N-hydroxyamidino)pyrid-2-yl)]-3-methyl-5-[(2'-tert-butylaminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 35 1-[(6-amidinopyrid-2-yl)]-3-methyl-5-[(2'-aminosulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 1-[6-amidinopyrid-2-yl]-3-methyl-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 40 1-(3-aminomethylphenyl)-3-methyl-5-((2-methoxy-4-(N-morpholino)phenyl)aminocarbonyl)pyrazole;
- 45 1-(3-aminomethylphenyl)-3-methyl-5-[4'-(3"-methyl-5"-oxo-3"-pyrazolin-2"-yl)-phenyl]aminocarbonyl]pyrazole;
- 1-[3-(aminomethyl)phenyl]-5-[(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 50 1-(3-aminomethyl-4-fluorophenyl)-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- 55 ethyl 1-[3-(aminomethyl)-phenyl]-5-[3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylate;

- 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-3-carboxylic acid;
- 5 1-[3-(aminomethyl)phenyl]-3-[aminocarbonyl]-5-[3-fluoro-(2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole;
- ethyl 1-[3-(aminomethyl)-phenyl]-3-trifluoromethyl-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]pyrazole-4-carboxylate;
- 10 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylthio)pyrazole;
- 15 1-[3-(aminomethyl)phenyl]-5-[(3-fluoro-2'-methylsulfonyl-[1,1']-biphen-4-yl)aminocarbonyl]-3-(methylsulfonyl)pyrazole;
- 20 1-[3-(aminomethyl)phenyl]-5-[(4-(5-(methoxyaminocarbonyl)imidazol-1-yl)phen-1-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole; and,
- 1-(3-aminomethylphenyl)-5-[(4-(5-methyl-1,2,3-triazol-1-yl)phen-1-yl)aminocarbonyl]-3-trifluoromethyl-pyrazole;
- 25 and pharmaceutically acceptable salts thereof.

24. A pharmaceutical composition, comprising: a

30 pharmaceutically acceptable carrier and a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof.

35 25. A method for treating or preventing a thromboembolic disorder, comprising: administering to a patient in need thereof a therapeutically effective amount of a compound according to Claim 1 or a pharmaceutically acceptable salt thereof.

40

# INTERNATIONAL SEARCH REPORT

Internat'l Application No  
PCT/US 97/22895

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D207/34 C07D231/14 C07D233/90 C07D249/06 C07D257/04  
A61K31/40 A61K31/41 C07D401/12 C07D403/12 C07D409/12  
C07D413/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	EP 0 418 845 A (FUJISAWA PHARMACEUTICAL CO., LTD.) 27 March 1991 see page 2, line 3; claims 1-11 ---	1-25
A	PATENT ABSTRACTS OF JAPAN vol. 017, no. 025, 18 January 1993 & JP 04 247081 A (TAKEDA CHEMICAL INDUSTRIES LTD.), 3 September 1992, see abstract ---	1-25
A	PATENT ABSTRACTS OF JAPAN vol. 096, no. 010, 31 October 1996 & JP 08 143565 A (FUJISAWA PHARMACEUTICAL CO., LTD.), 4 June 1996, see abstract ---	1-25
-/--		

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

31 March 1998

Date of mailing of the international search report

24. 04. 98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Herz, C

## INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 97/22895

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	PATENT ABSTRACTS OF JAPAN vol. 015, no. 177, 7 May 1991 & JP 03 041072 A (HOKKO CHEM. IND. CO., LTD.), 21 February 1991, see abstract ---	1-25
A	PATENT ABSTRACTS OF JAPAN vol. 014, no. 356, 2 August 1990 & JP 02 129171 A (NISSAN CHEM. IND., LTD.), 17 May 1990, see abstract ---	1-25
A	DE 36 33 840 A (HOECHST AG) 14 April 1988 see claims 1-10 ---	1-25
A	EP 0 554 829 A (FUJISAWA PHARMACEUTICAL CO., LTD.) 11 August 1993 see page 3, line 3; claims 1-12 ---	1-25
A	US 4 540 703 A (M. UCHIDA ET AL.) 10 September 1985 see claims 1-52 -----	1-25



# INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 97/ 22895

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
  
2. ☒ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:  
See further information sheet PCT/ISA/210
  
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
  
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
  
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
  
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/ US 97/22895

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Claims Nos.: 1(part)-18(part),24(part),25(part)

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

The breadth of the claims is so large and encompasses too broad a range of totally different chemical groups. the vast number of theoretically conceivable compounds resulting from the combination of all claimed substituents precludes a comprehensive search. Guided by the inventive concept as disclosed in the descriptive part of the present application a complete search has been limited to claims 19 to 23. Claims 1 to 18 and 14 to 23 have been only searched as far as specific examples disclosed in the application are concerned (cf. Articles 6, 15 and Rule 33 PCT, Guidelines Exam. Part B, Chapt. III, 3.6, 3.7).

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No.

PCT/US 97/22895

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 418845 A	27-03-91	AT 126216 T	15-08-95
		AU 637142 B	20-05-93
		AU 6307290 A	18-04-91
		CA 2025599 A	23-03-91
		CN 1050382 A	03-04-91
		DE 69021472 D	14-09-95
		DE 69021472 T	25-01-96
		ES 2088933 T	01-10-96
		HU 9500344 A	28-09-95
		IE 68857 B	24-07-96
		IL 95675 A	31-03-96
		JP 2586713 B	05-03-97
		JP 3141261 A	17-06-91
		NO 301006 B	01-09-97
		PT 95389 A	22-05-91
		RU 2021990 C	30-10-94
		RU 2059622 C	10-05-96
		US 5134142 A	28-07-92
-----			
DE 3633840 A	14-04-88	AU 610085 B	16-05-91
		AU 7930887 A	14-04-88
		DK 518287 A	05-04-88
		EP 0269806 A	08-06-88
		JP 63091373 A	22-04-88
		PH 25550 A	08-08-91
		US 4891057 A	02-01-90
		US 5082949 A	21-01-92
-----			
EP 554829 A	11-08-93	AU 663149 B	28-09-95
		AU 3217493 A	12-08-93
		CA 2088835 A	06-08-93
		CN 1075959 A	08-09-93
		HU 9500347 A	28-09-95
		IL 104311 A	13-07-97
		JP 5246997 A	24-09-93
		MX 9300579 A	30-09-93
		US 5550147 A	27-08-96
		US 5670533 A	23-09-97
		ZA 9300077 A	04-08-93
-----			

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 97/22895

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4540703 A	10-09-85	JP 1469540 C	14-12-88
		JP 55036447 A	14-03-80
		JP 63018588 B	19-04-88
		US 4663323 A	05-05-87
		US 4766120 A	23-08-88
-----			